Dysphagia rates after minimally invasive tubular approach to the anterior cervical spine

Tasas de disfagia posoperatoria en el abordaje tubular mínimamente invasivo de la columna cervical anterior

José A. I. Romero-Rangel^{1,2*}, Sergio Soriano-Solís³, Manuel Rodríguez-García³, Manuel E. Soto-García¹, Héctor Soriano-Solís³, Briscia Y. Aranda-Torres^{1,4} y José A. Soriano-Sánchez⁵

¹Centro Neurológico, American British Cowdray Medical Center IAP, Ciudad de México; ²Departamento de Enseñanza, Universidad de Sonora, Sonora; ³Cirugía Ortopédica, Centro Neurológico, American British Cowdray Medical Center, IAP, Ciudad de México; ⁴Facultad de Ciencias Nutricionales, Universidad de Sonora (UNISON), Sonora; ⁵Clínica de Cirugía de Columna Vertebral, Centro Neurológico, American British Cowdray Medical Center, IAP, Ciudad de México. México

Abstract

Introduction: Cervical spine surgery (anterior cervical spine surgery [ACSS]) is a commonly indicated to provide neurological decompression, correct deformity, and preserve stability. Current series report good to excellent results, but a number of patients suffer from dysphagia with reported rates as high as 80%. **Materials and methods:** Retrospective cohort of patients undergoing ACSS (anterior cervical discectomy and fusion [ACDF]; and arthroplasty, total disc replacement [TDR]) from 2013 to 2017 by a single surgeon. We collected demographic data, surgical outcomes, clinical outcomes, and dysphagia incidence and magnitude (visual analog scale [VAS]), we analyzed results with SPSS 22. **Results:** A total of 81 patients operated from 2013 to 2018, ACDF was performed in 70, and TDR was performed in 11, mean surgical time was 156.8, mean neck pain and radicular pain VAS reduced significantly in both groups ($p < 0.01$). Dysphagia was encountered in 18 ACDF patients, and one TDR patient. Overall dysphagia rate was 9.1% in day-1 post-operative, with mean 0.27 magnitude (standard deviation [SD] 0.4) in ACDF group and 0.18 (0.6 SD) in TDR group, 1-month follow-up dysphagia rate was 0%. **Conclusions:** Minimally invasive techniques may reduce post-operative dysphagia related to soft-tissue trauma, to 9% in day-1 post-operative (as compared to 39%), and to 0% (as compared to 8.5% reported worldwide) by 1-month follow-up.

Key Words: Anterior cervical. Tubular. Minimally invasive. Dysphagia. Spine surgery. Anterior cervical spine surgery.

Resumen

Antecedentes: La cirugía cervical anterior (CCA) esta comúnmente indicada para proporcionar descompresión neurológica, corregir deformidades y preservar la estabilidad. Las series actuales reportan tasas de disfagia de hasta el 80%. **Método:** cohorte retrospectiva de pacientes sometidos a CCA (discectomía y fusión cervical anterior [DFCA], y artroplastia total de disco [ATD]) de 2013 a 2017. Se recopilaron datos demográficos, quirúrgicos, clínicos y de incidencia y magnitud de la disfagia (escala visual analógica [EVA]). Se analizaron los resultados con SPSS 22. **Resultados:** 81 pacientes operados de 2013 a 2018, ACDF en 70 y ATD en 11, tiempo quirúrgico promedio de 156.8 minutos. La puntuación en la EVA cervical y radicular

Correspondencia:

*José A. I. Romero-Rangel

Avda. Carlos Fernández Graef, 154

Santa Fe, Col. Contadero

C.P. 05330, Ciudad de México, México

E-mail: gmaisrael@gmail.com

Fecha de recepción: 23-04-2019

Fecha de aceptación: 27-08-2019

DOI: 10.24875/CIRU.19001257

Cir Cir. 2020;88(3):286-291

Contents available at PubMed

www.cirugiaycirujanos.com

0009-7411/© 2019 Academia Mexicana de Cirugía. Publicado por Permayer. Este es un artículo open access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

se redujo significativamente en ambos grupos ($p < 0.01$). Hubo disfagia en 18 pacientes con DFCA y en uno con ATD. La tasa de disfagia fue del 9.1% en el primer día de posoperatorio, con EVA de 0.27 (desviación estándar [DE]: 0.4) en el grupo DFCA y de 0.18 (DE: 0.6) en el grupo ATD. La tasa de disfagia de seguimiento a 1 mes fue del 0%. **Conclusiones:** Las técnicas mínimamente invasivas reducen la disfagia posoperatoria en CCA al 9% en el primer día (en comparación con el 39%) y al 0% (en comparación con el 8.5% reportado mundialmente) al mes de seguimiento.

Palabras Clave: Cervical anterior. Tubular. Mínimamente invasiva. Disfagia. Cirugía de columna. Cirugía cervical anterior.

Introduction

Anterior cervical spine surgery (ACSS) is a commonly indicated procedure to solve a variety of cervical spine pathologies (vertebral body fractures, tumors, and degenerative disk disease)^{1,2}, at present, there exist a wide spectrum of surgical techniques able to be done through an anterior approach, whose main target is to provide neurological decompression of the neural axis, correct deformity, and preserve or attain mechanical stability³. Current series report good to excellent results on pain and neurological deficit improvement with high patient satisfaction scores⁴ irrespective of the chosen surgical technique, never the less, early in the post-operative period, and even in the long-term follow-up, a number of patients suffer from dysphagia, because significant retraction may be exerted on the esophagus during the surgical procedure or related to adverse outcomes from surgery such as post-operative hematoma, recurrent laryngeal nerve palsy, esophageal injury, post-operative adhesions, and device migration^{1,3,5}. Dysphagia, being a well-known complication of ACSS^{5,6}, can interfere with daily and social life, affecting prognosis adversely because of dehydration, malnutrition, or ineffective protection of airway from aspiration, ultimately becoming a life-threatening condition in severe cases^{1,2,7}.

Dysphagia is defined erratically as a subjective increase in the time or effort required to move food from mouth to the stomach or more practically as difficulty in swallowing solids and liquids, and as a matter of inconsistency in diagnostic criteria across the scientific literature, its ranges vary widely from 1% to 80%^{1,5,8}. In prospective studies, rates of dysphagia 1-month post-operatively are reported to be as high as 50%, which might decrease progressively overtime to 10-12.5% at 1 year⁹⁻¹¹. Risk factors to the development of dysphagia include female gender, age over 60 years, smoking, preexisting swallowing dysfunction, C4-C5 and C5-C6 involvement, and multilevel surgery^{9,12,13}.

Most anterior cervical surgery trials are focused in outcomes related to fusion rates and pain relief as

well as neurologic deficit recovery, but few researches have been undertaken to determine dysphagia rates consistently as well as in the measures to prevent its post-operative incidence⁹. There is some evidence about steroids use, graft type, low profile plates, and surgical approach regarding dysphagia rates with no guidelines established up to date^{3,8,14-20}. In this paper, we sought to describe our experience with a minimally invasive tubular approach for the anterior cervical procedures such as anterior cervical discectomy and fusion (ACDF) and total disc replacement (TDR) (arthroplasty), with the aim to compare our rates of dysphagia with those of available literature.

Materials and methods

The present work represents a retrospective observational cohort study with historical cohort comparison. We review the clinical files of patients undertaking a minimally invasive tubular approach to the anterior cervical spine to perform an ACDF or arthroplasty from 2013 to 2017. We collected data from clinical files and divided them by type of surgery, we then review variables such as sex, age, body mass index (BMI), surgical procedure, surgical time, bleeding, number of operated levels, radicular and neck pain visual analog scale (VAS), dysphagia VAS at post-operative day 1, 1 month, and end of follow-up. We processed data with SSPS 22 software, performing demographic statistical analysis and comparing incidence rates of radicular and neck pain VAS, dysphagia VAS from pre-operative to 1st-day post-operative, 1 month, and end of follow-up. We then compared dysphagia rates for the type of surgical procedure.

Results

We collected 81 patients operated by a minimally invasive approach from 2013 to 2017 by the senior author (JASS), 70 patients underwent surgery for

Figure 1. Patient distribution by operated levels and gender.

Figure 2. Immediate day 1 post-operative visual analog scale dysphagia distribution (0 means no dysphagia).

ACDF (44 females and 26 males) and 11 patients (8 females and three males) or arthroplasty.

Mean age in ACDF group was 52.5 years (11.9 standard deviation [SD]), mean BMI was 24.8 (3.7 SD), mean operated levels per patient were 2.01 (0.8 SD), mean surgical time was 156.8 min (SD 67.2), and mean surgical bleeding was 39.07 ml (SD 31.03). When analyzed by the number of levels, surgical time resulted to be 55.5 min (20.0 SD) per level, and bleeding was 23.23 ml per level (22.5 SD). Mean pre-operative radicular pain VAS was reported to be 6.77 (1.2 SD) as compared to 1.35 (1.1 SD) mean VAS at the end of follow-up ($p < 0.01$). Mean pre-operative neck pain was reported to be 4.16 (2.7 SD) as compared to 0.67 (0.9 SD) mean VAS at the end

Figure 3. Significant changes ($p < 0.01$) in mean dysphagia rates at day 1 post-operative and 1-month follow-up.

of follow-up ($p < 0.01$). Mean post-operative VAS dysphagia on day 1 post-operative was 0.27 (0.4 SD) with a minimum of 0 and a maximum of 2, and no dysphagia was encountered at 1-month post-operative ($p < 0.01$). The mean follow-up period was 32.7 months (13.0 SD) with a minimum of 9 months and a maximum of 60 months. Figure 1 shows patients distribution by operated level. Figure 2 demonstrates immediate day 1 post-operative VAS dysphagia distribution (0 means no dysphagia). Figure 3 shows significant changes ($p < 0.01$) in mean dysphagia rates at day 1 post-operative and 1-month follow-up.

Mean age in arthroplasty group was 42.7 years (13.0 SD), mean BMI was 24.44 (3.0 SD), mean operated levels per patient were 1.36 (0.6 SD), mean surgical time was 125.45 min (SD 34.7), and mean surgical bleeding was 30.64 ml (SD 14.5). When analyzed by the number of levels, surgical time resulted to be 54.77 min per level (16.0 SD), and bleeding was 23.74 ml per level (9.2). Mean pre-operative radicular pain VAS was reported to be 7.0 (1.1 SD) as compared to 0.64 (0.8 SD) mean VAS at the end of follow-up ($p < 0.01$). Mean pre-operative neck pain was reported to be 3.09 (2.5 SD) as compared to 0.64 (0.8 SD) mean VAS at the end of follow-up ($p < 0.01$). Mean post-operative VAS dysphagia on day 1 post-operative was 0.18 (0.6 SD) with a minimum of 0 and a maximum of 2, and no dysphagia was encountered at 1-month post-operative with no statistically significant difference ($p = 0.341$) due to the low sample size. The mean follow-up period was 26.7 months (6.7 SD) with a minimum of 18 months and a

Table 1. Dysphagia rates reported in current literature as compared to the present series (Modified from Shriver 2017). FU = Follow-Up

Author	Day-1 (95% CI)	End of FU Rates	FU Months
Soriano 2018	23.5 % (Mean) 9.1% (Arthroplasty) (Since 1-Month) 25.7% (ACDF)	0%	31.85
World Global Rate		8.5%	46.7
Graham 2014		8.1%	6
Burkus 2014		9.8%	84
Hou 2013		9.7%	22.5
Kepler 2012		39%	1.5
Coric 2011		6.1%	24
Cheng 2011		16.7%	36
McAfee 2010		8%	24
Anderson 2008		7.2%	24
Fernández-Fairen 2008		6.1%	24
Xie 2007		13.3%	24
Song 2006		2.6%	24
Bazaz 2002		20.2%	6
Bruneau 2001		1.9%	24.6
Bolesta 200		6.7%	42

Figure 4. Distribution by operated levels and gender.

maximum of 36 months. Figure 4 shows patients distribution by operated levels. Figure 5 demonstrates comparative day-1 post-operative and 1-month VAS dysphagia distribution between ACDF and arthroplasty. Table 1 shows dysphagia rates reported in current literature as compared to the present series (Modified from Shriver).

Figure 5. Comparative day-1 post-operative and 1-month visual analog scale dysphagia distribution between anterior cervical discectomy and fusion and total disc replacement.

Discussion

Current research on dysphagia after ACSS has focused on surgical approach (ACDF vs. TDR, vs. anterior cervical corpectomy and fusion vs. ZERO-P)²¹⁻²⁶, steroid use (intraoperative vs. post-operative, local or systemic)²⁷, and graft implications (bone morphogenetic proteins two use vs. autografts vs. allograft)^{18,28}, as well as overall incidence²⁸, but few researches have been taken over the impact of minimally invasive approaches to reduce incidence rates.

Minimally invasive approaches have the common aim of avoiding biomechanical complications and are associated with decreased pain and morbidity in the early post-operative period, as well as with faster recovery rates accounting for the increasing number of safe outpatient-procedures being performed nowadays even in the cervical spine^{29,30}. The present study was conceived based on the concept that minimally invasive approaches reduce soft-tissue trauma³¹, and so for reduce post-operative complications, such as dysphagia, a well-known complication related to soft-tissue edema (among other causes) after an ACSS⁶. Theoretically, a lesser rate of dysphagia in the early post-operative period would provide a proof of concept that minimally invasive tubular approaches, in fact, reduce soft-tissue trauma and their related complications. Worldwide rates of dysphagia are reported to be around 8.5 (5.7-11.3%, 95% confidence interval) as reported by Shriver et al. in their meta-analysis of 14 papers evaluating post-operative dysphagia after ACDF²⁸, among them, the greatest rate of dysphagia (39%) was reported by Kepler on a cohort study with

mean follow-up of 1.5 months¹, becoming also the paper with the shortest follow-up, which translates in a high dysphagia rate in the early post-operative period, providing evidence of the important role of soft-tissue trauma in the genesis of dysphagia, a fact that can help understand why steroids have proved effective in reducing early post-operative dysphagia³²⁻³⁴ driven by their known anti-inflammatory effects. Conversely, our present series results report a 9% rate of dysphagia by 1-month follow-up, presenting a 9% dysphagia rate in day-1 post-operative that fully recovered by the end of follow-up.

Conclusions

Based on this result, we can assume that minimally invasive techniques may provide an effective and safer surgical approach to the anterior cervical spine to reduce post-operative dysphagia related to soft-tissue trauma, as stated by the present series results effectively reducing early post-operative dysphagia to 9% in day-1 post-operative (as compared to 39%) and to 0% (as compared to 8.5% reported worldwide, 46.7 months mean follow-up) by 1-month follow-up. Further research with a level 1 evidence randomized clinical trial may help underscore the importance of adopting minimally invasive techniques in the armamentarium of spine surgeons to achieve better results in the anterior cervical spine, a surgical procedure that is well known by their excellent results, that can further been improved by reducing soft-tissue trauma-related complications that have potentially life-threatening outcomes such as dysphagia, and by the same time increasing patient satisfaction by a faster recovery rate and reducing hospital stay.

Acknowledgment

The authors would like to thank our families and friends for their time given and their support over all these years.

Conflicts of interest

The present article is a systematic review and so for an ethical approval was not required.

Funding

The present work had no other funding than the authors own.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that they have followed the protocols of their work center on the publication of patient data.

Right to privacy and informed consent. The authors declare that no patient data appear in this article.

References

1. Kepler CK, Rihn JA, Bennett JD, Anderson DG, Vaccaro AR, Albert TJ, et al. Dysphagia and soft-tissue swelling after anterior cervical surgery: a radiographic analysis. *Spine J.* 2012;12:639-44.
2. Kang SH, Kim DK, Seo KM, Kim KT, Kim YB. Multi-level spinal fusion and postoperative prevertebral thickness increase the risk of dysphagia after anterior cervical spine surgery. *J Clin Neurosci.* 2011; 18:1369-73.
3. Njoku I Jr, Alimi M, Leng LZ, Shin BJ, James AR, Bhargoo S, et al. Anterior cervical discectomy and fusion with a zero-profile integrated plate and spacer device: a clinical and radiological study: clinical article. *J Neurosurg Spine* 2014;21:529-37.
4. Kim Y, Boachie-Adjei O, Bridwell K, Lenke L, Kim Y, Cho S, et al. Selection of proximal fusion level in adult lumbar segmental instrumentation fusion to L5 or S1 for degenerative lumbar scoliosis. *Spine J.* 2009;9:96S-7S.
5. Kang SS, Lee JS, Shin JK, Lee JM, Youn BH. The association between psychiatric factors and the development of chronic dysphagia after anterior cervical spine surgery. *Eur Spine J.* 2014;23:1694-8.
6. Min Y, Kim WS, Kang SS, Choi JM, Yeom JS, Paik NJ. Incidence of dysphagia and serial videofluoroscopic swallow study findings after anterior cervical discectomy and fusion: a prospective study. *Clin Spine Surg.* 2016;29:E177-81.
7. Nanda A, Sharma M, Sonig A, Ambekar S, Bollam P. Surgical complications of anterior cervical discectomy and fusion for cervical degenerative disk disease: a single surgeon's experience of 1,576 patients. *World Neurosurg.* 2014;82:1380-7.
8. Lovasik BP, Holland CM, Howard BM, Baum GR, Rodts GE, Refai D. Anterior cervical discectomy and fusion: comparison of fusion, dysphagia, and complication rates between recombinant human bone morphogenetic protein-2 and beta-tricalcium phosphate. *World Neurosurg.* 2017; 97:674-830.
9. Koreckij TD, Davidson AA, Baker KC, Park DK. Retropharyngeal steroids and dysphagia following multilevel anterior cervical surgery. *Spine (Phila Pa 1976).* 2016;41:E530-4.
10. Papavero L, Heese O, Klotz-Regener V, Buchalla R, Schröder F, Westphal M. The impact of esophagus retraction on early dysphagia after anterior cervical surgery: Does a correlation exist? *Spine (Phila Pa 1976).* 2007;32:1089-93.
11. Radcliff KE, Bennett J, Stewart RJ, Kepler CK, Sidhu GS, Hilibrand AS, et al. Change in angular alignment is associated with early dysphagia after anterior cervical discectomy and fusion. *Clin Spine Surg.* 2016; 29:248-54.
12. Olsson EC, Jobson M, Lim MR. Risk factors for persistent dysphagia after anterior cervical spine surgery. *Orthopedics.* 2015;38:e319-23.
13. Shi S, Li XF, Zhao QT, Yang LL, Liu ZD, Yuan W. Risk factors for dysphagia after single-level anterior cervical decompression with arthroplasty or fusion: a prospective study comparing 2 zero-profile implants. *World Neurosurg.* 2016;95:148-55.
14. Fisahn C, Schmidt C, Rustagi T, Moisi M, Iwanaga J, Norvell DC, et al. Comparison of chronic dysphagia in standalone versus conventional plate and cage fusion. *World Neurosurg.* 2018;109:e382-8.
15. Li Y, Hao D, He B, Wang X, Yan L. The efficiency of zero-profile implant in anterior cervical discectomy fusion: a prospective controlled long-term follow-up study. *J Spinal Disord Tech.* 2015;28:398-403.
16. Lu DC, Tumialán LM, Chou D. Multilevel anterior cervical discectomy and fusion with and without rhBMP-2: a comparison of dysphagia rates and outcomes in 150 patients. *J Neurosurg Spine.* 2013;18:43-9.
17. Miao J, Shen Y, Kuang Y, Yang L, Wang X, Chen Y, et al. Early follow-up outcomes of a new zero-profile implant used in anterior cervical discectomy and fusion. *J Spinal Disord Tech.* 2013;26:E193-7.

18. Riederman BD, Butler BA, Lawton CD, Rosenthal BD, Balderama ES, Bernstein AJ. Recombinant human bone morphogenetic protein-2 versus iliac crest bone graft in anterior cervical discectomy and fusion: dysphagia and dysphonia rates in the early postoperative period with review of the literature. *J Clin Neurosci*. 2017;44:180-3.
19. Segebarth B, Datta JC, Darden B, Janssen ME, Murrey DB, Rhyné A, et al. Incidence of dysphagia comparing cervical arthroplasty and ACDF. *SAS J*. 2010;4:3-8.
20. Song KJ, Lee SK, Ko JH, Yoo MJ, Kim DY, Lee KB. The clinical efficacy of short-term steroid treatment in multilevel anterior cervical arthrodesis. *Spine J*. 2014;14:2954-8.
21. Anderson PA, Nassr A, Currier BL, Sebastian AS, Arnold PM, Fehlings MG, et al. Evaluation of adverse events in total disc replacement: a meta-analysis of FDA summary of safety and effectiveness data. *Global Spine J*. 2017;7:76S-83S.
22. Zhong ZM, Li M, Han ZM, Zeng JH, Zhu SY, Wu Q, et al. Does cervical disc arthroplasty have lower incidence of dysphagia than anterior cervical discectomy and fusion? A meta-analysis. *Clin Neurol Neurosurg*. 2016;146:45-51.
23. Fallah A, Akl EA, Ebrahim S, Ibrahim GM, Mansouri A, Foote CJ, et al. Anterior cervical discectomy with arthroplasty versus arthrodesis for single-level cervical spondylosis: a systematic review and meta-analysis. *PLoS One*. 2012;7:e43407.
24. Wang T, Wang H, Liu S, An HD, Liu H, Ding WY. Anterior cervical discectomy and fusion versus anterior cervical corpectomy and fusion in multilevel cervical spondylotic myelopathy: a meta-analysis. *Medicine (Baltimore)*. 2016;95:e5437.
25. Shao H, Chen J, Ru B, Yan F, Zhang J, Xu S, et al. Zero-profile implant versus conventional cage-plate implant in anterior cervical discectomy and fusion for the treatment of degenerative cervical spondylosis: a meta-analysis. *J Orthop Surg Res*. 2015;10:148.
26. Dong J, Lu M, Lu T, Liang B, Xu J, Zhou J, et al. Meta-analysis comparing zero-profile spacer and anterior plate in anterior cervical fusion. *PLoS One*. 2015;10:e0130223.
27. Zadegan SA, Jazayeri SB, Abedi A, Bonaki HN, Vaccaro AR, Rahimi-Movaghgar V. Corticosteroid administration to prevent complications of anterior cervical spine fusion: a systematic review. *Global Spine J*. 2018; 8:286-302.
28. Shriver MF, Lewis DJ, Kshetry VR, Rosenbaum BP, Benzel EC, Mroz TE. Dysphagia rates after anterior cervical discectomy and fusion: a systematic review and meta-analysis. *Global Spine J*. 2017; 7:95-103.
29. Banczerowski P, Czigléczi G, Papp Z, Veres R, Rappaport HZ, Vajda J. Minimally invasive spine surgery: systematic review. *Neurosurg Rev*. 2015;38:11-26.
30. Ban D, Liu Y, Cao T, Feng S. Safety of outpatient anterior cervical discectomy and fusion: a systematic review and meta-analysis. *Eur J Med Res*. 2016;21:34.
31. Pishnamaz M, Schemmann U, Herren C, Horst K, Lichte P, Hildebrand F, et al. Muscular changes after minimally invasive versus open spinal stabilization of thoracolumbar fractures: a literature review. *J Musculoskelet Neuronal Interact*. 2018;18:62-70.
32. Siasios I, Fountas K, Dimopoulos V, Pollina J. The role of steroid administration in the management of dysphagia in anterior cervical procedures. *Neurosurg Rev*. 2018;41:47-53.
33. Nam TW, Lee DH, Shin JK, Goh TS, Lee JS. Effect of intravenous dexamethasone on prevertebral soft tissue swelling after anterior cervical discectomy and fusion. *Acta Orthop Belg*. 2013;79:211-5.
34. Adenikinju AS, Halani SH, Rindler RS, Gary MF, Michael KW, Ahmad FU. Effect of perioperative steroids on dysphagia after anterior cervical spine surgery: a systematic review. *Int J Spine Surg*. 2017;11:9.