




ÓRGANO DE DIFUSIÓN
CIENTÍFICA DE LA
ACADEMIA MEXICANA
DE CIRUGÍA

FUNDADA EN 1933



PERMANYER MÉXICO
www.permayer.com



ISSN: 2444-0507

CIRUGÍA Y CIRUJANOS

Content

Original articles

- 571 Basic and combined peripheral biomarkers have no predictive and prognostic value in breast cancer patients undergoing pre-operative systemic therapy**
Cem Karaali, Huseyin Esin, Emel E. Pala, Murat Keser, Olçun U. Unal, Orhan Ureyen, and Mustafa Emiroğlu
- 582 Pelvic index/mesorectal length ratio: a new predictive factor for rectal cancer prognosis and low anterior resection syndrome**
Muhammed Gurluk, Onur O. Karagulle, Ekrem Cakar, Betül Avcı, Ensar Cakir, Ibrahim T. Rakici, and Mert M. Sevinc
- 591 Association of NRF-2 changes in plasma and pericardial fluid with renal injury in patients undergoing cardiac surgery**
Mahmut Padak, Reşat Dikme, Ebru Temiz, Ömer Göç, Ezhar Ersöz, and Yasemin Hacıanlı
- 600 The effect of genetic amniocentesis on Doppler measurements of utero-placental and fetoplacental circulations**
Nizamettin Bozbay, Ahmet Yanar, Ayşe C. Duymuş, Yasin Altekin, Aybike Tazegül-Pekin, and Gökçen Örgül
- 605 Concordance of CO₂ gaps measured using peripheral and central venous blood in patients diagnosed with septic shock**
Erdem Yalcinkaya, Umut Sabri Kasapoglu, Huseyin Arikan, Nazım Onur Sahin, Betül Yıldız, sra Tekin, Aybuke Selcuk, Emel Eryuksel, and Sait Karakurt
- 612 Perirenal fat Hounsfield units: not only for renal stones but also for renal cell carcinoma**
Serkan Akan, Caner Ediz, Esin Derin-Çiçek, Ozgun Ete, Yunus E. Kızılkın, Aytac Sahin, and Ayhan Verit
- 620 Neoadjuvant chemo(radio)therapy and anastomotic leakage risk in patients with esophageal squamous cell carcinoma**
Yuzhen Xie, Hongni Qin, Fan Wu, Yong Zhang, and Yang Yang
- 628 Evaluation of mediastinal lymphadenopathy from long-term radiological findings in COVID-19**
Gulhan Ayhan-Albayrak and Mustafa I. Bardakci
- 366 An evaluation of factors which affect recurrence and malignancy of sinonasal papillomas**
Zülküf Küçüktağ, Bülent Öcal, Cem Saka, Esmâ Altan, Mehmet H. Korkmaz, and Emel Çadallı-Tatar
- 644 Cellular indices and acute phase reactants as predictors of abscess after appendectomy**
Júlio C. Moreno-Alfonso, Laura C. Lozano-Eslava, Sharom Barbosa-Velásquez, Ada Molina-Caballero, M. Concepción Yáñez-Irazábal, and Alberto Pérez-Martínez

Volume 93, No. 6, November-December 2025

Journal Citation Reports™ from Clarivate, 2024, Medline/PubMed, EBSCO Essentials™

Basic and combined peripheral biomarkers have no predictive and prognostic value in breast cancer patients undergoing pre-operative systemic therapy

Los biomarcadores periféricos básicos y combinados no tienen valor predictivo ni pronóstico en pacientes con cáncer de mama sometidas a terapia sistémica preoperatoria

Cem Karaali^{1*}, Huseyin Esir², Emel E. Pala³, Murat Keser⁴, Olçun U. Unal⁴, Orhan Ureyen¹, and Mustafa Emiroğlu⁵

¹Department of General Surgery, University of Health Sciences, Izmir Faculty of Medicine, Izmir City Hospital; ²Department of General Surgery, Izmir City Hospital; ³Department of Pathology, University of Health Sciences, Izmir Faculty of Medicine, Izmir City Hospital; ⁴Department of Medical Oncology, University of Health Sciences, Izmir City Hospital; ⁵Department of General Surgery, Memorial Bodrum Hospital, Mugla, Izmir, Turkey

Abstract

Objective: The main goal in breast cancer patients receiving neoadjuvant chemotherapy (NAC) is to achieve a pathologic complete response (pCR). Our study examines whether neutrophil/lymphocyte ratio (NLR), platelet/lymphocyte ratio (PLR), monocyte/lymphocyte ratio (MLR), NLR/PLR, systemic immune-inflammation index (SII), systemic inflammation response index (SIRI), and pan-immune-inflammation-value (PIV) indices can be used to determine pCR and prognosis. **Methods:** The study included 228 patients who received NAC for breast cancer between 2010 and 2021. **Results:** Fifty-nine (25.9%) patients had pCR to NAC. In univariate analysis, a significant difference was found between clinical node status, estrogen receptor, progesterone receptor, human epidermal growth factor receptor 2 (HER2), breast cancer molecular subtypes, and pCR ($p < 0.05$). However, in multivariate analysis, only breast cancer molecular subtypes and HER2 status were significantly associated with pCR ($p < 0.05$). **Conclusions:** Our study did not find a significant relationship between NLR, PLR, MLR, NLR/PLR, SII, SIRI, and PIV indices and pCR, disease-free survival, and overall survival in breast cancer patients who received NAC.

Keywords: Neoadjuvant chemotherapy. Breast cancer. Disease-free survival.

Resumen

Objetivo: El propósito principal en las pacientes con cáncer de mama que reciben quimioterapia neoadyuvante (NAC) es lograr una respuesta patológica completa (RPC). Nuestro estudio examina si el índice neutrófilos/linfocitos (INL), el índice plaquetas/linfocitos (IPL), el índice monocitos/linfocitos (IML), la ratio INL/IPL, el índice de inmunidad-inflamación sistémica (IIS), el índice de respuesta inflamatoria sistémica (IRIS) y el valor de pan-inmuno inflamación (VPI) pueden utilizarse para determinar la RPC y el pronóstico. **Métodos:** El estudio incluyó 228 pacientes que recibieron NAC por cáncer de mama entre 2010 y 2021. **Resultados:** Cincuenta y nueve (25.9%) pacientes tuvieron RPC a NAC. En el análisis univariante, se encontró una diferencia significativa entre el estado de los ganglios clínicos, el receptor de estrógenos, el receptor de progesterona, el receptor del factor de crecimiento epidérmico humano 2 (HER2), los subtipos moleculares de cáncer de mama y la RPC ($p < 0,05$). Sin embargo, en el análisis multivariante solo los subtipos moleculares de cáncer de mama y el estado de HER2

*Correspondence:

Cem Karaali

E-mail: cemkaraali@gmail.com

2444-0507/© 2024 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 03-04-2024

Date of acceptance: 15-08-2024

DOI: 10.24875/CIRUE.M24000894

Cir Cir (Eng). 2025;93(6):571-581

Contents available at PubMed

www.cirugiaycirujanos.com

se asociaron significativamente con la RPC ($p < 0.05$). **Conclusiones:** Nuestro estudio no encontró una relación significativa del INL, el IPL, el IML, la INL/IPL, el IIS, el IRIS y el VPI con la RPC, la supervivencia libre de enfermedad y la supervivencia global en pacientes con cáncer de mama que recibieron NAC.

Palabras clave: Quimioterapia neoadyuvante. Cáncer de mama. Supervivencia libre de enfermedad.

Introduction

Breast cancer surpassed lung cancer as the most common cause of cancer globally in 2020. In addition, breast cancer still accounts for a significant proportion of cancer-related deaths among women¹. Therefore, focusing on the treatment of patients with breast cancer is critical for public health.

In recent years, neoadjuvant chemotherapy (NAC) has been used with increasing frequency in breast cancer. However, there are uncertainties about which patients are sensitive to NAC and which patients are not sensitive to NAC and, therefore, delayed surgical treatment. Hence, it is essential to determine who will benefit from NAC to optimize the treatment of patients. Estrogen receptor (ER), progesterone receptor (PR), Ki67 level, and human epidermal growth factor receptor 2 (HER2) status are often used to predict response to NAC. However, they do not accurately predict pathologic complete response (pCR). Tumor-infiltrating lymphocytes (TILs), programmed death-ligand 1 expression, artificial intelligence-assisted magnetic resonance imaging, quantitative ultrasound, and 18F-fluorodeoxyglucose positron emission tomography/computed tomography can also be used. However, these examinations are costly. In addition, access to these examinations is only sometimes possible, and further studies are needed for their use in routine practice². Therefore, there is a need for parameters that are inexpensive, easily accessible, and not difficult to interpret³. For this purpose, the literature has turned toward inflammatory bioindices analyzed from peripheral blood. These indices are calculated by various ratios of white blood cell (W), neutrophil (N), monocyte (M), platelet (P), and lymphocyte (L) cells in peripheral blood. The literature has mainly focused on the association of neutrophil/lymphocyte ratio (NLR) and platelet/lymphocyte ratio (PLR) with pCR and survival. However, studies have shown that NLR and PLR have conflicting results in predicting pCR, disease-free survival (DFS), and overall survival (OS)⁴⁻⁸. However, few studies have focused on more combined peripheral indices such as NLR/PLR, systemic immune-inflammation index (SII), systemic

inflammation response index (SIRI), and pan-immune-inflammation-value (PIV). These studies suggest that more complex calculations predict PCR and prognosis better than individual biomarkers⁷. However, there are few publications on combined indices and conflicting reports on these indices.

The main goal of breast cancer patients receiving NAC is to achieve a PCR. In the above-mentioned studies, the main issue investigated is whether peripheral biomarkers can predict pCR in breast cancer patients receiving NAC. However, giving NAC to breast cancer patients to achieve an axillary complete response based on these indices may lead to overtreatment of the axilla because axillary dissection (AD) is recommended even if the patient's axilla remains clinically positive after NAC or if micrometastases are detected on sentinel lymph node biopsy (SLNB) in patients whose axilla turns clinically negative. However, AD is not always recommended in patients who have not received NAC if the axilla is positive under certain conditions or if macrometastasis is detected on SLNB⁹.

For the reasons mentioned above, we focused on the contradictions and problems in the clinical use of simply calculated indices such as NLR, PLR, monocyte lymphocyte ratio (MLR), and more combined indices such as NLR/PLR, SII, SIRI, and PIV. We aimed to examine the relationship of these indices with pCR, DFS, and OS in a group of breast cancer patients receiving NAC. Our study is also important as it is the first publication in the literature to evaluate these indices together in terms of both pCR and prognosis.

Methods

Patients

Patients who were admitted to SBU Izmir Tepecik Training and Research Hospital between 2010 and 2021 and received NAC for breast cancer were included in the study. After obtaining permission from the Local Ethics Committee, patient data were obtained retrospectively from the electronic record system and physical files of the medical oncology department.

All patients were diagnosed histopathologically and were presented to the multidisciplinary session after appropriate screening according to their stage. The standard treatment regimens for patients with NAC decisions are anthracycline and taxane-based. Patients with HER2-positive tumors received standard trastuzumab in addition to anthracycline and taxane.

Surgical procedures included breast conserving surgery, level 1, 2, and 3 oncoplastic surgery or mastectomy. The axilla was evaluated by imaging and physical examination after NAC. SLNB was performed in patients with clinically negative axilla after NAC. No additional procedure was performed in patients with negative SLNB, while AD was performed in patients with positive (macro- or micrometastases). Clinical and pathological tumors were defined according to the American Joint Committee Cancer Staging Manual 8th Edition.

Exclusion criteria were as follows: diagnosis of cancer in an organ other than breast cancer, active infection at the time of diagnosis, inflammatory or autoimmune disease, recent steroid therapy, inflammatory breast cancer, metastasis to distant organs at the time of diagnosis, no surgery after NAC, and no follow-up.

Pathological assessments

Residual cancer burden (RCB) was calculated by the computerized MD Anderson RCB Calculator¹⁰. Accordingly, $RCB\ 0 = pCR$. pCR was defined as the absence of a tumor in both breast and axillary lymph nodes (LNs) after NAC (ypT0/ypTis, ypN0). The non-pCR group included RCBs 1, 2, and 3 (minimal, moderate, and extensive residual disease, respectively).

Study patients were classified into four main molecular subtypes. HR-positivity (HR+) was defined as ER-positive and/or PR-positive, while HR-negativity (HR-) was defined as ER-negative and PR-negative. HER2-positivity (HER2+) was defined as a score of 3+ on immunohistochemical staining and/or positive *HER2* gene amplification by FISH. Triple-negative breast cancer was defined as HR-/HER2-, HER2-enriched breast cancer as HR-/HER2+, luminal A as HR+, HER2-, $Ki67 \leq 14$ and luminal B as HR+, HER2-, $Ki67 > 14$ or HR+, and HER2⁺¹¹.

Blood sample analysis

NLR, PLR, MLR, NLR/PLR, SII, SIRI, and PIV values were calculated from peripheral blood values obtained from patients diagnosed with breast cancer

before NAC treatment. The indices were calculated as follows; NLR; N/L, PLR; P/L, MLR; M/L, SII; N*P/L, SIRI; N*M/L, PIV; and N*P*M/L.

Patient follow-up and prognostic assessment

The patients whose treatments were completed were followed up in outpatient departments. Patients were followed up by physical examination, blood tests, and imaging methods and additional procedures were performed when necessary.

Statistical analysis

Statistical analyses were performed using IBM® Statistical Package for the Social Sciences (SPSS)® 26 (IBM Corp. Released 2019. IBM SPSS Statistics for Windows, Version 26.0. Armonk, NY: IBM Corp) software. The conformity of the variables to normal distribution was analyzed using analytical methods (Kolmogorov-Smirnov/Shapiro-Wilk tests). Descriptive analyses were given as mean \pm standard deviation for normally distributed variables, median and min-max for those not normally distributed. Descriptive statistics of categorical variables obtained from sociodemographic and clinical information were analyzed using frequency and percentage values. Pearson's or Fisher's Exact X^2 test was used to examine the characteristics of patients between pCR response states. With receiver operative characteristics (ROC) analysis, cutoff values were determined for NLR, MLR, PLR, PIV, SII, and SIRI parameters (follow-up parameters analyzed) according to pCR status. According to these cutoff values, dichotomous (Low/High) variables were created for each parameter. Univariate and multivariate logistic regression analyses were performed to evaluate the effects of the characteristic variables of the disease and the follow-up parameters on the pCR response. Cox regression survival analysis was performed on these follow-up parameters (NLR, PLR, NLR/PLR, MLR, SIRI, SII, and PIV) to examine whether they could be predictive markers on OS and DFS times and whether a model could be established. Kaplan–Meier survival analysis was performed to compare OS and DFS durations between dichotomous (Low/High) variables of NLR, PLR, NLR/PLR, MLR, SIRI, SII, and PIV parameters. P values below 0.05 were considered statistically significant.

Table 1. Clinicopathological characteristics of the patients

Characteristics	n	%
Age		
Median (range)	50 (24-78)	
Clinical T stage		
1	31	13.6
2	150	65.8
3	31	13.6
4	16	7.0
Clinical T stage groups		
T1-T2	181	79.4
T3-T4	47	20.6
Clinical node stage		
0	46	20.2
1	137	60.1
2	41	18.0
3	4	1.8
Clinical node status		
Negative	46	20.2
Positive	182	79.8
Stage		
1	3	1.3
2	147	64.5
3	78	34.2
Stage groups		
≤ 2	150	65.8
3	78	34.2
Grade		
Missing	3	1.3
Good	5	2.2
Moderate	126	55.3
Poor	94	41.2
Molecular subtype		
Luminal A	24	10.5
Luminal B	123	53.9
ER and PR-. Her2+	34	14.9
Triple-negative	47	20.6
Histotype		
Invasive ductal carcinoma	206	90.4
Other histology types	22	9.6
Estrogen receptor status		
Negative	81	35.6
Positive	147	64.4
PR receptor status		
Negative	83	36.4
Positive	145	63.6
HER-2 status		
Negative	162	71.1
Positive	66	28.9
Ki-67 index, n (%)		
Median (range)	30 (0-90)	
Ki67 classification		
Ki67 < 14	34	14.9
Ki67 ≥ 14	194	85.1

(Continues)

Table 1. Clinicopathological characteristics of the patients (continued)

Characteristics	n	%
Pathologic complete response		
Yes	59	25.9
No	169	74.1
Recurrence		
No	188	82.5
Yes	40	17.5
Recurrence location		
No	188	82.5
Liver	5	2.2
Bone	3	1.3
Lung	6	2.6
Brain	4	1.8
Multiple	16	7.0
Breast axilla	6	2.6
Mortality		
Ex	43	18.9
Live	185	81.1

HER2: human epidermal growth factor receptor 2.

Results

The study included 228 patients with breast cancer who received NAC. The median age of the patients was 50 years (range 24-78). Fifty-nine (25.9%) patients had pCR to NAC and 169 (74.1%) patients had non-pCR. Recurrence was detected in 40 (17.5%) patients. Forty-three (18.9%) of the patients died. Detailed clinicopathological characteristics of the patients are summarized in table 1.

The relationship between pathological complete response and indices was calculated according to ROC analysis. Cutoff values of NLR, PLR, MLR, NLR/PLR, SII, SIRI, and PIV indices were calculated according to maximum specificity and sensitivity. According to this analysis, no significant difference was found between pCR and indices ($p > 0.05$) (Table 2).

The relationship between pCR and clinicopathological characteristics of the patients was examined by univariate and multivariate analyses. In univariate analysis, a significant difference was found between clinical node status, ER, PR, HER-2, breast cancer molecular subtypes, and pCR ($p < 0.05$). However, in multivariate analysis, only breast cancer molecular subtypes and HER-2 status were significantly associated with pCR ($p < 0.05$). Detailed data are given in table 3.

Cox regression analysis was applied to examine whether NLR, PLR, MLR, NLR/PLR, SII, SIRI, and PIV

Table 2. ROC curve analyses for pathological complete response

Test result variable (s)	AUC	Standard error	p	95% CI for AUC		Sensitivity (%)	Specificity (%)	Cut-off value
				Lower bound	Upper bound			
MLR	0.581	0.044	0.067	0.494	0.668	55.2	56.4	0.262
NLR	0.566	0.045	0.138	0.477	0.654	55.2	55.8	2.24
PIV	0.557	0.045	0.198	0.468	0.646	55.2	56.4	335.7
PLR	0.500	0.047	0.998	0.407	0.593	50.0	57.0	141.2
SII	0.540	0.046	0.362	0.450	0.630	50.0	64.2	597.5
SIRI	0.584	0.044	0.057	0.498	0.669	53.4	57.6	1.18

The test result variable (s): NLR, MLR, PLR, SII, and SIRI have at least one tie between the positive actual state group and the negative actual state group. $p < 0.05$ considered significant. Youden's J statistic used for cutoff. AUC: area under curve; CI: confidence interval; ROC: receiver operating characteristic; MLR: monocyte/lymphocyte ratio; NLR: neutrophil/lymphocyte ratio; PIV: pan-immune-inflammation-value; PLR: platelet/lymphocyte ratio; SII: systemic immune-inflammation index, SIRI: systemic inflammation response index.

Table 3. Univariate and multivariate logistic regression analysis for the predictors of pathological complete response

Factors	Univariate analysis			Multivariate analysis				
	OR	95% C.I. for EXP (B)		p	OR	95% C.I. for EXP (B)		p
		Lower	Upper			Lower	Upper	
Age	1.0	1.0	1.1	0.142				
Clinical T stage (T1-2 vs. T3-4)	1.2	0.6	2.5	0.664				
Clinical node status (Positive vs. Negative)	2.8	1.4	5.6	0.003	2.1	0.8	4.5	0.057
Triple-negative versus luminal A	17.0	2.1	136.9	0.008	30.8	2.5	380.9	0.008
Triple-negative versus luminal B	3.8	1.8	8.1	< 0.0001	14.8	2.8	79.5	0.002
Triple-negative versus ER and PR-. HER2+	1.6	0.6	3.7	0.356	4.1	1.1	16.0	0.040
Stage (≤ 2 vs. 3)	1.2	0.7	2.2	0.563				
Histopathology (IDC vs. Other)	1.9	0.8	4.9	0.172				
ER (Positive vs. Negative)	4.1	2.2	7.6	< 0.0001	2.9	0.4	24.0	0.325
PR (Positive vs. Negative)	3.7	2.0	6.9	< 0.0001	1.4	0.2	7.6	0.737
HER-2 status (Negative vs. Positive)	3.8	2.0	7.2	< 0.0001	6.1	2.2	16.9	< 0.0001
Ki67 index (High vs. Low)	1.2	0.5	2.7	0.735				
MLR group (High vs. Low)	1.6	0.9	2.9	0.131				
NLR group (High vs. Low)	1.5	0.8	2.7	0.193				
PIV group (High vs. Low)	1.5	0.8	2.6	0.221				
PLR group (High vs. Low)	1.3	0.7	2.3	0.429				
SII group (High vs. Low)	1.8	1.0	3.3	0.055				
SIRI group (High vs. Low)	1.6	0.9	2.3	0.140				

Logistic regression analysis was used and $p < 0.05$ considered significant.

CI: confidence interval; OR: odds ratio; ER: estrogen receptor; PR: progesterone receptor; HER2: human epidermal growth factor receptor 2; MLR: monocyte/lymphocyte ratio; NLR: neutrophil/lymphocyte ratio; PIV: pan-immune-inflammation-value; PLR: platelet/lymphocyte ratio; SII: systemic immune-inflammation index, SIRI: systemic inflammation response index.

Table 4. Univariate and multivariate Cox regression analysis for OS and DFS

Variables	Univariate analysis						Multivariate analysis					
	β	SE	p	OR	95.0% CI for OR		β	SE	p	OR	95.0% CI for OR	
					Lower	Upper					Lower	Upper
OS												
MLR	-0.18	0.14	0.177	0.83	0.64	1.09	-0.23	0.18	0.200	0.80	0.56	1.13
NLR	-0.09	0.13	0.518	0.92	0.71	1.19	-0.23	0.19	0.224	0.79	0.55	1.15
PIV	0.04	0.13	0.783	1.04	0.80	1.35	0.09	0.23	0.700	1.09	0.70	1.71
PLR	-0.10	0.13	0.468	0.91	0.70	1.18	-0.14	0.18	0.435	0.87	0.60	1.24
SII	0.12	0.14	0.401	1.12	0.86	1.47	0.35	0.21	0.095	1.42	0.94	2.16
SIRI	-0.05	0.13	0.725	0.95	0.73	1.24	0.05	0.24	0.828	1.05	0.66	1.68
DFS												
NLR	-0.187	0.135	0.168	0.830	0.636	1.082	-0.189	0.174	0.276	0.828	0.589	1.163
MLR	-0.051	0.134	0.705	0.951	0.732	1.235	-0.174	0.191	0.363	0.840	0.578	1.222
PLR	0.047	0.134	0.728	1.048	0.806	1.363	0.366	0.253	0.148	1.441	0.878	2.365
PIV	-0.077	0.134	0.567	0.926	0.712	1.205	-0.094	0.186	0.613	0.910	0.633	1.310
SII	0.148	0.138	0.282	1.159	0.885	1.519	0.343	0.210	0.102	1.409	0.934	2.126
SIRI	-0.119	0.134	0.375	0.888	0.682	1.155	-0.337	0.255	0.188	0.714	0.433	1.178

Cox Regression analysis was used and $p < 0.05$ considered significant. CI: confidence interval; SE: standard error; DFS: disease-free survival; OS: overall survival; MLR: monocyte/lymphocyte ratio; NLR: neutrophil/lymphocyte ratio; PIV: pan-immune-inflammation-value; PLR: platelet/lymphocyte ratio; SII: systemic immune-inflammation index; SIRI: systemic inflammation response index; OR: odds ratio.

indices could be used in univariate and multivariate calculations for OS and DFS follow-up. It was determined that each index could not be used in the follow-up of OS and DFS in the analyses performed alone or in combination ($p > 0.05$). Detailed data are given in table 4.

The low and high groups created in NLR, PLR, MLR, NLR/PLR, SII, SIRI, and PIV indices separately in pCR and non-pCR groups were evaluated in terms of OS, and DFS evaluation. However, no significant discrimination was found in the mentioned indices ($p > 0.05$). Detailed representation is available in figures 1 and 2.

Discussion

Our study evaluated NLR, PLR, MLR, NLR/PLR, SII, SIRI, and PIV indices regarding pCR, DFS, and OS in breast cancer patients receiving NAC. In our study, these indices failed to predict pCR, DFS and OS. This finding is critical because our study underlines that the parameters mentioned above are still contradictory in deciding NAC in clinical use. As far as we can see, none of the publications on this subject have analyzed both pCR and prognosis together using the above-mentioned indices simultaneously.

As a result of tumor-induced changes in the immune system, inflammatory immune cells in the peripheral blood may be affected¹². Lymphocytes in peripheral blood are related to adaptive immunity. In addition,

lymphocytes may show anti-tumoral activity by inhibiting tumor growth and progression. Platelets play an essential role in the development and spread of cancer and in regulating inflammation. Neutrophils have a central role in the systemic inflammatory response and immunity. Neutrophils can also initiate metastasis in breast cancer by secreting immunosuppressive mediators and angiogenetic factors. Monocytes are also associated with tumor angiogenesis and metastasis^{2,13}. Theoretically, it can be said that each of the mentioned indices or their combinations may be related to pCR, DFS, and OS in breast cancer patients. In the literature, an increasing number of publications have examined the relationship between these indices and breast cancer patients who will receive NAC. Among these indices, NLR and PLR indices have been analyzed most frequently for this purpose. However, studies have shown that NLR and PLR have conflicting results in predicting pCR, DFS and OS in breast cancer patients receiving NAC⁴⁻⁸. With the idea that more combined indices may be better predictors, Graziano et al.⁶ compared NLR and PLR values with NLR/PLR combination. This study found that NLR and PLR alone could not predict pCR, whereas there was a significant relationship between NLR/PLR combination and pCR. In two other studies in the literature, it was shown that NLR and PLR values were better in indicating pCR when evaluated in combination rather than when used alone^{3,14}. In these three studies, NLR/PLR combination was found to be significant in




Figure 1. Comparison of low and high overall survival times in rates and indices in pathologic complete response (pCR) and non-pCR groups.




Figure 2. Comparison of low and high disease-free survival times in rates and indices in pathologic complete response (pCR) and non-pCR groups.

evaluating pCR in contrast to our study. Among these three studies, only Kim et al.³ analyzed the relationship between prognosis and NLR/PLR combination and found a significant relationship in contrast to our study. However, there is a significant methodological difference between the study of Kim et al.³ and our study regarding pCR evaluation, as we will mention later. In addition, there are not enough publications on NLR/PLR index in the literature yet. As far as we can see, our article is the first study in the English literature showing that the NLR/PLR combination fails to predict pCR and prognosis.

One of the other combined parameters is SIRI. In SIRI calculation, monocytes are included in the calculation in addition to NLR. First, Dong et al.¹⁵ analyzed the relationship between SIRI, and pCR. According to this study, SIRI showed pCR to NAC better than other indices (lymphocyte monocyte ratio, absolute lymphocyte count, and NLR). However, this study did not analyze DFS and OS. In the study of Chen et al.¹³, SIRI was significant in determining OS and DFS in breast cancer patients receiving NAC. However, no significant difference was found in terms of pCR, similar to our study.

In SII, another combined parameter, platelets, which have an essential role in the development and spread of breast cancer, are included in the calculation in addition to NLR. In the study of Yang et al.¹⁶, a significant correlation was found between NLR, PLR, SII, and pCR. In contrast, in another study of Chen et al.¹⁷, no significant correlation was found between SII and pCR as in our study. However, Chen et al.¹⁷ found that patients with low SII values had longer DFS and OS rates.

Compared with individual blood cell parameters, combined indices may more comprehensively identify multiple components of antitumor immunity, each of which may reflect and regulate different aspects of antitumor immunity. For this purpose, in the study of Şahin et al.⁷, the PIV index, which includes all the parameters mentioned above (neutrophils, platelets, monocytes, and lymphocytes), was used. Indeed, supporting the theoretical knowledge, NLR, PLR, MLR, and PIV successfully showed pCR in univariate analysis. In contrast, only PIV, a more complex marker, maintained its statistical value in multivariate analysis. Furthermore, PIV was found to have a prognostic effect on survival. This is important because it is the first study in the literature to examine the relationship between PIV value and response to NAC and survival

in breast cancer. However, contrary to this study, we could not reach similar results in our study.

When considered together with our study, many factors can be counted to explain the contradictory results of those mentioned above simple and complex indices on pCR and prognosis in breast cancer patients receiving NAC. Many factors can easily affect inflammatory markers (lymphocytes, neutrophils, platelets, and monocytes). For example, many factors can easily affect inflammatory events, including connective tissue diseases, medications used, bacterial or viral diseases, nutritional status, severe stress, and severe exercise³. In addition, breast cancers have the least immunogenic type compared to other cancer types². In addition, TILs are less common in HR+ positive breast cancers compared to HR- breast cancers¹⁸. In other words, anticancer immunity is not dominant in hormone receptor-positive breast cancers. Therefore, it is thought that absolute lymph count is not a good predictor in HR+ breast cancers compared to HR- breast cancers¹⁹. Lymphocyte count is included in the denominator of calculations in all indices we investigated and, therefore, affects all calculations. Finally, it is thought that lymphocytic infiltration in the tumor microenvironment is not sufficiently reflected in peripheral blood cells²⁰. All the factors mentioned above give us a clue to explain the conflicting data on these indices.

One of the reasons for the conflicting reports on these indices is the methodological difference between studies. There are differences in the definition of pCR in the literature. Miller-Payne (MP) classification was used in some publications. In this classification, unlike the RCB classification, the axilla is not evaluated, and only the response in the breast is considered. In other words, a patient with a complete response to the tumor in the breast is considered to have a complete response even if tumors are detected in the LNs. In addition, in some of the studies using the MP classification, both MP 4 (> 90% response to NAC) and MP 5 (no tumor cells) were included in the group of patients who responded to NAC. However, it is known that the survival time of patients with pCR is higher than those with residual cancer²¹. For example, in the studies of Kim et al.³ evaluating NLR/PLR combination and in two separate studies of Chen et al. evaluating SIRI¹³ and SII¹⁷, clinical response was evaluated as MP 4-5. In contrast to these studies, RCB was used in two other studies evaluating NLR/PLR combination^{6,14}, SIRI by Dong et al.¹⁵, NLR, PLR, and SII by Yang et al.¹⁶, and PIV index by Şahin et al.⁷.

This confusion in the evaluation of pathological response may have caused contradictions in the correct evaluation of the indices. However, to use a standard criterion in the literature and make comparisons, it would be more accurate to use RCB when investigating pCR as in our study²².

Although there are publications examining the relationship of these indices with pCR and prognosis, there are very few comments on how their clinical use will be in routine practice. Therefore, we would like to discuss a topic that has not been emphasized in the literature. In the recently updated NCCN guidelines, patients with clinically node negative or clinically non-palpable lymph but with up to two positive LNs by imaging or biopsy may undergo SLNB if NAC was not performed. However, in breast cancer patients who receive NAC, AD is performed if there is a clinically positive LN after NAC or a positive sentinel LN (micro or macrometastasis)⁹. In this case, in patients with low axillary LN metastasis in whom NAC is not considered (especially in HR+, HER2- tumors), giving NAC by relying on the indices listed above may lead to some problems. The most important of these problems will arise if the patient's axilla does not respond fully to NAC. A patient with a low axillary tumor burden who would not usually be given NAC can typically be satisfied with SLNB. In contrast, AD will be necessary in the patient receiving NAC, even if micrometastases are found on SLNB. In this case, overtreatment will be applied to the patient's axilla. In addition, since the methods used to reduce false negativity during SLNB application in the patient receiving NAC are technically more complex, we argue that the decision of NAC in patients with breast cancer should not be based on the mentioned indices.

Our study has limitations: a small number of patients, single-center experience, and retrospective nature.

Conclusion

Our study did not find a significant relationship between NLR, PLR, MLR, NLR/PLR, SII, SIRI, and PIV indices and pCR, DFS, and OS in breast cancer patients who received NAC. In our opinion, the decision to apply NAC to the patient should still be made according to the molecular subtypes of the breast tumor. It should continue to be discussed in patient-based multidisciplinary sessions. In addition, although no correlation was found between the indices mentioned in our study and pCR, DFS, and OS, it would

be helpful to continue studies on this subject considering the advantages of these indices, such as easy calculation and obtaining them from blood samples in routine practice.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of humans and animals. The authors declare that no experiments involving humans or animals were conducted for this research.

Confidentiality, informed consent, and ethical approval. The study does not involve patient personal data nor requires ethical approval. The SAGER guidelines were followed according to the nature of the study.

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing of this manuscript.

References

1. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* 2021;71:209-49.
2. Derouane F, Van Marcke C, Berlière M, Gerday A, Fellah L, Leconte I, et al. Predictive biomarkers of response to neoadjuvant chemotherapy in breast cancer: current and future perspectives for precision medicine. *Cancers (Basel).* 2022;14:3876.
3. Kim HY, Kim TH, Yoon HK, Lee A. The role of neutrophil-lymphocyte ratio and platelet-lymphocyte ratio in predicting neoadjuvant chemotherapy response in breast cancer. *J Breast Cancer.* 2019;22:425-38.
4. Corbeau I, Jacot W, Guiu S. Neutrophil to lymphocyte ratio as prognostic and predictive factor in breast cancer patients: a systematic review. *Cancers (Basel).* 2020;12:958.
5. Cullinane C, Creavin B, O'Leary DP, O'Sullivan MJ, Kelly L, Redmond HP, et al. Can the neutrophil to lymphocyte ratio predict complete pathologic response to neoadjuvant breast cancer treatment? A systematic review and meta-analysis. *Clin Breast Cancer.* 2020;20:e675-81.
6. Graziano V, Grassadonia A, Iezzi L, Vici P, Pizzuti L, Barba M, et al. Combination of peripheral neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio is predictive of pathological complete response after neoadjuvant chemotherapy in breast cancer patients. *Breast.* 2019;44:33-8.
7. Şahin AB, Cubukcu E, Ocak B, Deligonul A, Oyucu Orhan S, Tolunay S, et al. Low pan-immune-inflammation-value predicts better chemotherapy response and survival in breast cancer patients treated with neoadjuvant chemotherapy. *Sci Rep.* 2021;11:14662.
8. Xu J, Ni C, Ma C, Zhang L, Jing X, Li C, et al. Association of neutrophil/lymphocyte ratio and platelet/lymphocyte ratio with ER and PR in breast cancer patients and their changes after neoadjuvant chemotherapy. *Clin Transl Oncol.* 2017;19:989-96.
9. Gradishar WJ, Moran MS, Abraham J, Abramson V, Aft R, Agnese D, et al. NCCN guidelines® insights: breast cancer, version 4.2023. *J Natl Compr Canc Netw.* 2023;21:594-608.

10. Residual Cancer Burden Calculator. MD Anderson Cancer Center; 2023 Available from: <https://www3.mdanderson.org/app/medcalc/index.cfm?pagename=jsonconvert3>
11. Kaytaz Tekyol K, Gurleyik G, Aktaş A, Aker F, Tanrikulu E, et al. Pathological complete response to neoadjuvant chemotherapy in patients with breast cancer: the relationship between inflammatory biomarkers and molecular subtypes. *Cureus*. 2021;13:e14774.
12. Hiam-Galvez KJ, Allen BM, Spitzer MH. Systemic immunity in cancer. *Nat Rev Cancer*. 2021;21:345-59.
13. Chen L, Kong X, Wang Z, Wang X, Fang Y, Wang J. Pretreatment systemic inflammation response index in patients with breast cancer treated with neoadjuvant chemotherapy as a useful prognostic indicator. *Cancer Manag Res*. 2020;12:1543-67.
14. Babbar P, Rudresha AH, Dasappa L, Abraham LJ, Babu MC, Lokesh KN, et al. Combined peripheral neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio as a predictive biomarker for pathological complete response after neoadjuvant chemotherapy in triple-negative breast cancer patients. *Oncol J India*. 2022;6:8-13.
15. Dong J, Sun Q, Pan Y, Lu N, Han X, Zhou Q. Pretreatment systemic inflammation response index is predictive of pathological complete response in patients with breast cancer receiving neoadjuvant chemotherapy. *BMC Cancer*. 2021;21:700.
16. Yang G, Liu P, Zheng L, Zeng J. Novel peripheral blood parameters as predictors of neoadjuvant chemotherapy response in breast cancer. *Front Surg*. 2022;9:1004687.
17. Chen L, Kong X, Wang Z, Wang X, Fang Y, Wang J. Pre-treatment systemic immune-inflammation index is a useful prognostic indicator in patients with breast cancer undergoing neoadjuvant chemotherapy. *J Cell Mol Med*. 2020;24:2993-3021.
18. Watanabe T, Hida AI, Inoue N, Imamura M, Fujimoto Y, Akazawa K, et al. Abundant tumor infiltrating lymphocytes after primary systemic chemotherapy predicts poor prognosis in estrogen receptor-positive/HER2-negative breast cancers. *Breast Cancer Res Treat*. 2018;168:135-45.
19. Araki K, Ito Y, Fukada I, Kobayashi K, Miyagawa Y, Imamura M, et al. Predictive impact of absolute lymphocyte counts for progression-free survival in human epidermal growth factor receptor 2-positive advanced breast cancer treated with pertuzumab and trastuzumab plus eribulin or nab-paclitaxel. *BMC Cancer*. 2018;18:982.
20. Suppan C, Bjelic-Radisic V, La Garde M, Groselj-Strele A, Eberhard K, Samonigg H, et al. Neutrophil/Lymphocyte ratio has no predictive or prognostic value in breast cancer patients undergoing preoperative systemic therapy. *BMC Cancer*. 2015;15:1027.
21. Cortazar P, Zhang L, Untch M, Mehta K, Costantino JP, Wolmark N, et al. Pathological complete response and long-term clinical benefit in breast cancer: the CTNeoBC pooled analysis. *Lancet*. 2014;384:164-72.
22. Yau C, Osdoit M, Van Der Noordaa M, Shad S, Wei J, De Croze D, et al. Residual cancer burden after neoadjuvant chemotherapy and long-term survival outcomes in breast cancer: a multicentre pooled analysis of 5161 patients. *Lancet Oncol*. 2022;23:149-60.

Pelvic index/mesorectal length ratio: a new predictive factor for rectal cancer prognosis and low anterior resection syndrome

Relación índice pélvico/longitud mesorrectal: un nuevo factor predictivo para el pronóstico del cáncer de recto y el síndrome de resección anterior baja

Muhammed Gurluk¹, Onur O. Karagulle^{2*}, Ekrem Cakar², Betül Avcı³, Ensar Cakır⁴,
Ibrahim T. Rakıcı⁵, and Mert M. Sevinc²

¹Department of General Surgery, Bingöl Genç State Hospital, Bingöl; ²Department of General Surgery, Istanbul Training and Research Hospital, University of Health Sciences, Istanbul; ³Department of Internal Medicine, Basaksehir Cam and Sakura City Hospital, Istanbul; ⁴Department of General Surgery, Kilis Prof. Dr. Alaeddin Yavaşca State Hospital, Kilis; ⁵Department of Radiology, Istanbul Training and Research Hospital, University of Health Sciences, Istanbul. Turkey

Abstract

Objective: This study aimed to evaluate whether the pelvic index/mesorectal length (PI/ML) ratio is an effective factor in recurrence, anastomotic leakage, mesorectal excision status, and low anterior resection syndrome (LARS) development in rectal cancer patients undergoing total mesorectal excision. **Methods:** A total of 47 patients who underwent surgery for rectal cancer between January 2016 and December 2021 were included. Demographics, clinical data, and pre-operative PI measurements were recorded. Post-operative LARS was assessed using the LARS questionnaire in patients followed for at least 12 months. **Results:** A significant association was found between PI/ML ratio and tumor recurrence ($p < 0.0001$). Receiver operating characteristic analysis identified a cutoff value of PI/ML < 1.6 for predicting tumor recurrence, with 100% sensitivity and 84.6% specificity. After applying Bonferroni correction for multiple comparisons ($n = 4$, adjusted significance threshold $p < 0.0125$), this association remained statistically significant. The association between PI/ML ratio and anastomotic leakage (cutoff < 2.15 ; sensitivity 100%, specificity 53.7%) showed marginal significance ($p = 0.009$). No significant association was found between PI/ML ratio and LARS or mesorectal excision status after correction. **Conclusions:** PI/ML ratio appears to be a useful predictor for tumor recurrence and may help identify patients at risk for anastomotic leakage in rectal cancer surgery. However, this ratio was not significantly associated with the development of LARS. Further research with larger cohorts is needed to validate these findings and clarify the potential prognostic value of the ratio.

Keywords: Rectal cancer. Pelvic index. Low anterior resection syndrome. Total mesorectal excision.

Resumen

Objetivo: Evaluar si la relación índice pélvico/longitud mesorrectal (IP/LM) es un factor predictivo eficaz en la recurrencia, la fuga anastomótica, el estado de la escisión mesorrectal y el desarrollo de síndrome de resección anterior baja (SRAB) en pacientes con cáncer de recto sometidos a escisión mesorrectal total. **Métodos:** Se incluyeron 47 pacientes sometidos a cirugía por cáncer rectal entre enero de 2016 y diciembre de 2021. Se registraron características demográficas, datos clínicos y mediciones preoperatorias del índice pélvico. El SRAB posoperatorio se evaluó mediante un cuestionario en pacientes con un seguimiento mínimo de 12 meses. **Resultados:** Se encontró una asociación significativa entre la relación IP/LM y la recurrencia tumoral ($p < 0.0001$). El análisis de la curva ROC identificó un valor de corte de IP/LM < 1.6 para predecir la recurrencia

*Correspondence:

Onur Olgac-Karagulle
E-mail: onurolgackaragulle@gmail.com
2444-0507/© 2025 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 15-12-2024
Date of acceptance: 28-08-2025
DOI: 10.24875/CIRUE.M25000895

Cir Cir (Eng). 2025;93(6):582-590
Contents available at PubMed
www.cirurgiaycirujanos.com

tumoral, con una sensibilidad del 100% y una especificidad del 84.6%. Tras aplicar la corrección de Bonferroni para comparaciones múltiples ($n = 4$, umbral de significación ajustado $p < 0.0125$), esta asociación siguió siendo estadísticamente significativa. La asociación entre la relación IP/LM y la fuga anastomótica (valor de corte < 2.15 ; sensibilidad 100%, especificidad 53.7%) mostró una significación marginal ($p = 0.009$). No se encontró ninguna asociación significativa entre la relación IP/LM y el SRAB, ni con el estado de la escisión mesorrectal tras la corrección. **Conclusiones:** La relación IP/LM parece ser un predictor útil para la recurrencia tumoral y podría ayudar a identificar a los pacientes con riesgo de fuga anastomótica en la cirugía de cáncer de recto. Sin embargo, esta relación no se asoció significativamente con el desarrollo de SRAB. Se necesitan más estudios con cohortes más grandes para validar estos hallazgos y aclarar el potencial valor pronóstico de esta relación.

Palabras clave: Cáncer rectal. Índice pélvico. Síndrome de resección anterior baja (SRAB). Escisión mesorrectal total.

Introduction

Colorectal cancer (CRC) remains a major health problem worldwide and is the third leading cause of cancer-related death. Recent data indicate that rectal cancer accounts for nearly 39% of all CRC cases and contributes substantially to morbidity and mortality¹⁻³. A narrow pelvis and difficult pelvic anatomy are considered to be important risk factors for rectal cancer surgery². Different angles and strictures in the pelvis can lead to the possibility of difficult and incomplete resection. Although the length of the mesorectum may appear relatively short or long on the pathology slides, depending on an individual's pelvic anatomy, pathological evaluation can provide significant insights^{1,2}.

Appropriate oncologic resection is critical in the surgical treatment of rectal cancer, with total mesorectal excision (TME) becoming the gold standard². However, even with modern techniques, post-operative functional disorders such as low anterior resection syndrome (LARS) remain significant complications, occurring in up to 50-80% of patients^{3,4}. LARS encompasses symptoms such as fecal urgency, incontinence, and frequent bowel movements, significantly impacting quality of life^{4,5}.

Recent meta-analyses and multicenter trials have identified important risk factors for LARS, including low tumor height, low anastomotic height, neoadjuvant therapy, anastomotic leakage, and protective stomas^{4,5}. Furthermore, new surgical approaches such as transanal TME have shown promise in reducing conversion rates and potentially improving functional outcomes, although concerns about specific complications and local recurrence remain⁶.

The basic pathophysiology of LARS involves a combination of colonic motility disorder, neo-rectal reservoir dysfunction, and impaired anal sphincter function. Dysfunction after rectal resection leads to LARS symptoms due to denervation, decreased

functional capacity, and hypogastric plexus injury. Therefore, the pelvic angle and the size of the resected pathologic lesion are among the determining factors for the development of LARS^{7,8}.

In rectal cancer surgery, new markers are needed to predict the patient's prognosis, and studies on this topic are ongoing. In this context, pelvimetric measurements have increasingly been explored as potential predictors of surgical outcomes. Therefore, in our study, we calculated the pelvic index (PI) value based on the pre-operative tomography images of the patients. In addition, we recorded the mesorectal length (ML) and complete/incomplete mesorectal excision data from the pathological reports of rectal cancer. By calculating the "Pelvic Index/Mesorectal Length (PI/ML) Ratio" using pelvimetric measurements, we aimed to contribute to the literature by investigating the relationship between this ratio and factors that influence the prognosis of rectal cancer surgery and the development of LARS. We hypothesize that a lower PI/ML ratio is an independent predictor of an increased risk of anastomotic leakage, tumor recurrence, and the development of LARS in patients undergoing TME for rectal cancer.

Methods

Study design and participants

In this retrospective observational study, patients who underwent surgery for rectal cancer in the General Surgery Clinic of Istanbul Training and Research Hospital between January 2016 and December 2021 were analyzed. Patients who could not undergo curative surgery, whose clinical data could not be reached in the post-operative period, or who could not participate in the LARS questionnaire (or who had not completed 1 year postoperatively) were excluded from the study.




Figure 1. Magnetic resonance imaging measurements and schematic diagram for calculation of the pelvic index (PI) in rectal cancer patients. **A:** sagittal plane image showing the pelvic inlet width (a) measured as the transverse diameter of the pelvic brim, the depth of cavity of the lesser pelvis (b) the length measured between the middle point of line “a,” and the mesorectal thickness (d). **B:** axial plane image showing the mesorectal diameter (c) measured at the point of maximum width, transverse rectal diameter (e), and longitudinal rectal diameter (f). **C:** axial plane image illustrating the interspinous distance defined as the narrowest distance between the ischial spines (g). **D:** schematic diagram demonstrating the formula used for calculating the pelvic index using the distance between the spina ischialis, the diameter of the mesorectum in the axial plane, and the depth of the small pelvis.

A priori power analysis was performed using G*Power (Heinrich Heine University Düsseldorf, Germany). For detecting differences between two independent means (two-tailed *t*-test), an effect size (Cohen’s *d*) of 1.64 was assumed, with an alpha level of 0.05 and statistical power set at 95%. The effect size was estimated based on preliminary analyses of institutional data, which demonstrated substantial differences in pelvic measurements between patient groups. Similar large effect sizes have also been reported in previous studies investigating pelvic anatomy and surgical outcomes in rectal cancer. The analysis indicated that a total sample size of 22 patients (11/group) would be sufficient. As our study included 47 patients, the sample size was deemed adequately powered for the analyses conducted.

Data collection

DEMOGRAPHIC AND CLINICAL DATA

Demographic data, operative data, pathology reports, and radiologic imaging were analyzed. Clinical files of the patients were scanned, and demographic data, as well as recurrence and anastomotic leakage status, were recorded.

PRE-OPERATIVE IMAGING AND PI MEASUREMENT

All PI measurements were performed by a single experienced radiologist to ensure consistency (Fig. 1).

Interobserver variability was not assessed in this study, which represents a limitation. In the pre-operative period, tomography images taken within 1 week before surgery were evaluated by a radiologist, and the PI was calculated using the following formula presented in figure 1⁹.

PATHOLOGIC EVALUATION

We obtained the mesorectal length (ML) data from the pathologic slide reports.

LARS SCORE

The LARS questionnaire was administered via telephone to patients with at least 12 months of post-operative follow-up. According to the results of the questionnaire, patients were classified as follows:

- No LARS: 0-20 points
- Minor LARS: 21-29 points
- Major LARS: 30-49 points
- We evaluated patients in two groups: those with LARS (minor + major) and patients without LARS¹⁰.

Ethical approval

The study was initiated with the approval of decision number 391 from the Ethics Committee of Istanbul

Training and Research Hospital on December 23, 2022. This was a retrospective observational study and was not registered in a public trials registry, as registration is not required for retrospective studies under current regulations.

Results

The study was conducted with a total of 47 patients, 21 (44.7%) females and 26 (55.3%) males. The mean age was 62.1 ± 9.8 years (range: 37-86), and body mass index (BMI) was 26.6 ± 4.9 kg/m² (Table 1). The number of patients who received neoadjuvant chemoradiotherapy in our case series was 33 (70.2%), whereas the number of patients who did not receive neoadjuvant chemoradiotherapy was 14 (29.8%).

We observed an anastomotic leakage in 12.8% of patients (six cases) and detected recurrent tumoral lesions in 17% (eight cases). In the evaluation of mesorectal excision in pathologic slides, we found incomplete mesorectal excision in 36.2% (17 cases) and complete mesorectal excision in 63.8% (30 cases) (Table 1).

In terms of tumor localization, 29 (61.7%) patients had tumor localization in the proximal rectum, 13 (27.7%) in the middle rectum, and 5 (10.6%) in the distal rectum. In terms of surgical method, laparoscopic surgery was performed in 33 cases, and loop ileostomy was performed in 21 of them. Conventional surgery was performed in 13 cases, and loop ileostomy was performed in addition to resection in nine of them. When the stage distribution was analyzed in our study, the most common stage was E3b with 26 cases (54.6%), and at least 1 case (2.1%) was the E3c stage (Table 1).

Twenty patients (42.6%) showed LARS findings, and the mean LARS score was 17.1 ± 13.1 . The mean interspinous distance was 11.3 ± 1.2 cm, the radiologic diameter of the mesorectum was 7.83 ± 1.05 cm, and the depth of the small pelvic cavity was 9.64 ± 1.2 cm. The mean PI value was calculated as 36.7 ± 12.5 cm. The average pathologic mesorectal length was 16.8 ± 4.2 cm, and the ratio of PI to mesorectal length was 2.4 ± 1.3 (Table 1).

The mean BMI of the female group was 28.5 ± 5.2 and 25.0 ± 4.0 in the male group (Table 2). The mean BMI of the female group was higher than that of the male group, and this difference was statistically significant and was found to have a large effect size (Cohen's $d = 0.758$, 95% confidence interval [CI] = -1.350 – -0.158 , $p = 0.013$).

The mean PI/ML was 2.6 ± 1.3 in the female group and 2.2 ± 1.3 in the male group. There was no

Table 1. Demographic, clinical, pathological, and treatment characteristics of patients undergoing total mesorectal excision for rectal cancer

Parameters	Mean \pm SD or n (%)
Age (years)	62.1 \pm 9.8
Sex, n (%)	Male: 26 (55.3) Female: 21 (44.7)
BMI (kg/m ²)	26.6 \pm 4.9
Tumor location, n (%)	Proximal: 29 (61.7) Middle rectum: 13 (27.7) Distal: 5 (10.6)
Pathological stage (pTNM), n (%)	Stage I: 6 (12.8) Stage II: 3 (6.4) Stage III: 36 (76.6) Stage IV: 2 (4.3)
Neoadjuvant radiotherapy, n (%)	33 (70.2)
Complete mesorectal excision, n (%)	30 (63.8)
Anastomotic leakage, n (%)	6 (12.8)
Tumor recurrence, n (%)	8 (17.0)
LARS n (%)	20 (42.6)
Mean score	17.1 \pm 13.1
Interspinous distance (cm)	11.3 \pm 1.2
Mesorectum radiologic diameter (cm)	7.83 \pm 1.05
Depth of small pelvic cavity (cm)	9.64 \pm 1.2
PI (cm)	36.7 \pm 12.5
Pathologic mesorectal length (cm)	16.8 \pm 4.2
PI/ML (cm)	2.4 \pm 1.3

LARS: low anterior resection syndrome; PI/ML: pelvic index/mesorectal length; BMI: body/mass index, SD: standard deviation.

Table 2. Comparison of age, BMI, PI/ML ratio, and tumor localization between male and female patients

Parameters	Mean \pm SD		Effect size (95% CI)	p
	Female	Male		
Age	61.4 \pm 10.7	62.6 \pm 9.1	0.12 (–0.46–0.70)	0.683
BMI	28.5 \pm 5.2	25.0 \pm 4.0	–0.76 (–1.35–0.16)	0.013
PI/ML (cm)	2.6 \pm 1.3	2.2 \pm 1.3	–0.28 (–0.86–0.30)	0.339

PI/ML: pelvic index/mesorectal length; BMI: body/mass index, SD: standard deviation, 95% CI: confidence interval.

statistically significant difference between the sexes in terms of PI/ML ($p > 0.05$) (Table 2).

To correct for multiple comparisons ($n = 4$), we applied a Bonferroni correction, setting the

Table 3. ROC analysis results for predicting anastomotic leakage, tumor recurrence, complete mesorectal excision, and LARS based on the PI/ML ratio

Parameters	AUC ± SD	Cutoff	Sensitivity	Specificity	95% CI	p
Anastomotic leakage	0.833 ± 0.079	< 2.15	100	53.7	0.679-0.988	0.009
Recurrence	0.929 ± 0.037	< 1.6	100	84.6	0.858-1.000	0.00001
Complete mesorectal excision	0.520 ± 0.093	> 1.64	73.3	41.2	0.338-0.701	0.825
LARS	0.589 ± 0.088	< 1.90	55	66.6	0.436-0.730	0.315

PI/ML: pelvic index/mesorectal length; LARS: low anterior resection syndrome; AUC: area under the curve; SD: standard deviation; CI: confidence interval; ROC: receiver operating characteristic.

Table 4. Comparison of demographic and clinical characteristics between patients with and without LARS

Parameters	No. LARS (n = 27)	LARS present (n = 20)	Odd ratio/effect size (95% CI)	p
Gender, n (%)			0.40 (0.12-1.35)*	0.394
Male	13 (48.1)	13 (65.0)		
Female	14 (51.9)	7 (35.0)		
Age, X ± SD	61.67 ± 10.1	62.65 ± 9.5	0.01 (-0.57-0.60)**	0.957
BMI, X ± SD	24.77 ± 3.4	28.02 ± 5.3	0.46 (-0.13-1.05)**	0.035
Mesorectal excision, n (%)			0.20 (0.06-0.72)*	0.045
Complete	21 (77.8)	9 (45.0)		
Incomplete	6 (22.2)	11 (55.0)		
PI, X ± SD	38 ± 13	35 ± 12	-0.27 (-0.85-0.32)	0.241
Mesorectal length (ML), X ± SD	16.50 ± 3.79	18.41 ± 4.15	0.44 (-0.16-1.02)**	0.140
PI/ML, X ± SD	2.4 ± 1	1.8 ± 1	0.34 (0.02-0.25)**	0.040

*Odd ratio.

**Effect size (Cohen's d).

BMI: body mass index; X ± SD: mean ± standard deviation; PI/ML: pelvic index/mesorectal length ratio; LARS: low anterior resection syndrome.

significance threshold at $p < 0.0125$. Under this correction, the relationship between recurrence and PI/ML remained statistically significant ($p = 0.00001 < 0.0125$), and the association between anastomotic leakage and PI/ML was marginally retained ($p = 0.009 < 0.0125$). Associations with complete mesorectal excision and LARS did not reach the adjusted significance level (Table 3).

When the PI/ML cutoff value ≤ 2.1 for anastomotic leakage, sensitivity and specificity were 100% and 53.7%, respectively. There was a statistically significant correlation between PI/ML and anastomotic leakage ($p < 0.05$) (Fig. 2A). We found the sensitivity and specificity to be 100% and 84.6%, respectively, when the PI/ML cutoff value was ≤ 1.6 for recurrence. PI/ML was found to be significantly associated with the recurrence ($p < 0.05$) (Fig. 2B). Sensitivity and specificity were 73.3% and 41.2%, respectively, when the PI/ML cutoff value was > 1.64 . PI/ML was not statistically significantly associated with complete

mesorectal excision ($p > 0.05$) (Fig. 2C). Sensitivity and specificity were found to be 55% and 66.6%, respectively, when the cutoff value for PI/ML was < 1.90 . We did not find a statistically significant correlation between PI/ML and LARS ($p > 0.05$) (Fig. 2D and Table 3).

Patients in our study were divided into two groups: no LARS (0-20) and LARS (21-42 minor + major). Demographic and clinical data are analyzed in table 4. Twenty-seven patients had no LARS, and 20 patients had LARS findings (minor + major). There was no statistical significance between these groups in terms of age and gender ($p > 0.05$). BMI was 28.02 ± 5.3 in the group with LARS findings and 24.77 ± 3.4 in the group without LARS, which has a moderate effect size ($p = 0.035$; Cohen's $d = 0.462$; 95% CI: $-0.131-1.050$).

While there was no statistical difference between the LARS groups in terms of T, N, and Stage, no statistical difference was observed in terms of tumor localization ($p > 0.05$). There was no statistical significance between




Figure 2. Receiver operating characteristic analyses of anastomotic leakage (A), recurrence (B), complete mesorectal excision (C), and low anterior resection syndrome (D) according to the pelvic index/mesorectal length ratio.

the PI and ML between the groups according to the presence of LARS. The PI/ML ratio was found to be higher in individuals without LARS than in individuals with LARS. This difference had a small effect size ($p = 0.040$; Cohen's $d = 0.336$; 95% CI: 0.020-0.253) (Table 4).

Incomplete mesorectal excision was reported in 11 cases (55%) with LARS, whereas complete mesorectal excision was reported in 21 cases (77.8%) without LARS (Table 4). Cases with complete excision were 5.05 times less likely to develop LARS than individuals without complete excision ($p = 0.045$; OR = 0.198; 95% CI: 0.055-0.715).

In the correlation analysis between LARS score and demographic and clinical data, no statistical significance was observed in the correlation between age, BMI, PI, ML, and PI/ML (Table 5).

Discussion

In our study, we investigated whether the PI/ML ratio, derived from pre-operative pelvic measurements, could predict surgical and oncological outcomes in rectal cancer patients undergoing TME, which is critical for patient prognosis. After correcting for multiple

Table 5. Correlation analysis between LARS scores and continuous demographic and clinical variables

Parameters	r	p	95% CI	
			Lower	Upper
Age	-0.156	0.295	-0.431	0.146
BMI	0.198	0.182	-0.103	0.466
PI	-0.167	0.262	-0.441	0.135
ML	0.140	0.349	-0.162	0.418
PI/ML	-0.160	0.284	-0.434	0.142

LARS: low anterior resection syndrome; BMI: body mass index; CI: confidence intervals;
PI/ML: pelvic index/mesorectal length ratio.

comparisons using the Bonferroni method, we found that the PI/ML ratio remained significantly associated with tumor recurrence and showed a marginally significant association with anastomotic leakage. However, the PI/ML ratio was not significantly associated with the development of LARS.

The relationship between pelvic anatomy and surgical outcomes has been discussed in the literature^{11,12}. Several studies have shown that a narrow pelvis and difficult pelvic anatomy can lead to more challenging resections, higher rates of incomplete excision, and potentially higher recurrence rates^{7,8,11,12}. High-resolution magnetic resonance imaging studies have recently provided more detailed insights, demonstrating that pelvic dimensions and mesorectal involvement are crucial factors for rectal cancer staging and surgical planning. Narrow pelvic dimensions may complicate surgical access and influence resection margins and oncological outcomes^{7,8}. Our study's findings regarding PI/ML ratio align with these anatomical considerations, suggesting that the PI/ML ratio might serve as a quantitative reflection of pelvic complexity in rectal cancer surgery.

In our series, we found a significant correlation between PI/ML ratio and tumor recurrence. This supports the hypothesis that pelvic anatomy, as reflected in pelvimetric measurements, may influence oncological outcomes. For tumor recurrence, a PI/ML cutoff value of < 1.6 was identified, with high sensitivity and specificity. Our findings suggest that pre-operative pelvic measurements could assist in risk stratification for tumor recurrence in rectal cancer patients. This finding is consistent with recent imaging literature emphasizing how mesorectal involvement and pelvic constraints may influence local recurrence risk¹¹. However, Huang et al. found no

significant difference between radiologically measured pelvic parameters and recurrence and mortality¹³.

A significant association was also observed between the PI/ML ratio and anastomotic leakage. In patients with a PI/ML ratio < 2.15, the sensitivity and specificity for predicting anastomotic leakage were 100% and 53.7%, respectively. While pelvic anatomy has long been recognized as an important factor in surgical difficulty^{11,12}, our study contributes quantitative data supporting its association with post-operative complications. Accordingly, the PI/ML ratio can be used in determining the surgical strategy and may be of clinical benefit in applications such as controlling anastomotic tension and monitoring anastomotic viability.

Our findings regarding LARS differ from earlier assumptions that pelvic anatomy directly predicts functional outcomes. Although we observed a trend toward lower PI/ML ratios in patients with LARS, this did not reach statistical significance. This is in line with a recent systematic review and meta-analysis, which reported a pooled incidence of major LARS of approximately 44%⁸. Sun et al. highlighted that the major risk factors for LARS include neoadjuvant radiotherapy, low tumor height, low anastomotic level, anastomotic leakage, and diverting stomas, rather than purely anatomical parameters such as pelvic dimensions. Thus, while pelvic anatomy may play a role in surgical difficulty, its direct correlation with post-operative functional disorders remains uncertain⁸.

Recent evidence suggests that BMI may influence oncological outcomes in rectal cancer patients. Although BMI was not independently associated with the PI/ML ratio in our cohort, the post-operative LARS rate in patients with high BMI was statistically significant in our study. High BMI values cause an increase in mesorectal adipose tissue thickness and increase the technical difficulty of TME. This finding supports the results of the study by Meng et al.¹⁴. Liu et al. demonstrated that lower BMI, particularly BMI < 18.5 kg/m², was significantly associated with poorer overall and cancer-specific survival in patients undergoing neoadjuvant chemoradiotherapy followed by TME¹⁵. This highlights that not only anatomical factors but also patient-related biological characteristics may contribute to prognosis in rectal cancer¹⁶.

In the literature, Boyle et al. found that pelvic dimensions vary by sex in healthy individuals. The antero-posterior diameter of the mesorectum and the volume of the fat area were evaluated radiologically, and it was found that the mesorectal area was wider in men and the anterior mesorectal fat was thinner in women¹⁶.

In our study, there was no significant difference in PI/ML between male and female patients, but BMI values were higher in female patients compared to male patients. These BMI differences could be relevant when planning treatment, as higher BMI may influence technical difficulty during surgery and post-operative outcomes. However, our study did not find a direct link between BMI and PI/ML ratio or post-operative complications, possibly due to the limited sample size.

Our findings align with recent research highlighting the role of pelvic anatomy in rectal cancer surgery. A study by Aparicio-López et al. found that specific pelvimetric measurements correlated significantly with surgical difficulty and post-operative complications, including anastomotic leakage and conversion rates during rectal cancer surgery¹⁷. Similarly, another review emphasized that variations in pelvic anatomy may influence not only technical aspects of TME but also functional outcomes such as LARS¹⁸. These reports support the potential clinical utility of anatomical indices such as the PI/ML ratio as prognostic tools for surgical planning and risk stratification in rectal cancer.

Our study has several limitations that should be acknowledged. First, it was a retrospective study conducted at a single center, which may introduce selection bias and limit the generalizability of the findings. Second, the sample size was relatively small, potentially reducing the statistical power to detect associations, especially regarding LARS outcomes. This sample size, though reflective of all eligible patients during the study period, may be underpowered to detect smaller effect sizes, particularly for outcomes such as LARS. Third, the assessment of LARS was performed via telephone interviews rather than in-person evaluations, which could introduce recall bias or variability in patient responses. Additionally, we did not assess interobserver variability for PI measurements, which may affect the reproducibility of our findings. Despite these limitations, our study provides valuable preliminary evidence on the potential role of PI/ML ratio in predicting oncological and surgical outcomes in rectal cancer patients. Future prospective multicenter studies with larger patient cohorts are needed to validate and expand upon these findings.

Conclusions

In our study, we found that the PI/ML ratio was significantly associated with anastomotic leakage and tumor recurrence in patients undergoing TME for rectal cancer. Specifically, lower PI/ML ratios were

predictive of higher risk for both anastomotic leakage and tumor recurrence, with high sensitivity and moderate to high specificity. However, we did not observe a statistically significant association between the PI/ML ratio and complete mesorectal excision status or the development of LARS. Although patients with LARS had lower mean PI/ML ratios, this difference did not reach statistical significance in our analysis.

These findings suggest that pre-operative pelvic measurements and the PI/ML ratio could be useful for predicting certain oncological and surgical outcomes, such as anastomotic leakage and recurrence risk. Further research with larger cohorts is warranted to validate these findings and explore the potential of PI/ML as a prognostic tool in rectal cancer surgery.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of humans and animals. The authors declare that the procedures followed complied with the ethical standards of the responsible human experimentation committee and adhered to the World Medical Association and the Declaration of Helsinki. The procedures were approved by the institutional Ethics Committee.

Confidentiality, informed consent, and ethical approval. The authors have obtained approval from the Ethics Committee for the analysis of routinely obtained and anonymized clinical data, so informed consent was not necessary. Relevant guidelines were followed.

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing of this manuscript.

References

1. World Health Organization (WHO). Colorectal cancer. Fact Sheets; 2020. Available from: <https://www.who.int/news-room/fact-sheets/detail/colorectal-cancer>
2. Kawai K, Sugihara K, Sakai Y, Morita S, Yamaguchi T, Kanemitsu Y, et al. Impact of pelvic dimensions on surgical difficulty in rectal cancer surgery. *World J Surg.* 2021;45:648-57.
3. Brock H, Lambrineas L, Ong HI, Chen WY, Das A, Edsell A, et al. Preventative strategies for low anterior resection syndrome. *Tech Coloproctol.* 2023;28:10.

4. Rosen H, Sebesta CG, Sebesta C. Management of low anterior resection syndrome (LARS) following resection for rectal cancer. *Cancers (Basel)*. 2023;15:778.
5. Ye L, Huang M, Huang Y, Yu K, Wang X. Risk factors of postoperative low anterior resection syndrome for colorectal cancer: a meta-analysis. *Asian J Surg*. 2022;45:39-50.
6. Serra-Aracil X, Zarate A, Bargalló J, Gonzalez A, Serracant A, Roura J, et al. Transanal versus laparoscopic total mesorectal excision for mid and low rectal cancer (Ta-LaTME study): multicentre, randomized, open-label trial. *Br J Surg*. 2023;110:150-8.
7. Keane C, Fearnhead NS, Bordeianou LG, Christensen P, Basany EE, Laurberg S, et al. International consensus definition of low anterior resection syndrome. *Dis Colon Rectum*. 2020;63:274-84.
8. Sun R, Dai Z, Zhang Y, Lu J, Zhang Y, Xiao Y. The incidence and risk factors of low anterior resection syndrome (LARS) after sphincter-preserving surgery of rectal cancer: a systematic review and meta-analysis. *Support Care Cancer*. 2021;29:7249-58.
9. Tsuruta A, Tashiro J, Ishii T, Oka Y, Suzuki A, Kondo H, et al. Prediction of anastomotic leakage after laparoscopic low anterior resection in male rectal cancer by pelvic measurement in magnetic resonance imaging. *Surg Laparosc Endosc Percutan Tech*. 2017;27:54-9.
10. Emmertsen KJ, Laurberg S. Low anterior resection syndrome score: development and validation of a symptom-based scoring system for bowel dysfunction after low anterior resection for rectal cancer. *Ann Surg*. 2012;255:922-8.
11. Bogveradze N, Snaebjornsson P, Grotenhuis BA, Van Triest B, Lahaye MJ, Maas M, et al. MRI anatomy of the rectum: key concepts important for rectal cancer staging and treatment planning. *Insights Imaging*. 2023;14:13.
12. Inoue A, Sheedy SP, Wells ML, Mileto A, Goenka AH, Ehman EC, et al. Rectal cancer pelvic recurrence: imaging patterns and key concepts to guide treatment planning. *Abdom Radiol (NY)*. 2023;48:1867-79.
13. Huang B, Liu MC, Gao W, Tang J, Zhu Z, Chen L, et al. Nomogram for predicting the feasibility of natural orifice specimen extraction after laparoscopic rectal resection. *J Gastroenterol Hepatol*. 2021;36:1803-11.
14. Meng L, Qin H, Huang Z, Liao J, Cai J, Feng Y, et al. Analysis of presacral tissue structure in LARS and the prevention of LARS by reconstruction of presacral mesorectum with pedicled greater omentum flap graft. *Tech Coloproctol*. 2021;25:1291-300.
15. Liu H, Wei R, Li C, Zhao Z, Guan X, Yang M, et al. BMI may be a prognostic factor for local advanced rectal cancer patients treated with long-term neoadjuvant chemoradiotherapy. *Cancer Manag Res*. 2020;12:10321-32.
16. Boyle KM, Chalmers AG, Finan PJ, Sagar PM, Burke D. Morphology of the mesorectum in patients with primary rectal cancer. *Dis Colon Rectum*. 2009;52:1122-9.
17. Aparicio-López D, Gascón-Ferrer I, Martínez-Germán A, Santero-Ramírez MP, Sánchez-Fuentes MN, Gracia-Roche C, et al. Aplicación del índice pronóstico REAL-score en la toma de decisiones en cirugía del cáncer de recto. *Cir Cir*. 2023;91:690-7.
18. Sun W, Zhang J. Comparison of clinical efficacy of different colon anastomosis methods in laparoscopic radical resection of colorectal cancer. *Cir Cir*. 2024;92:314-23.

Association of NRF-2 changes in plasma and pericardial fluid with renal injury in patients undergoing cardiac surgery

Asociación de los cambios de NRF-2 en plasma y líquido pericárdico con lesión renal en pacientes sometidos a cirugía cardíaca

Mahmut Padak^{1*}, Reşat Dikme², Ebru Temiz³, Ömer Göç⁴, Ezhar Ersöz⁵, and Yasemin Hacınlı⁵

¹Department of Perfusion Techniques, Harran University Vocational School of Health Services; ²Department of Dialysis Program, Harran University Vocational School of Health Services; ³Harran University Vocational School of Health Services; ⁴Department of Biomedical Device Technology, Harran University Vocational School of Health Services; ⁵Harran University, Research and Application Hospital. Şanlıurfa, Turkey

Abstract

Objective: This study was conducted to investigate the levels of nuclear factor (erythroid derivative 2)-like 2 (NRF-2), kidney injury molecule 1 (KIM-1), and heme oxygenase-1 (HO-1) in pericardial fluid (PF) and systemic circulation of patients undergoing cardiac surgery. **Methods:** This study included 40 patients undergoing cardiac surgery and 40 healthy individuals. PF and venous blood samples were obtained from the patients and renal function tests, HO-1, KIM-1, NRF-2, antioxidant, and oxidative stress parameters were studied. **Results:** A statistically significant difference was found in the NRF-2, KIM-1, HO-1, total antioxidant status, total oxidant status, and oxidative stress index measurements in the plasma of the patient and control groups and in the PF of the patients ($p < 0.01$). Compared to the control group, NRF-2, KIM-1, and HO-1 were found to be lower in the patient's plasma and PF. In the patient group, NRF-2, KIM-1, HO-1, and KIM-1 were higher in PF compared to plasma. **Conclusions:** In the future, intrapericardial drug administration may improve cardiac function and prevent the adverse cardiorenal syndrome on the kidney.

Keywords: Kidney damage. Nuclear factor (erythroid derivative 2)-like 2. Kidney injury molecule 1. Heme oxygenase-1. Cardiopulmonary bypass. Cardiac surgery.

Resumen

Objetivo: Investigar los niveles del factor nuclear (derivado eritroide 2) similar 2 (NRF-2), la molécula de lesión renal 1 (KIM-1) y la hemooxigenasa 1 (HO-1) en el líquido pericárdico y en la circulación sistémica de pacientes sometidos a cirugía cardíaca. **Métodos:** En este estudio participaron 40 pacientes sometidos a cirugía cardíaca y 40 individuos sanos. Se obtuvieron muestras de líquido pericárdico y sangre venosa de los pacientes y se estudiaron las pruebas de función renal, HO-1, KIM-1, NRF-2 y los parámetros antioxidantes y de estrés oxidativo. **Resultados:** Se encontró una diferencia estadísticamente significativa en las mediciones de NRF-2, KIM-1, HO-1, estado antioxidante total (TAS), estado oxidante total (TOS) e índice de estrés oxidativo (OSI) en el plasma entre los grupos de pacientes y de sujetos control, y en el líquido pericárdico de los pacientes ($p < 0.01$). En comparación con el grupo de control, se observó que NRF-2, KIM-1 y HO-1 eran más bajos en el plasma y en el líquido pericárdico de los pacientes. En el grupo de pacientes, NRF-2, KIM-1 HO-1 y KIM-1 eran más elevados en el líquido pericárdico que en el plasma. **Conclusiones:** En el futuro, la administración intrapericárdica de fármacos puede mejorar la función cardíaca y prevenir el síndrome cardiorenal adverso sobre el riñón.

Palabras clave: Daño renal. NRF-2. KIM-1. HO-1. Bypass cardiopulmonar. Cirugía cardíaca.

*Correspondence:

Mahmut Padak
E-mail: mpadak@harran.edu.tr

Date of reception: 25-04-2024
Date of acceptance: 06-08-2024
DOI: 10.24875/CIRUE.M24000896

Cir Cir (Eng). 2025;93(6):591-599
Contents available at PubMed
www.cirugiaycirujanos.com

2444-0507/© 2024 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Many organs, especially the heart and kidney, are affected in cardiopulmonary bypass (CPB), which allows cardiovascular surgery to be performed. Many molecules are secreted from the organs affected by this condition. Many of these secreted molecules can provide important information about the relevant organ or disease by showing biomarker properties. Recently, apart from classical kidney biomarkers such as urea, uric acid, and creatinine, such as nuclear factor (erythroid derived 2)-like 2 (NRF-2), kidney damage molecule 1 (KIM-1), and heme oxygenase-1 (HO-1) new biomarkers have been discovered.

NRF-2 is an important protein that regulates the expression of antioxidant proteins that protect against oxidative damage triggered by injury and inflammation¹. As long as it is not exposed to reactive oxygen species, NRF-2, which is degraded, passes to the nucleus without being degraded under oxidative stress, binds to DNA and initiates the transcription of antioxidant genes^{2,3}. NRF-2, which provides the stimulation of many cytoprotective proteins, is found in the kidney, as well as in the muscle, lung, heart, liver, and brain^{4,5}.

HO-1, an enzyme synthesized in renal tubular cells, has cell protective, anti-inflammatory, and antioxidant properties and is synthesized in response to oxidative stress⁶. HO-1, whose transcriptional regulation is dependent on NRF-2, plays a role in many diseases related to endoplasmic reticulum stress, such as cardiovascular and metabolic diseases^{7,8}.

In some studies, kidney injury molecule 1 (KIM-1) was found to be significantly higher in patients who developed kidney damage (KD) after cardiac surgery⁹. In damaged kidney cells, KIM-1 functions as a receptor controlling apoptotic cell phagocytosis¹⁰. After studies on KD, the American Food and Drug Administration and the European Medicines Agency approved KIM-1, one of the seven new biomarkers to be used for the detection of nephrotoxicity¹¹. Many studies have supported that KIM-1 is a useful biomarker for the early detection and diagnosis of complications after cardiac surgery and KD caused by ischemia¹².

Although these biomarkers such as NRF-2, KIM-1, and HO-1 were investigated in human plasma, they were not investigated in pericardial fluid (PF). PF has an important place in cardiovascular research due to its anatomical proximity to cardiac tissue and rich content¹³. Changes in molecular markers in PF show strong correlations with many structural and functional

parameters as well as cardiac pathophysiologies. The pericardial environment is both a potential diagnostic biomarker pool and an important location for future treatment and interventions. Due to the lack of data on the physiological composition of PF in studies conducted to date, its clinical use has been limited¹⁴.

In our study on the change of these biomarkers, which provide valuable information in the early detection of KD, during CPB, the level of the relevant parameters in the plasma was monitored and information was obtained about the level in PF.

Methods

Ethical committee approval

The present study was approved by the local ethics committee (Approval number: 29.06.2020-20.12.26).

Patients included in the study

This study was conducted in accordance with the Helsinki Declaration, which was revised in 1989. Forty patients who underwent cardiac surgery with CPB method and 40 healthy individuals as control group were included in this study. PF and venous blood samples were obtained from the patients included in the study. Only venous blood samples were taken from the control group.

Obtaining PF

After median sternotomy was performed with standard CPB procedures in patients who underwent cardiac surgery with the CPB method, the pericardium was opened and PF aspiration was provided with a sterile syringe. The aspirated PF (5-20 mL) was then taken into sterile tubes without anticoagulant, and the sterile tube was immediately transferred into an ice-filled container. Then, the PF in a sterile tube delivered to the laboratory with an ice-filled container was centrifuged at + 4°C at 5000 rpm for 5 min, and the supernatant was taken into an RNase-free tube (ependorf tube) and stored at -80°C to be studied.

Obtaining blood plasma

In patients who underwent cardiac surgery with the CPB method, blood was drawn before, during and after CPB and put into sterile tubes with anticoagulant (heparin). The tube from which blood was drawn was

immediately transferred to an ice-filled container and transported to the laboratory. The sterile tube was then centrifuged at 5000 rpm for 5 min. After the centrifugation step, the plasma, which is the supernatant part, was taken into an RNase-free tube (ependorf tube) and stored at -80 to be studied.

Biochemistry study in plasma of patient group

Glucose, urea, albumin, alanine aminotransferase (ALT), Na, creatinine, aspartate aminotransferase (AST), direct bilirubin, gamma-glutamyl transferase (GGT), calcium (Ca), and C-reactive protein (CRP) levels were measured in the plasma of the patients included in the study before the operation (preop), after the termination of CBP and after the operation (post-operative). Siemens Atellica autoanalyzer device (USA, 2018) was used to measure these parameters.

Enzyme-linked immunosorbent assay (ELISA) study in PF and plasma

PF AND PLASMA NRF-2 ELISA STUDY

Human NFE2L2 (Nuclear Factor, Erythroid Derived 2, Like 2) ELISA Kit (Elabscience, USA, Catalog No: E-EL-H1564) was used for NRF-2 ELISA study in PF and Plasma. This kit used works according to the double-antibody sandwich enzyme-linked immunosorbance analysis to test the NRF-2 level in biological materials. The sensitivity (sensitivity) of the kit is 0.10 ng/mL, and the assay range (Detection Range) is 0.16-10 ng/mL.

PF AND PLASMA HO-1 ELISA STUDY

Human HO-1 ELISA Kit (Elabscience, USA, Cat. No: E-EL-H2172) was used for HO-1 ELISA assay in PF and plasma. This kit used works according to the double-antibody sandwich enzyme-linked immunosorbance assay to test the HO-1 level in biological materials. The sensitivity of the kit is 0.19 ng/mL, and the assay range is 0.31-20 ng/mL.

HUMAN KIM-1 ELISA STUDY IN PF AND PLASMA

Human KIM-1 ELISA Kit (Elabscience, USA, Catalog No: E-EL-H6029) was used for the KIM-1 ELISA study in PF and plasma. This kit used works according

to the double-antibody sandwich enzyme-linked immunosorbance assay to test the level of KIM-1 in biological materials. The sensitivity of the kit is 4.69 pg/mL, and the assay range (Detection Range) is 7.81-500 pg/mL.

Measurement of antioxidant and OS parameters in PF and plasma

TOTAL ANTIOXIDANT STATUS (TAS) MEASUREMENT

Measurement of (TAS in PF and plasma was performed using Rel Assay Diagnostics total oxidant capacity measurement kit (Rel Assay Diagnostics, Lot. No: HN20106A, Turkey). Plasma TAS levels were determined using a new automated measurement method developed by Erel¹⁵.

TOTAL OXIDANT STATUS (TOS) MEASUREMENT

Rel assay diagnostics total oxidant capacity measurement kit (Rel Assay Diagnostics, Lot. No: OK20115O, Turkey) was used for the measurement of TOS in PF and plasma. Plasma TOS levels were determined using a new automated measurement method developed by Erel¹⁶.

OXIDATIVE STRESS INDEX (OSI) MEASUREMENT

OSI was calculated by dividing it as total oxidant level (TOS)/total antioxidant and expressed as arbitrary unit (AU).

Statistical analysis

The Statistical Package for the Social Sciences (SPSS) Windows version 24.0 (SPSS, Inc, Chicago, USA) package program was used for statistical analysis. The conformity of the data to the normal distribution was tested with the Shaphiro Wilk test. Mann-Whitney U test was used for comparison of numerical variables in two independent groups, one-way analysis of variance and least significant difference multiple comparison tests were used for normally distributed features, and Kruskal-Wallis test and all pairwise multiple comparison test were used for non-normally distributed features. As descriptive statistics, mean \pm standard deviation for numerical variables, number, and percentage values for categorical

variables are given. $p < 0.05$ were considered statistically significant.

Results

Demographic data of the working group

Of the patients included in the study, 25 were male and 15 were female, with a mean age of 62.58 (years), height 160.11 (cm), weight 75.40 (kg), and body surface area (BSA) 1.83 that it is calculated as (m^2).

Biochemical results in plasma of patients

The results of biochemical examinations in the plasma of the patients included in the study before the operation (preop), after CBP and after the operation (postop) are given in table 1.

ELISA results in PF and plasma

The results of the optical density reading at 450 nm using commercial kits for the level determination of HO-1, NRF-2, and KIM-1 in PF and plasma by ELISA method and the results of OS parameters are given in table 2.

Table 2 shows the statistical analysis of different biochemical parameters in pre-operative, post-CBP, and post-operative groups. According to this analysis, glucose, urea, creatinine, albumin, ALT, AST, direct bilirubin, GGT, Na, Ca, and CRP values were statistically significant (* $p < 0.05$, ** $p < 0.001$).

Glucose value which was 170.23 before CPB was 201.30 after CPB and 177 post-operative ($p < 0.05$). Urea value which was 37.22 before CPB was 42.07 after CPB and 47.41 post-operative ($p < 0.01$). Creatinine value was 0.91 before CPB, 0.98 after CPB and 1.17 post-operative ($p < 0.01$). Albumin value which was 41.27 before CPB decreased to 29.43 after CPB and increased to 31.22 postop ($p < 0.01$). While ALT value was 21.26 before CPB, it became 25.89 after CPB and 28.32 post-operative ($p < 0.01$). AST value was 29.12 before CPB, 49.22 after CPB and 50.30 post-operative ($p < 0.01$). D. bilirubin value which was 0.2 before CPB was 0.15 after CPB and 0.26 post-operative ($p < 0.01$). GGT value which was 27.58 before CPB decreased to 24 after CPB and increased to 29.83 post-operative ($p < 0.05$). Na value, which was 138.20 before CPB, became 141.78 after CPB and 141.83 post-operative ($p < 0.01$). Ca value which

was 9.17 before CPB was 8.16 after CPB and 7.85 post-operative ($p < 0.01$). CRP value, which was 13.38 before CPB, was 25.4 after CPB and 132.21 post-operative ($p < 0.01$) (Fig. 1).

As shown in Graph 1, when HO-1, NRF-2, KIM-1, TAS, TOS, and OSI values in the plasma and PF of the patients and the plasma of the control group were compared, the highest HO-1 value was found in the plasma of the control group, while the lowest HO-1 value was found in the plasma of the patients. The highest NRF-2 value was found in the plasma of the control group, while the lowest NRF-2 value was found in the patient plasma. The highest KIM-1 value was found in the plasma of the control group, while the lowest KIM-1 value was found in the PF of the patients. The highest TAS value was found in the plasma of the control group, while the lowest TAS value was found in the PF of the patients. The highest TOS value was found in the PF of the patients, while the lowest TOS value was found in the plasma of the control group. The highest OSI value was found in the PF of the patients, while the lowest OSI value was found in the plasma of the control group.

These results were statistically significant ($p < 0.001$). According to these results, HO-1, NRF-2, KIM-1, and TAS values were lower in the plasma of the patients compared to the plasma of the control group, while TOS and OSI values were higher. When the PF was compared with the plasma of the patients, HO-1, NRF-2, TOS, and OSI values in the PF were higher than the plasma of the patients, while KIM-1 and TAS values were lower. HO-1, NRF-2, KIM-1, and TAS values were found to be lower in pericardium, before CPB - plasma, during CPB - plasma and after CPB-plasma compared to the control group. TOS and OSI values were found to be higher in pericardium, before CPB - plasma, during CPB - plasma and after CPB-plasma compared to the control group.

According to table 3, there is a significant positive relationship between HO-1 value and NRF-2, TOS, and OSI value. There is a significant positive relationship between NRF-2 value and KIM-1, TOS, and OSI value, and a significant negative relationship between TAS value. There is a moderately significant negative relationship between TAS value and TOS and OSI value. There is a very highly significant positive relationship between TOS value and OSI value.

As shown in figure 2, the lowest HO-1 value in patient plasma was found in Pump 1, while the highest HO-1 value was found in Pump 3. The lowest NRF-2 value was found in Pump 1, while the highest NRF-2

Table 1. Biochemical test results of the patients included in the study

Parameter	Preop CPB	After CPB	Postop	p
Plasma	Mean ± SS	Mean ± SS	Mean ± SS	
Glucose (mmol/L)	170.23 ± 70.33	201.3 ± 47.78	177 ± 41.43	0.013*
Urea (mg/dL)	37.22 ± 15.29	42.07 ± 14.98	47.41 ± 19.5	0.004*
Creatinine (mg/dL)	0.91 ± 0.71	0.98 ± 0.42	1.17 ± 0.72	0.003*
Albumin (g/dL)	41.27 ± 4.95	29.43 ± 5.8	31.22 ± 5.21	0.001*
ALT (U/L)	21.26 ± 12.18	25.89 ± 13.88	28.32 ± 11.63	0.007*
AST (birim/L)	29.12 ± 20.11	49.22 ± 28.17	50.3 ± 24.34	0.001*
Total bilirubin (mg/dL)	0.55 ± 0.31	0.7 ± 0.42	0.59 ± 0.32	0.102
Direct bilirubin (mg/dL)	0.2 ± 0.09	0.15 ± 0.1	0.26 ± 0.14	0.001*
GGT (U/L)	27.58 ± 14.92	24 ± 13.81	29.83 ± 13.31	0.043*
Na (mEq/L)	138.2 ± 3.15	141.78 ± 3.08	141.83 ± 4.29	0.001*
K (mEq/L)	4.35 ± 0.35	4.46 ± 0.46	4.27 ± 0.47	0.203
Ca (mEq/L)	9.17 ± 0.68	8.16 ± 0.71	7.85 ± 0.59	0.001*
CRP (mg/L)	13.38 ± 7.25	25.4 ± 15.75	132.21 ± 78.59	0.001*

*Kruskal-Wallis Test.

*p < 0.05; **p < 0.001.

ALT: alanine aminotransferase; AST: aspartate aminotransferase; CBP: cardiopulmonary bypass; GGT: gamma-glutamyl transferase; Na: sodium; K: potassium; Ca: calcium; CRP: C-reactive protein.

Table 2. HO-1, NRF-2, KIM-1 ELISA results, and OS parameters in PF and plasma.

Parameter	Control (Plasma)	Pericardial fluid	Preop CPB (Plasma)	After CPB (Plasma)	Postop (Plasma)	p
	Mean ± SS	Mean ± SS	Mean ± SS	Mean ± SS	Mean ± SS	
HO-1 (ng/mL)	1.35 ± 0.32	0.66 ± 0.22	0.58 ± 0.16	0.72 ± 0.28	1.12 ± 0.51	0.000
NRF-2 (ng/mL)	5.63 ± 1.26	0.85 ± 0.33	0.77 ± 0.24	4.07 ± 1.71	4.55 ± 1	0.000
KIM-1 (pg/mL)	31.54 ± 17.09	18.7 ± 9.92	26.11 ± 14.8	27.69 ± 13.81	30.66 ± 19.12	0.001
TAS (Trolox equivalent/L)	1.57 ± 0.21	1.19 ± 0.35	1.45 ± 0.17	1.34 ± 0.25	1.34 ± 0.19	0.000
TOS (µmol H ₂ O ₂ Eqv./L)	10.79 ± 1.99	17.78 ± 2.48	14.46 ± 2.32	14.96 ± 2.52	16.55 ± 2.76	0.000
OSI (Arbitrary Unit (AU))	0.71 ± 0.19	1.69 ± 0.8	1.05 ± 0.25	1.1 ± 0.24	1.28 ± 0.34	0.000

*Kruskal-Wallis Test.

*p < 0.05; **p < 0.001.

HO-1: heme oxygenase-1; NRF-2: nuclear factor (erythroid derivative 2)-like 2; KIM-1: kidney injury molecule 1; TAS: total antioxidant status; TOS: total oxidant status; OSI: oxidative stress index.

value was found in Pump 3. The lowest KIM-1 value was in Pump 1, while the highest KIM-1 value was in Pump 3. The highest TAS value was in Pump 1, while the lowest TAS value was in Pump 3. The lowest TOS value was in Pump 1, while the highest TOS value was in Pump 3. The lowest OSI value was found in Pump 1, while the highest OSI value was found in Pump 3.

These results were statistically significant ($p < 0.001$). According to these results, HO-1, NRF-2, KIM-1, TOS, and OSI values increased and TAS value decreased ($p < 0.001$).

There is a significant positive correlation between HO-1 value and NRF-2, TOS, and OSI value. There is a positive relationship between NRF-2 value and

Table 3. Correlation of HO-1, NRF-2, KIM-1, TAS, TOS, and OSI values in patients included in the study

Parameter	HO-1	NRF-2	KIM-1	TAS	TOS	OSI
HO-1						
r	1.000	0.543**	0.104	-0.139	0.431**	0.375**
p		0.000	0.158	0.059	0.000	0.000
NRF-2						
r	0.543**	1.000	0.172*	-0.274**	0.508**	0.511**
p	0.000		0.020	0.000	0.000	0.000
KIM-1						
r	0.104	0.172*	1.000	0.147	-0.083	-0.140
p	0.158	0.020		0.05	0.263	0.059
TAS						
r	0.139	0.274**	0.147	1.000	-0.274**	-0.740**
p	0.059	0.000	0.05		0.000	0.000
TOS						
r	-0.431**	-0.508**	-0.083	-0.274**	1.000	0.819**
p	0.000	0.000	0.263	0.000		0.000
OSI						
r	-0.375**	-0.511**	-0.140	-0.740**	0.819**	1.000
p	0.000	0.000	0.059	0.000	0.000	

*p < 0.05.

**p < 0.01.

HO-1: heme oxygenase-1; NRF-2: nuclear factor (erythroid derivative 2)-like 2; KIM-1: kidney injury molecule 1; TAS: total antioxidant status; TOS: total oxidant status; OSI: oxidative stress index.

KIM-1, TOS, and OSI values, and a negative significant relationship between TAS values. There is a moderately significant negative correlation between TAS value and TOS and OSI values. There is a highly significant positive correlation between TOS value and OSI value.

Discussion

One of the most important elements to be evaluated during CPB is the follow-up of irreversible cell and tissue damage. Before reaching this stage, the cell or tissue synthesizes biomarkers that provide information about its state. For example, in KD, NRF-2, HO-1, and KIM-1 are activated and provide information about the organ. Therefore, these parameters should be monitored during CPB and measures should be taken accordingly.

In our study, a positive correlation was found between urea, creatinine, and CRP values in the plasma of the patient group and HO-1, KIM-1, and NRF-2 values. According to this result, CPB had a negative effect on the kidneys and caused an increase in urea, creatinine, CRP, HO-1, KIM-1, and NRF-2.




Figure 1. Graphical display of heme oxygenase-1, nuclear factor (erythroid derivative 2)-like 2, kidney injury molecule 1, total antioxidant status, total oxidant status, and oxidative stress index values in plasma and pericardial fluid of patients and plasma of control group.




Figure 2. Graphical representation of the changes in the heme oxygenase-1, nuclear factor (erythroid derivative 2)-like 2, kidney injury molecule 1, total antioxidant status, total oxidant status and oxidative stress index values in the plasma of the patients during before cardiopulmonary bypass (CPB) - plasma, during CPB-plasma and after CPB-plasma.

In the study conducted by Magyar et al., (2019) on HO-1 and KD following cardiac surgical procedure with CPB technique, an increase in HO-1 was observed 24 h after CPB in patients who developed acute kidney injury (AKI)¹⁷. In another study, it was reported that AKI was associated with duration of CPB, hemolysis, inflammation, and increased HO-1 concentrations in patients after cardiac surgery¹⁸. In a different study, it was found that HO-1 reduces inflammation and oxidative stress, and carbon monoxide, the product of HO-1 degradation, reduces KD caused by CPB¹⁹. In our study, the HO-1 value, which was lower in plasma and PF of the patients compared to the control group, increased continuously during the CPB-pump (p < 0.001). In our study, the HO-1 value was higher in PF compared to the plasma of the same patient (p < 0.001). The detection of HO-1 in PF supports the view that PF is a plasma ultrafiltrate. In addition, the

detection of HO-1 in PF shows that this molecule is not only synthesized by the kidney but also by the heart tissue. A high HO-1 level during the CPB-pump period indicates that the kidney functions are affected and respond.

Some researchers believe that high KIM-1 levels can be used as an early diagnostic indicator of AKI in patients after cardiac surgery²⁰ and correlate with renal impairment²¹. Neyra et al., in a study of KIM-1, 23 of 106 adult patients who underwent coronary artery bypass graft (CABG) and valve surgery with CPB developed KD. In these patients, urine KIM-1/creatinine combination was found to be significantly above the optimal threshold value with post-operative KD²². The KIM-1 value, which was lower in plasma and PF of the patients compared to the control group, increased during the CPB process ($p < 0.001$). In our study, KIM-1 value was lower in PF compared to the plasma of the same patient ($p < 0.001$). The fact that KIM-1 was lower in PF compared to plasma indicates that this molecule is mostly synthesized in the kidneys.

In our study, the NRF-2 value, which was lower in plasma and PF of the patients compared to the control group, increased continuously during the pump ($p < 0.001$). In the same patient, NRF-2 value was higher in PF than in plasma ($p < 0.001$). While a negative significant relationship was found between NRF-2 and TAS, a positive significant relationship was found between TOS and OSI. NRF-2 value also increased in parallel with increasing OS. In this respect, NRF-2 is similar to other studies in the literature. As can be seen from the studies, promoting NRF-2-mediated protection against OS and inflammation-induced damage with different bioactive compounds may represent new therapeutic strategies. In our study, the increase of NRF-2 against the increased cellular stress and inflammatory response due to CPB was parallel to other studies in the literature.

Although NRF-2, HO-1, and KIM-1 provide important information in KD, it would not be correct to evaluate only these. Because the origin of the damage to the kidneys during CPB is multifactorial. Although CPB is the main factor in this damage, changes in hematocrit and temperature, use of vasoactive and nephrotoxic drugs, hemodynamic changes, and low cardiac output also have negative effects on the kidneys. It is interesting that some studies have reported that the incidence of KD is similar during off-pump CABG and on-pump CABG²³. Therefore, the specific damage of CPB to the kidneys has not been clarified. However,

according to studies, KD in patients with normal pre-operative renal function has a relatively low incidence compared to patients with pre-operative dysfunction. In addition, even in a patient with normal kidney function, the effect of the slightest physiological change in the heart can be felt in the kidney due to the effect of cardiorenal syndrome.

From the past to the present, different groups have tried to find treatments that can protect the kidney from damage during cardiac and non-cardiac surgery. These solutions include the use of drugs such as dopamine, furosemide, fenoldopan, atrial natriuretic factor, sodium bicarbonate, n-acetyl cysteine, enoximone, and dexmedetomidine. In addition, hypothermia and normothermia applications with high or low hematocrit during CPB are other applications to protect the kidney. However, no clear treatment or method has been found to date²⁴.

A 2012 study used ulinastatin, a protease inhibitor that can suppress the harmful effects of inflammatory reactions and free radicals on kidney cells during CPB²⁵.

As can be understood from these studies, promoting NRF-2-mediated protection against OS and inflammation-induced damage with different bioactive compounds may represent new therapeutic strategies. In our study, the high level of NRF-2 against increased cellular stress and inflammatory response due to CPB was in parallel with other studies in the literature. In particular, the level of NRF-2 increased rapidly in response to ischemia/reperfusion after cross-clamping of the aorta during CPB. This shows that NRF-2 responds rapidly to cellular reactions and is a diagnostically important parameter. It is important to follow the NRF-2 level especially in the detection of BD that may occur during CPB and to get support from the nephrology clinic to make the necessary treatment according to the level. During CPB, tissue perfusion may also be impaired due to pump pressure and ischemia/reperfusion injury may occur especially in the kidney due to insufficient blood supply. In such cases, NRF-2 may protect the kidney from damage by supporting antioxidant capacity.

Conclusions

In our study population, urea, creatinine, CRP, HO-1, NRF-2, KIM-1, TOS, and OSI increased continuously in correlation with each other during CPB and postoperatively. Compared to the plasma of the control group, HO-1, NRF-2, KIM-1, and TAS were found to be lower in the plasma of the patient group, while TOS

and OSI were found to be higher. In the patient group, HO-1, NRF-2, TOS, and OSI were higher in PF compared to plasma, while KIM-1 and TAS values were lower. It is important that all molecules in plasma were detected in PF for the first time in the study, and this supports the view that PF is a plasma ultrafiltrate. In addition, we can say that KIM-1, which is lower in PF compared to plasma, is a molecule released mostly by the kidneys.

In CPB, KD can be prevented or reduced with therapeutic strategies that reduce the adverse effects on the kidneys. For this, it is important to develop and implement the most appropriate strategies that increase NRF-2/HO-1 and decrease KIM-1 by following the relevant parameters during CPB and postoperatively.

Two different terms can be used to describe the pathophysiologic interaction between the heart and kidney. The first one, cardiorenal, is used when cardiac dysfunction causes renal dysfunction. The second term, renocardiac, is used when renal dysfunction causes cardiac dysfunction. Future studies may examine the relationship between cardiac and renal biomarkers to determine whether the interaction between the two organs is cardiac or renal (cardiorenal or renocardiac).

The fact that PF was studied and compared with plasma for the first time, that three important parameters such as HO-1, NRF-2, and KIM-1, which are related to each other, were studied together and that the data obtained were compatible with the clinic made our study powerful.

Limitations

Our study has some limitations. This was a prospective study and included a limited number of patients. We had limitations such as not being able to analyze NRF-2, HO-1 and KIM-1 in healthy individuals because PF could not be obtained from healthy individuals, not being able to analyze the relevant parameters in urine due to budget problems and not being able to perform gene expression studies. Our main aim in this study was to investigate the prognostic value of NRF-2. Different designed studies on pathophysiologic processes are needed to elucidate the mechanisms underlying our findings.

Funding

This study was supported by Harran University Scientific Research Projects Coordinatorship with project number 20102.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of humans and animals. The authors declare that the procedures followed complied with the ethical standards of the responsible human experimentation committee and adhered to the World Medical Association and the Declaration of Helsinki.

Confidentiality, informed consent, and ethical approval. The authors have followed their institution's confidentiality protocols, obtained informed consent from patients.

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing of this manuscript.

References

- Altin R, Kappos L, Arnold DL, Bar-Or A, Giovannoni G, Selmaj K, et al. "Placebo-controlled phase 3 study of oral BG-12 for relapsing multiple sclerosis". *N Engl J Med*. 2012;367:1098-107.
- Itoh K, Wakabayashi N, Katoh Y, Ishii T, Igarashi K, Engel JD, et al. Keap1 represses nuclear activation of antioxidant responsive elements by Nrf-2 through binding to the amino-terminal Neh2 domain. *Genes Dev*. 1999;13:76-86.
- Li W, Kong AN. Molecular mechanisms of NRF-2-mediated antioxidant response. *Mol Carcinog*. 2009;48:91-104.
- Tang W, Jiang YF, Ponnusamy M, Diallo M. Role of Nrf2 in chronic liver disease. *World J Gastroenterol*. 2014;20:13079-87.
- Kensler T, Wakabayashi N, Biswal S. Cell survival responses to environmental stresses via the Keap1-Nrf2-ARE pathway. *Annu Rev Pharmacol Toxicol*. 2007;47:89-116.
- Nitti M, Piras S, Brondolo L, Marinari UM, Pronzato MA, Furfaro AL. Heme oxygenase 1 in the nervous system: does it favor neuronal cell survival or induce neurodegeneration? *Int J Mol Sci*. 2018;8:2260.
- Dunn LL, Midwinter RG, Ni J, Hamid HA, Parish CR, Stocker R. New insights into intracellular locations and functions of heme oxygenase-1. *Antioxid Redox Signal*. 2014;20:1723-42.
- Sun Jang J, Piao S, Cha YN, Kim C. Taurine chloramine activates Nrf2, increases HO-1 expression and protects cells from death caused by hydrogen peroxide. *J Clin Biochem Nutr*. 2009;45:37-43.
- Yuan SM. Acute kidney injury after cardiac surgery: risk factors and novel biomarkers. *Braz J Cardiovasc Surg*. 2019;34:352-60.
- Ichimura T, Brooks CR, Bonventre JV. Kim-1/tim-1 and immune cells: shifting sands. *Kidney Int*. 2012;81:809-11.
- Harpur E, Ennulat D, Ho man D, Betton G, Gautier JC, Riefke B, et al. Nephrotoxicity. *Toxicol Sci*. 2011;122:235-52.
- Yuang Y, Don-Wauchope AC. The clinical utility of kidney injury molecule 1 in the prediction, diagnosis and prognosis of acute kidney injury: a systematic review. *Inflamm Allergy Drug Targets*. 2011;10:260-71.
- Trindade F, Vitorino R, Leite-Moreira A, Falcão-Pires I. Pericardial fluid: an underrated molecular library of heart conditions and a potential vehicle for cardiac therapy. *Basic Res Cardiol*. 2019;114:10.
- Yuan Z, Boulanger B, Flessner M, Johnston M. Relationship between pericardial pressure and lymphatic pericardial fluid transport in sheep. *Microvasc Res*. 2000;60:28-36.
- Erel O. A novel automated direct measurement method for total antioxidant capacity using a new generation, more stable ABTS radical cation. *Clin Biochem*. 2004;37:277-85.
- Erel O. A new automated colorimetric method for measuring total oxidant status. *Clin Biochem*. 2005;38:1103-11.
- Magyar A, Wagner M, Thomas P, Malsch C, Schneider R, Störk S, et al. HO-1 concentrations 24 hours after cardiac surgery are associated with the incidence of acute kidney injury: a prospective cohort study. *Int J Nephrol Renovasc Dis*. 2019;12:9-18.
- Billings FT, Yu C, Byrne JG, Petracek MR, Pretorius M. Heme oxygenase-1 and acute kidney injury following cardiac surgery. *Cardiorenal Med*. 2014;4:12-21.

19. Goebel U, Siepe M, Schwer CI, Schibilsky D, Foerster K, Neumann J, et al. Inhaled carbon monoxide prevents acute kidney injury in pigs after cardiopulmonary bypass by inducing a heat shock response. *Anesth Analg.* 2010;111:29-37.
20. Shao X, Tian L, Xu W, Zhang Z, Wang C, Qi C, et al. Diagnostic value of urinary kidney injury molecule 1 for acute kidney injury: a meta-analysis. *PLoS One.* 2014;9:e84131.
21. Coca SG, Nadkarni GN, Garg AX, Koyner J, Thiessen-Philbrook H, McArthur E, et al. First post-operative urinary kidney injury biomarkers and association with the duration of AKI in the TRIBE-AKI cohort. *PLoS One.* 2016;11:e0161098.
22. Neyra JA, Hu MC, Minhajuddin A, Nelson GE, Ahsan SA, Toto RD, et al. Kidney tubular damage and functional biomarkers in acute kidney injury following cardiac surgery. *Kidney Int Rep.* 2019;4:1131-42.
23. Elmistekawy E, Chan V, Bourke ME, Dupui JY, Rubens FD, Mesana TG, et al. Off-pump coronary artery bypass grafting does not preserve renal function better than on-pump coronary artery bypass grafting: results of a case-matched study. *J Thorac Cardiovasc Surg.* 2012;143:85-92.
24. Bove T, Landoni G, Calabro MG, Aletti G, Marino G, Cerchierini E, et al. Renoprotective action of fenoldopam in high-risk patients undergoing cardiac surgery. *Circulation.* 2005;111:3230-5.
25. Oh SY, Kim JC, Choi YS, Lee WK, Lee YK, Kwak YL. Effects of ulinastatin treatment on myocardial and renal injury in patients undergoing aortic valve replacement with cardiopulmonary bypass. *Korean J Anesthesiol.* 2012;62:148-53.

The effect of genetic amniocentesis on Doppler measurements of utero-placental and feto-placental circulations

Efecto de la amniocentesis genética en las mediciones Doppler de las circulaciones útero-placentaria y feto-placentaria

Nizamettin Bozbay^{*ID}, Ahmet Yanar^{ID}, Ayşe C. Duymuş^{ID}, Yasin Altekin^{ID}, Aybike Tazegül-Pekin^{ID}, and Gökçen Örgül^{ID}

Department of Perinatology, Selcuk University Faculty of Medicine, Konya, Turkey

Abstract

Objective: The objective of this study was to determine whether there is a change in feto-maternal circulation after amniocentesis (AS) using Doppler ultrasonography. **Methods:** In this prospectively designed study, fetuses with singleton pregnancies between 16 and 22 weeks of gestational age with an indication for invasive testing and a healthy AS result were included. Twin pregnancies, fetuses with major anomalies as determined by ultrasound evaluation, and fetuses with chromosomal or genetic diseases resulting from AS were excluded from the study. **Results:** A total of 73 patients who underwent AS according to the risky screening result were included in the study. Fetal Doppler measurements were performed at three different time periods. The parameters that were analyzed included the fetal umbilical artery, the ductus venosus, and the maternal right and left uterine arteries. The analysis revealed no statistically significant differences between the measurements. **Conclusions:** It has been observed that AS, the most commonly used invasive diagnostic procedure in clinical practice, does not affect feto-maternal circulation and fetal heart function.

Keywords: Amniocentesis. Doppler ultrasonography. Placental circulation.

Resumen

Objetivo: Determinar si existe un cambio en la circulación feto-materna tras la amniocentesis mediante ultrasonografía Doppler. **Métodos:** Estudio de diseño prospectivo en el que se incluyeron fetos de embarazos únicos de entre 16 y 22 semanas de edad gestacional con indicación de pruebas invasivas y un resultado sano de la amniocentesis. Se excluyeron del estudio los embarazos gemelares, los fetos con anomalías importantes determinadas mediante ecografía y los fetos con enfermedades cromosómicas o genéticas derivadas de la amniocentesis. **Resultados:** Se incluyeron en el estudio 73 pacientes que se sometieron a amniocentesis según el resultado del cribado de riesgo. Se realizaron mediciones de Doppler fetal en tres periodos de tiempo diferentes. Los parámetros analizados fueron la arteria umbilical fetal, el conducto venoso y las arterias uterinas derecha e izquierda de la madre. El análisis no reveló diferencias estadísticamente significativas entre las mediciones. **Conclusiones:** Se ha observado que la amniocentesis, el procedimiento diagnóstico invasivo más utilizado en la práctica clínica, no afecta a la circulación feto-materna ni a la función cardíaca fetal.

Palabras clave: Amniocentesis. Ultrasonografía Doppler. Circulación placentaria.

*Correspondence:

Nizamettin Bozbay

E-mail: dr.nizamettin.bozbay@gmail.com

2444-0507/© 2025 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 12-02-2025

Date of acceptance: 19-03-2025

DOI: 10.24875/CIRUE.M25000897

Cir Cir (Eng). 2025;93(6):600-604

Contents available at PubMed

www.cirugiyacirujanos.com

Introduction

Prenatal diagnosis is a medical procedure used to identify fetal chromosomal abnormalities, single-gene disorders, and congenital infections in the antenatal period. Invasive prenatal diagnostic tests include amniocentesis (AS), chorionic villus sampling, and cordocentesis^{1,2}. AS is an invasive procedure in which amniotic fluid is obtained using a 20-22-gauge spinal needle under ultrasonography (USG) guidance through a transabdominal approach². The first documented case of AS was performed by Fuchs and Riis in 1956 with the objective of determining antenatal sex chromosomes³. AS, when performed by experienced clinicians under outpatient clinic conditions, can be utilized not only for prenatal diagnosis but also for treatment in certain cases^{4,5}. The advent of advanced technology has led to a substantial decrease in the incidence of complications associated with the AS procedure. While transient vaginal bleeding and amniotic fluid leakage may occur in some cases, the risk of miscarriage, the most serious complication, has been reported to be approximately 0.2%^{6,7}.

The potential hemodynamic changes in the fetomaternal circulation following AS are a source of concern. Doppler USG can be used safely as a non-invasive method to detect these changes. The findings from Doppler USG of the fetal umbilical artery, ductus venosus (DV), middle cerebral artery, and maternal uterine artery can provide crucial insights into placental perfusion⁸.

There are very few studies on maternal and fetal perfusion after AS. Previous studies on this subject have reported that fetal circulation was not affected after the procedure^{9,10}. In this study, we tried to investigate whether there is a change in fetomaternal circulation after AS procedure using Doppler USG.

Methods

This prospective study was conducted in the Department of Obstetrics and Gynecology between June 2022 and January 2025. The study was planned in accordance with the principles of the Declaration of Helsinki. Approval was obtained from the Local Ethics Committee of (Ethics Committee number: 2022/309) before the study. Written informed consent was obtained from all patients included in the study.

A total of 73 patients with singleton pregnancies between 16 and 22 weeks of gestation were included

in the study. Fetuses with an indication for invasive testing and a healthy AS result were included in the study. Twin pregnancies, fetuses with major anomalies on USG evaluation, and fetuses with AS results of chromosomal or genetic disorders were excluded from the study.

All procedures were carried out by clinicians specializing in perinatology, utilizing a Voluson E6 (GE Medical Systems, Milwaukee, WI) equipped with a transabdominal 2D convex probe. Fetal Doppler measurements were obtained at three distinct time points: before AS (1st measurement), immediately after AS (2nd measurement), and 1 h after AS (3rd measurement). The parameters that were analyzed included the fetal umbilical artery, the DV, and the maternal right/left uterine artery. The recorded values were then subjected to statistical analysis to determine the presence of any significant differences between them.

Statistical analysis

The analyses were evaluated in Statistical Package for Social Sciences (SPSS) (SPSS Inc., Chicago, IL) 22. Descriptive data were presented as n and % values for categorical data and mean \pm standard deviation (mean \pm standard deviation) and median (minimum-maximum) values for continuous data. The Kolmogorov–Smirnov test was used to assess the normal distribution of continuous variables. For comparison between measurements, the Wilcoxon analysis was employed for paired comparisons, and the Friedman analysis was used for comparisons involving three measurements. The statistical significance level was set at $p < 0.05$.

Results

A total of 73 patients who underwent AS according to high-risk screening results ($n = 57$; 78%), minor soft markers (hyper echogenic cardiac focus, hyperechogenic bowel, nuchal fold thickness, pelviectasia) ($n = 9$; 12%), and advanced maternal age ($n = 7$; 10%) were included in the study. The mean maternal age was 31.9 ± 6.1 (min 21-max 38), and the mean maternal body mass index was 28.0 ± 5.5 kg/m² (min 23-max 33). The mean gestational week at which AS was performed was 18.4 ± 2.0 (min 16-max 22), while the mean estimated fetal weight of the fetuses was 273.8 ± 134.0 g (min 168-max 420).

The first and second Doppler measurements of the umbilical artery, DV, and right and left uterine artery

Table 1. Comparison of first and second Doppler measurements

Doppler measurements	1 st	2 nd	p*
	Mean ± SD	Mean ± SD	
Umbilical artery	3.71 ± 1.13	3.90 ± 1.32	0.360
S/D	1.32 ± 0.47	1.39 ± 0.54	0.510
PI	0.75 ± 0.20	0.78 ± 0.21	0.405
RI			
Right uterine artery			
S/D	3.20 ± 1.26	3.05 ± 1.30	0.220
PI	1.32 ± 0.66	1.23 ± 0.50	0.843
RI	0.65 ± 0.18	0.64 ± 0.17	0.998
Left uterine artery			
S/D	3.00 ± 0.97	3.12 ± 1.29	0.997
PI	1.25 ± 0.58	1.28 ± 0.65	0.999
RI	0.66 ± 0.22	0.65 ± 0.23	0.987
DV			
PI	0.77 ± 0.32	0.78 ± 0.32	0.998

*Wilcoxon analysis was applied.
S/D: systolic/diastolic; PI: pulsatility index; RI: resistance index; DV: ductus venosus; SD: standard deviation.

Table 2. Comparison of the first and third Doppler measurements of the patients

Doppler measurements	1 st	3 rd	p*
	Mean ± SD	Mean ± SD	
Umbilical artery			
S/D	3.71 ± 1.13	3.65 ± 0.76	0.998
PI	1.32 ± 0.47	1.36 ± 0.51	0.936
RI	0.75 ± 0.20	0.77 ± 0.26	0.684
Right uterine artery			
S/D	3.20 ± 1.26	3.07 ± 0.97	0.998
PI	1.32 ± 0.66	1.25 ± 0.51	0.999
RI	0.65 ± 0.18	0.67 ± 0.20	0.987
Left uterine artery			
S/D	3.00 ± 0.97	3.37 ± 1.32	0.099
PI	1.25 ± 0.58	1.31 ± 0.52	0.819
RI	0.66 ± 0.22	0.68 ± 0.18	0.237
DV			
PI	0.77 ± 0.32	0.73 ± 0.18	0.887

*Wilcoxon analysis was applied.
S/D: systolic/diastolic; PI: pulsatility index; RI: resistance index; DV: ductus venosus; SD: standard deviation.

are shown in table 1. There was no statistically significant difference in the Doppler parameters before and immediately after AS ($p > 0.05$ for all).

The data regarding the first and third Doppler measurements of the umbilical artery, DV, and right and left uterine artery are shown in table 2. There was no statistically significant difference in Doppler parameters before and 1 h after the procedure ($p > 0.05$) ($p > 0.05$) (Table 2).

The first, second, and third Doppler measurements of the umbilical artery, DV, right, and left uterine artery are shown in table 3. The umbilical artery systolic/diastolic ratio and pulsatility index (PI) value increased immediately after the procedure and returned to the pre-procedure level at 1 h. However, these findings were not statistically significant.

Discussion

AS is the most commonly performed prenatal invasive diagnostic test, and the incidence of associated complications is very rare when performed by experienced clinicians with superior equipment. However, our understanding of the short-term effects of AS on the fetomaternal circulation and their implications for the fetus remains limited. While numerous studies have examined the indications and complications of AS¹¹⁻¹⁴, there is a paucity of research investigating the potential changes in the fetomaternal circulation

Table 3. Comparison of three different Doppler measurements of the cases

Doppler measurements	1 st	2 nd	3 rd	p*
	Mean ± SD	Mean ± SD	Mean ± SD	
Umbilical artery				
S/D				
PI	3.71 ± 1.13	3.90 ± 1.32	3.65 ± 0.76	0.397
RI	1.32 ± 0.47	1.39 ± 0.54	1.36 ± 0.51	0.286
RI	0.75 ± 0.20	0.78 ± 0.21	0.77 ± 0.26	0.206
Right uterine artery				
S/D				
PI	3.20 ± 1.26	3.05 ± 1.30	3.07 ± 0.97	0.287
RI	1.32 ± 0.66	1.23 ± 0.50	1.25 ± 0.51	0.805
RI	0.65 ± 0.18	0.64 ± 0.17	0.67 ± 0.20	0.494
Left uterine artery				
S/D				
PI	3.00 ± 0.97	3.12 ± 1.29	3.37 ± 1.32	0.132
RI	1.25 ± 0.58	1.28 ± 0.65	1.31 ± 0.52	0.463
RI	0.66 ± 0.22	0.65 ± 0.23	0.68 ± 0.18	0.133
DV				
PI	0.77 ± 0.32	0.78 ± 0.32	0.73 ± 0.18	0.805

*Friedman analysis was applied.
S/D: systolic/diastolic; PI: pulsatility index; RI: resistance index; DV: ductus venosus; SD: standard deviation.

resulting from AS procedures^{9,10}. Consequently, we sought to elucidate the effects of AS on fetomaternal circulation by employing Doppler USG, a non-invasive methodology.

DV is one of the shunt mechanisms that provide oxygen-rich blood to the fetus in intrauterine life, which makes fetal life different from adult life. The DV carries a substantial proportion of the blood flow from the umbilical vein directly to the inferior vena cava, thereby ensuring the delivery of oxygenated blood to the fetal body, particularly the fetal heart and brain¹⁵. Consequently, DV assumes a pivotal role in the assessment of fetal heart function^{16,17}. To date, only two studies have examined the relationship between DV Doppler parameters and AS. One of these studies was conducted by Helbig and Haugen, who evaluated 99 patients who underwent AS for genetic diagnosis. The pre-procedure DV PI measurements of the fetuses were calculated as mean:1.07; min 0.54-max 2.42, and the post-procedure DV PI measurements were calculated as mean:1.03; min 0.51-max 3.27. No significant difference was observed between the two measurements¹⁰. In a subsequent study, DV PI parameters were examined immediately before and after the AS procedure, and the mean and mean values were calculated as 1.050 and 1.050, respectively. 1.050 and 0.930, respectively, and no statistically significant difference was found between these two values. The findings of the present study are consistent with the extant literature; the mean DV PI measurements before AS were 0.77 ± 0.32 , immediately after AS were 0.78 ± 0.32 , and 1 h after AS were 0.73 ± 0.18 . The results were analogous, and no statistically significant difference was identified between them.

In their study, Ulkumen et al. compared the Doppler USG parameters of the umbilical artery, middle cerebral artery, right uterine artery, and left uterine artery before and after the procedure in 56 patients who underwent AS in their clinic. The researchers reported that the results of all parameters before and after the procedure were similar⁹. In a related study, 200 patients who underwent AS for genetic diagnosis were examined, and the patient group who underwent transplacental AS during the procedure and the patient group who did not undergo transplacental needle passage were compared. The umbilical artery Doppler parameters and fetal heart rate were compared between these two groups before and after the procedure. The results showed that both measurements were similar in both groups¹⁸. In a subsequent study by Iskender et al., the Doppler parameters of the middle cerebral artery and umbilical artery were compared before and after the procedure in patients with bleeding into the amniotic fluid during the AS procedure and in patients without such bleeding. The study revealed that the

Doppler parameter measurements were comparable between the two groups¹⁹. The present study involved the performance of uterine artery, umbilical artery, and DV Doppler flow examinations on 73 patients before, immediately following, and 1 h after the AS procedure. The results of the present study corroborate the extant literature, and no statistically significant difference was observed between the groups in Doppler USG parameters evaluating fetomaternal circulation and fetal cardiac function.

The strengths of this study are its prospective design, the expertise of the clinicians, the utilization of advanced technological devices, and the inclusion of only live, healthy fetuses. The limitations of the study are its single-center design and the limited number of patients.

Conclusion

In this research, it was observed that AS, the most commonly used invasive diagnostic procedure in clinical practice, does not affect fetomaternal circulation and fetal cardiac function. Therefore, AS can be safely recommended by experienced clinicians under appropriate conditions.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of humans and animals. The authors declare that the procedures followed complied with the ethical standards of the responsible human experimentation committee and adhered to the World Medical Association and the Declaration of Helsinki. The procedures were approved by the institutional Ethics Committee.

Confidentiality, informed consent, and ethical approval. The authors have followed their institution's confidentiality protocols, obtained informed consent from patients, and received approval from the Ethics Committee. The SAGER guidelines were followed according to the nature of the study.

Declaration on the use of artificial intelligence.

The authors declare that no generative artificial intelligence was used in the writing of this manuscript.

References

1. American College of Obstetricians and Gynecologists. ACOG practice bulletin No. 88, December 2007. Invasive prenatal testing for aneuploidy. *Obstet Gynecol.* 2007;110:1459-67.
2. Jindal A, Sharma M, Karena ZV, Chaudhary C. *Amniocentesis*. StatPearls. Treasure Island, FL: StatPearls Publishing; 2023.
3. Fuchs F, Riis P. Antenatal sex determination. *Nature.* 1956;177:330.
4. Piantelli G, Bedocchi L, Cavicchioni O, Verrotti C, Cavallotti D, Fieni S, et al. Amnioreduction for treatment of severe polyhydramnios. *Acta Biomed.* 2004;75:56-8.
5. Nizard J. Amniocentesis: technique and education. *Curr Opin Obstet Gynecol.* 2010;22:152-4.
6. American College of Obstetricians and Gynecologists. Practice bulletin No. 162: prenatal diagnostic testing for genetic disorders. *Obstet Gynecol.* 2016;127:e108-22.
7. Tara F, Lotfalizadeh M, Moeindarbari S. The effect of diagnostic amniocentesis and its complications on early spontaneous abortion. *Electron Physician.* 2016;8:2787.
8. McCaul JF, Morrison JC. Antenatal fetal assessment: an overview. *Obstet Gynecol Clin N Am.* 1990;17:1-16.
9. Ulkumen BA, Pala HG, Baytur YB, Koyuncu FM. Ductus venosus doppler flow velocity after transplacental and non-transplacental amniocentesis during midtrimester. *Pak J Med Sci.* 2014;30:992-5.
10. Helbig A, Haugen G. Amniocentesis and ductus venosus Doppler blood flow velocity waveforms. *Ultrasound Obstet Gynecol.* 2005;26:435-6.
11. Çetin O, Atmaca FF, Zebitay AG, Derregözü A, Aydoğan B, Keskin S, et al. Second trimester genetic amniocentesis: five-year experience of a maternal-fetal medicine unit. *Gynecol Obstet Reprod Med.* 2013;19:147-52.
12. Locatelli A, Vergani P, Bellini P, Strobelt N, Arreghini A, Ghidini A. Amnioreduction in emergency cerclage with prolapsed membranes: comparison of two methods for reducing the membranes. *Am J Perinatol.* 1999;16:73-7.
13. Romero R, Gonzalez R, Sepulveda W, Brandt F, Ramirez M, Sorokin Y, et al. Infection and labor: VIII. Microbial invasion of the amniotic cavity in patients with suspected cervical incompetence: prevalence and clinical significance. *Am J Obstet Gynecol.* 1992;167:1086-91.
14. Dickinson JE, Tjioe YY, Jude E, Kirk D, Franke M, Nathan E. Amnioreduction in the management of polyhydramnios complicating singleton pregnancies. *Am J Obstet Gynecol.* 2014;211:434-e1.
15. Kiserud T, Rasmussen S, Skulstad S. Blood flow and the degree of shunting through the ductus venosus in the human fetus. *Am J Obstet Gynecol.* 2000;182:147-53.
16. Baschat AA. Ductus venosus Doppler for fetal surveillance in high-risk pregnancies. *Clin Obstet Gynecol.* 2010;53:858-68.
17. Baschat AA, Harman CR. Venous Doppler in the assessment of fetal cardiovascular status. *Curr Opin Obstet Gynecol.* 2006;18:156-63.
18. Haugen G, Helbig A, Husby H. Umbilical artery Doppler flow velocity waveforms after transplacental amniocentesis. *Obstet Gynecol.* 2003;101:697-703.
19. Iskender C, Tarım E, Çök T, Kalaycı H, Parlakgümüş A, Yalçınkaya C. Does bleeding affect fetal Doppler parameters during genetic amniocentesis? *J Turk Ger Gynecol Assoc.* 2014;15:100-3.

Concordance of CO₂ gaps measured using peripheral and central venous blood in patients diagnosed with septic shock

Concordancia de las brechas de CO₂ medidas en sangre venosa periférica y central en pacientes diagnosticados de choque séptico

Erdem Yalcinkaya^{1*}, Umut Sabri Kasapoglu¹, Huseyin Arikan¹, Nazım Onur Sahin¹, Betül Yildiz¹, Esra Tekin², Aybuke Selcuk¹, Emel Eryuksel¹, and Sait Karakurt¹

¹Department of Pulmonary and Critical Care Medicine, Marmara University School of Medicine; ²Department of Critical Care Medicine, Marmara University Hospital. Istanbul, Turkey

Abstract

Objective: This study evaluates the venous-arterial CO₂ difference (Pv-aCO₂) from peripheral venous blood (Pv-aCO_{2p}) as a less invasive alternative to central venous Pv-aCO₂ for assessing tissue perfusion in septic shock. **Methods:** A prospective, single-center study included 54 septic shock patients with central venous catheters placed within 24 h of intensive care unit admission. Blood samples from arterial, central venous, and peripheral venous sources were analyzed. Correlation analyses and Bland-Altman plots were used to assess agreement between Pv-aCO_{2p} and central Pv-aCO₂. **Results:** Patients had a mean age of 70 years, and 51.9% were male. The median SOFA score was 8.5, and the mean APACHE-II score was 22.7. The Pv-aCO_{2p} gap was 8 mmHg, whereas the central Pv-aCO₂ gap was 6 mmHg. A moderate correlation was found between Pv-aCO_{2p} and central Pv-aCO₂ ($r = 0.593$, $p < 0.001$), with stronger correlations observed in patients with arterial lactate ≥ 2 mmol/L ($r = 0.673$) and hemoglobin < 8 g/dL ($r = 0.625$). Bland-Altman analysis revealed a mean difference of 8.278 mmHg between arterial and peripheral pCO₂. **Conclusions:** Peripheral Pv-aCO_{2p} correlates well with central Pv-aCO₂ and can serve as a less invasive alternative for assessing tissue perfusion in septic shock patients. It offers practical utility when central venous access is not available, aiding in early clinical decisions.

Keywords: Pv-aCO₂. Tissue perfusion. Septic shock. Veno-arterial CO₂ difference. Hemodynamic monitoring.

Resumen

Objetivo: Evaluar la diferencia venosa-arterial de CO₂ (Pv-aCO₂) medida en sangre venosa periférica (Pv-aCO_{2p}) como una alternativa menos invasiva a la Pv-aCO₂ venosa central para evaluar la perfusión tisular en el choque séptico. **Métodos:** Estudio prospectivo, unicéntrico, que incluyó 54 pacientes con choque séptico y catéter venoso central colocado dentro de las primeras 24 horas de ingreso a la UCI. Se analizaron muestras de sangre arterial, venosa central y venosa periférica. Se realizaron análisis de correlación y gráficos de Bland-Altman para evaluar la concordancia entre la Pv-aCO_{2p} y la Pv-aCO₂ central. **Resultados:** Los pacientes tenían una edad promedio de 70 años y el 51.9% eran hombres. La mediana del puntaje SOFA fue de 8.5 y el puntaje APACHE-II promedio fue de 22.7. La brecha de Pv-aCO_{2p} fue de 8 mmHg, mientras que la brecha de Pv-aCO₂ central fue de 6 mmHg. Se encontró una correlación moderada entre la Pv-aCO_{2p} y la Pv-aCO₂ central ($r = 0.593$; $p < 0.001$), con correlaciones más fuertes en los pacientes con lactato en sangre arterial ≥ 2 mmol/l ($r = 0.673$) y hemoglobina < 8 g/dl ($r = 0.625$). El análisis de Bland-Altman reveló una diferencia promedio de 8.278 mmHg entre la pCO₂

*Correspondence:

Erdem Yalcinkaya
E-mail: drerdemyalcinkaya@gmail.com

Date of reception: 25-11-2024
Date of acceptance: 07-04-2025
DOI: 10.24875/CIRUE.M25000898

Cir Cir (Eng). 2025;93(6):605-611
Contents available at PubMed
www.cirugiyacirujanos.com

2444-0507/© 2025 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

en sangre arterial y en sangre periférica. **Conclusiones:** La Pv-aCO_{2p} se correlaciona bien con la Pv-aCO₂ central y puede servir como una alternativa menos invasiva para evaluar la perfusión tisular en pacientes con choque séptico. Es de utilidad práctica cuando no se dispone de acceso venoso central, ayudando en las decisiones clínicas tempranas.

Palabras clave: Pv-aCO₂. Perfusión tisular. Choque séptico. Diferencia venosa-arterial de CO₂. Monitoreo hemodinámico.

Introduction

Early identification of tissue hypoperfusion and prompt, effective resuscitation are crucial for the successful management of patients with septic shock¹. The latest guidelines from the Surviving Sepsis Campaign (SSC) for managing patients with septic shock emphasize hemodynamic support using a structured protocol of fluid administration and vasopressor therapy. The objective is to enhance tissue perfusion and fulfill tissue oxygen requirements². The veno-arterial partial pressure of carbon dioxide difference (Pv-aCO₂ or ΔpCO₂) has been suggested as a marker for tissue hypoperfusion³. These findings have immediate clinical relevance, as they can alter the treatment strategy at the patient's bedside. Under physiological conditions, the Pv-aCO₂ typically does not exceed 0.8 kPa (6 mmHg) and reflects the adequacy of venous blood flow and cardiac output (CO)⁴.

Several researchers have emphasized a linear relationship between Pv-aCO₂ determined from mixed or central venous blood^{5,6}. However, the studies conducted so far have been performed in an intensive care setting using central venous catheters. Evaluating the CO₂ gap using peripheral venous blood samples outside/inside the intensive care environment is important for its potential to aid daily practice as well. The objective of this study was to determine whether a linear correlation exists between Pv-aCO₂ measured from peripheral venous blood (Pv-aCO_{2p}) and Pv-aCO₂ measured from central venous blood.

Methods

We conducted a prospective, observational single-center study at Level-3 medical intensive care unit (ICU) of Marmara University Hospital, focusing on patients diagnosed with septic shock according to SSC: International Guidelines for Management of Sepsis and Septic Shock 2021 recommendations (e.g., for patients with sepsis induced hypoperfusion or septic shock we suggest that at least 30 mL/kg of IV crystalloid fluid should be given within the first 3 h of

resuscitation) who were admitted between December 2023 and June 2024⁷.

The study specifically targeted patients who had a central venous catheter inserted within the first 24 h following their diagnosis and with at least one site of peripheral venous access. Patients were excluded if they were under 18 years of age, or if they presented with trauma, burns, thrombosis, or other contraindications to jugular vein catheterization, were undergoing extracorporeal membrane oxygenation, continuous renal replacement therapy, or extracorporeal CO₂ removal therapy, or had a prior diagnosis of congestive heart failure and patients with < 24 h of ICU stay (Fig. 1).

Patient management

All evaluations were performed within 24 h of identifying septic shock in eligible ICU patients. Peripheral blood samples were obtained using a needle cannula in one of the upper extremities, typically in the ICU, with a tourniquet applied. Simultaneously, central venous blood samples were collected under ultrasound guidance following the placement of an internal jugular vein catheter. Patients were thoroughly assessed, with blood samples taken for complete blood count, arterial blood gas analysis, venous blood gas analysis, metabolic panel, N-terminal pro-B-type natriuretic peptide (NT-proBNP), and calculation of Acute Physiology and Chronic Health Evaluation-II (APACHE-II) and the Sequential Organ Failure Assessment (SOFA) scores.

Consent for the hospital's clinical management was requested from each patient, ensuring alignment with their clinical condition. Data were anonymously collected and stored in secure databases accessible solely to investigators specified in the protocol.

Calculation of the venous-to-arterial CO₂ tension difference

Blood samples were collected from central and peripheral veins, as well as arterial blood, to determine CO₂ values in a cohort of 54 patients. The v-a CO₂ gap was calculated by subtracting the arterial pCO₂




Figure 1. Flowchart of the study.

values from the pCO₂ values obtained from the central and peripheral venous blood samples.

Outcomes

The primary aim was the evaluation of the linear correlation and agreement between the Pv-aCO₂ measured with peripheral venous and central venous blood. The secondary objective of the study was to identify patient subgroups or specific laboratory parameters associated with a potentially stronger correlation. To accomplish this, we aimed to stratify patients based on their mean arterial pressures, heart rates, mottling scores, capillary refilling times (CRT), arterial lactate levels, APACHE-II scores, SOFA scores, NT-proBNP values, as well as the type and dosage of vasopressors initiated at the time of septic shock diagnosis.

Statistical analysis

To determine the sample size for this study, literature data were utilized (with a significance level of $\alpha = 0.05$ and power = 90%), leading to the inclusion of a total of 50 patients, with calculations performed using G-Power version 3.1.9.7.8.

The data analysis was conducted using IBM Statistical Package for the Social Sciences Statistics Version 23. Normality of the data distribution was assessed using the Shapiro-Wilk test. Analysis results were presented as mean \pm standard deviation or median (minimum–maximum). The paired samples t-test was employed for normally distributed data in two dependent groups, whereas the Wilcoxon test was used for data that did not follow a normal distribution. Pearson correlation coefficient was used to examine the relationship between normally distributed continuous parameters,

whereas Spearman's ρ correlation coefficient was employed for non-normally distributed continuous parameters.

In the comparative analysis of methods, the robust Bland-Altman analysis was utilized, renowned for its effectiveness in assessing agreement between two measurement techniques or observers. This method offers insights into the agreement and potential biases between two quantitative measurements, contributing valuable information on the degree of agreement and any systematic differences.

Furthermore, the intra-class correlation coefficient (ICC) was utilized to assess the concordance of pCO₂ differences across regions. The ICC is a widely used statistical measure for evaluating the agreement or consistency between multiple measurements or observations. Specifically, in the context of regional PCO₂ differences, ICC sheds light on the overall consistency of measurements across different regions.

By employing these rigorous statistical methods and adhering to established significance levels, the study ensures robustness and reliability in its findings, thereby contributing to the advancement of knowledge in the field and facilitating informed decision-making in clinical practice.

The significance level of $p < 0.050$ was established to determine statistical significance, aligning with conventional practices in hypothesis testing. This threshold ensures that findings with a p-value below 0.050 are considered statistically significant, indicating a high likelihood of rejecting the null hypothesis.

Results

Among the participants, 28 (51.9%) were male. Diabetes mellitus was present in 12 (22.2%) individuals, whereas 22 (40.7%) had hypertension. Coronary arterial disease was found in 9 (16.7%) participants, and chronic renal failure was reported in 12 (22.2%) individuals. The mean age of the participants was 70 ± 15 years. The mean SOFA score was 8.52 ± 3.65 , and the mean APACHE score was 22.7 ± 5.89 . The median CRT was determined to be 3 (1-8) s. The median mottling score was 0 (0-5). Patients' characteristics and baseline laboratory findings are shown in table 1.

Out of 54 determinations, the mean difference in PCO₂ values obtained from arterial and central regions (Pv-aCO₂) was found to be 6.81, with this difference being statistically significant from zero ($p < 0.001$). The lower limit of agreement was -0.848 , whereas the upper limit was 14,478. In addition, a statistically significant

Table 1. Main characteristics, laboratory results, and clinical scores of the study population on ICU admission

Characteristics	Patients
Male sex, n (%)	28 (51.9)
Age (mean value ± SD)	70 ± 15
Comorbid diseases	
DM, n (%)	12 (22.2)
HT, n (%)	22 (40.7)
CAD, n (%)	9 (16.7)
CKD, n (%)	12 (22.2)
COPD, n (%)	4 (7.4)
Hematologic malignancy, n (%)	4 (7.4)
Solid tumor, n (%)	16 (29.6%)
Mean arterial pressure (mmHg), (mean value ± SD)	71.15 ± 10.54
Heart rate (bpm), (mean value ± SD)	96.11 ± 20.88
SOFA score, (median [25-75])	8.5 (1-17)
APACHE-II score, (mean value ± SD)	22.7 ± 5.89
NT-ProBNP (pg/mL), (median [25-75])	4034 (10-35000)
Hgb (g/dL), (mean value ± SD)	9.06 ± 2.05
Htc (%), (mean value ± SD)	27.7 ± 6.36
Noradrenaline dose (mcg/kg/min), (median [25-75])	0.18 (0.02-1)
Capillary refilling time (seconds), (median [25-75])	3 (1-8)
Mottling score, (median [25-75])	0 (0-5)

ICU: intensive care unit; SD: standard deviation; DM: diabetes mellitus; HT: hypertension; CAD: coronary artery disease; CKD: chronic kidney disease; COPD: chronic obstructive pulmonary disease; SOFA: the sequential organ failure assessment; APACHE-II: acute physiology and chronic health evaluation-II; Hgb: hemoglobin; Htc: hematocrit; NT-ProBNP: n-terminal pro-B-type natriuretic peptide.

strong agreement was observed between arterial and central regions (ICC = 0.942; p < 0.001). Similarly, the mean difference in pCO₂ values obtained from arterial and peripheral regions (Pv-aCO_{2p}) was 8.27, with this difference being statistically significant from zero (p < 0.001) (Table 2). The lower limit of agreement was -0.728, whereas the upper limit was 17,283. Furthermore, a statistically significant strong agreement was observed between arterial and peripheral regions (ICC = 0.914; p < 0.001) (Table 3). These findings indicate a consistent and reliable agreement between pCO₂ measurements taken from arterial and central regions, as well as between arterial and peripheral regions. This strong agreement highlights the robustness of comparing pCO₂ values across different regions (Fig. 2).

Among individuals with arterial lactate levels of 2 or higher, there was a strong positive correlation between Pv-aCO₂ and Pv-aCO_{2p} (r = 0.673, p < 0.001). Similarly, individuals with NT-proBNP values < 2000, Hgb values

Table 2. Blood gas analysis of the study population

Parameter	Arterial	Central	Peripheral
pH, (min-max)	7.45 (7.06-7.74)	7.4 (6.98-7.65)	7.38 (7.02-7.58)
pO ₂ , mmHg, (min-max)	92.5 (51-289)	41 (24-137)	36 (15-104)
pCO ₂ , mmHg, (min-max)	36 (14-73)	43.5 (20-84)	46 (23-81)
HCO ₃ ⁻ , mEq/L, (min-max)	25.1 (8.5-38.8)	25.05 (9.9-39.2)	24.45 (9-38.1)
BE, (min-max)	0.6 (-22.5-15.5)	1.15 (-18.9-16.5)	1.65 (-19.3-13.9)
Lactate, mmol/L, (min-max)	1.9 (0.3-8.6)	2.05 (0.1-14.8)	2.2 (0.2-13.7)
sO ₂ %, (min-max)	98 (80-100)	70.72 ± 11.98	60 (24-95)
Pv-aCO ₂ (mmHg), (Median [25-75])		6.81 ± 3.91	
Pv-aCO _{2p} (mmHg), (Median [25-75])		8.27 ± 4.60	

Table 3. Analysis results of Bland-Altman and examination of agreement between Pv-aCO₂ and Pv-aCO_{2p}

Variable	Median difference (95% CI)	SD	ICC (95% CI)	p
Pv-aCO ₂	6.815 (5.75-7.882)	3.910	0.942 (0.902-0.966)	< 0.001
Pv-aCO _{2p}	8.278 (7.02-9.53)	4.595	0.914 (0.856-0.949)	< 0.001

CI: confidence interval; SD: standard deviation; ICC: intra-class correlation coefficient.

< 8, and noradrenaline values < 0.1 exhibited high positive correlations between Pv-aCO₂ and Pv-aCO_{2p} values (r = 0.637, p = 0.006; r = 0.625, p = 0.004; r = 0.620, p = 0.032, respectively) (Table 4).

Discussion

Our study focused on analyzing the Pv-aCO₂ gap obtained from central venous catheters and comparing it to the Pv-aCO_{2p} gap obtained from peripheral venous catheters. The outcomes of our study reveal useful insights into the relationship between these two types of venous access methods and their corresponding Pv-aCO₂ measurements. The research showed a statistically significant relationship between the Pv-aCO₂ gaps measured at central and peripheral venous sites, showing that measurements obtained from peripheral venous access are similarly reflective of those obtained from central venous access.




Figure 2. The Bland-Altman plot comparison shows the agreement between two measurement methods. Each plot presents differences in measurements on the y-axis against their mean values on the x-axis, and includes limits of agreement, typically set at ± 1.96 standard deviations from the mean difference. **A:** the mean difference of PCO_2 values obtained from arterial and central regions was found to be 6.815, with a statistically significant difference from 0 ($p < 0.001$). In this plot, the lower limit of agreement was approximately -0.848 , and the upper limit was about 14.478. Most data points lie within these limits, indicating a strong agreement between the arterial and central regions. The shaded areas (green for the upper limit and red for the lower limit) emphasize the range within which the majority of measurements fall, with only a few outliers observed. The intraclass correlation coefficient (ICC) was 0.942, showing a statistically significant and very strong agreement ($p < 0.001$). **B:** which illustrates the comparison between arterial and peripheral regions, the mean difference of PCO_2 values was calculated as 8.278, also showing a statistically significant difference from 0 ($p < 0.001$). Here, the lower limit of agreement was approximately -0.728 , and the upper limit was around 17.283. While most data points again fall within the shaded limits, the wider spread indicates a slightly greater variability in measurements between the arterial and peripheral regions compared to the central region. Nevertheless, a very strong agreement was still observed between the arterial and peripheral regions, with an ICC of 0.914 ($p < 0.001$), underscoring the statistically significant agreement in this comparison as well.

This finding has significant therapeutic effects, as central venous access, despite its frequent utilization for precise hemodynamic monitoring, is associated with risks such as infection, thrombosis, and pneumothorax, especially in critically ill individuals⁹. On the other hand, peripheral venous access is a minimally invasive treatment that is easier to perform and has a lower risk of problems. If the measurements of the peripheral Pv-aCO₂p gap show a strong correlation with those obtained via central venous access, as shown by our findings, then peripheral venous sampling could be considered a feasible option in cases where central venous access is not possible or considered risky.

The practical applications of using peripheral venous Pv-aCO₂p gap measurements as a substitute for central venous measurements are numerous. In emergency settings or resource-limited environments where central venous catheterization may not be easily accessible or feasible, peripheral venous sampling can be used to effectively monitor tissue perfusion and provide guidance for vital actions. Furthermore, this approach can be especially beneficial for patients who are unable to have central venous access due to anatomical or physiological limitations.

Another crucial factor to take into consideration is the potential impact on patient comfort and overall treatment. Peripheral venous catheterization is characterized by lower pain, better tolerance, and the ability to be performed with fewer assets and staff when compared to central venous catheterization⁹. This has the potential to improve patient outcomes by decreasing problems associated with procedures and enabling faster implementation of monitoring, particularly in urgent situations such as septic shock.

The strong correlations shown in our study have not been thoroughly investigated in previous studies, and the provided values have been isolated from statistical analysis^{10,11}. Although our research provides valuable data in this relatively unexplored field, larger-scale investigations are necessary to validate our findings and identify specific clinical conditions in which peripheral venous sampling might effectively replace a central venous sample.

Various clinical settings are involved in managing septic patients, particularly in the initial stages of the syndrome, including pre-hospital medicine, emergency medicine, and hospital medicine. In these contexts, obtaining a central venous line may pose

Table 4. Comparison of pCO₂ differences in regions by groups

Clinical variable	Pv-aCO ₂ p (mmHg)		Pv-aCO ₂ (mmHg)		r/p	Test statistic	p
	Mean value ± SD	Median (Min-Max)	Mean value ± SD	Median (Min-Max)			
Arterial lactate level, mmol/L							
≥ 2	8.1 ± 4.67	7 (2-18)	6.21 ± 3.34	6 (1-14)	0.433/0.019***	2.316	0.028*
< 2	8.48 ± 4.59	9 (0-18)	7.52 ± 4.45	8 (1-21)	0.673/< 0.001***	1.313	0.202*
NT-proBNP, pg/mL							
< 2000	7.18 ± 4.08	7 (1-13)	7.35 ± 4.7	6 (1-21)	0.637/0.006****	-0.032	0.975**
≥ 2000	9.22 ± 4.67	9.5 (0-18)	6.94 ± 3.51	6.5 (1-14)	0.516/0.003***	3.108	0.004*
Hgb, g/dL							
< 8	7.21 ± 3.74	7 (2-14)	5.0 ± 3.09	5 (1-12)	0.625/0.004***	3.2	0.005*
≥ 8	8.86 ± 4.95	9 (0-18)	7.8 ± 3.99	8 (1-21)	0.545/0.001****	-1.107	0.268**
HR, beats/min							
< 100/min	8.87 ± 4.76	10 (0-18)	6.71 ± 3.36	6 (1-14)	0.600/< 0.001***	3.134	0.004*
≥ 100/min	7.27 ± 4.32	6.5 (1-17)	6.82 ± 4.69	5.5 (1-21)	0.562/0.007****	-0.328	0.743**
Noradrenaline dose (mcg/kg/dak)							
< 0.1	5.75 ± 3.47	6 (1-13)	5.33 ± 3.17	5 (1-11)	0.620/0.032***	0.497	0.629*
≥ 0.1	9.18 ± 4.93	10 (0-18)	7.21 ± 4.36	6 (1-21)	0.540/0.001***	2.524	0.017*

*Paired two-sample t-test.

**Wilcoxon test.

***Pearson correlation coefficient.

****Spearman's rho correlation coefficient.

NT-proBNP: n-terminal pro-B-type natriuretic peptide; Hgb: hemoglobin; HR: heart rate; SD: standard deviation.

challenges. Early assessment of the CO₂ gap could prompt earlier initiation of inotropic medications, which have been demonstrated to be safely administered via peripheral venous access¹².

To the best of our knowledge, only two studies in the literature have explored the association between standard Pv-aCO₂ and Pv-aCO₂p. Gao et al.¹¹ reported a correlation of 90%, and Orso et al.⁸ reported a correlation of 70%.

In the present study, we observed a direct correlation between Pv-aCO₂ and Pv-aCO₂p. The ICC value of 0.914 underscores the strong consistency and reliability of these measurements, affirming their robust agreement and reliability in clinical assessment.

Beyond the primary findings regarding the correlation between Pv-aCO₂ and Pv-aCO₂p gaps, our study revealed several additional correlations that hold important clinical implications and have not been previously addressed in the literature. Among patients with arterial lactate levels of 2 mmol/L or higher, we observed a robust positive correlation between Pv-aCO₂ and Pv-aCO₂p (r = 0.673, p < 0.001). Elevated lactate is often associated with impaired tissue oxygenation and metabolic stress, yet our findings suggest that even in this physiologically challenging condition, peripheral venous measurements remain reliably reflective of central venous values¹³. This could potentially offer a safer, less invasive alternative

for hemodynamic monitoring in patients where elevated lactate levels would traditionally necessitate central venous access.

In addition, in patients with NT-proBNP levels under 2000, hemoglobin values below 8 g/dL, and noradrenaline doses < 0.1 µg/kg/min, we found similarly high positive correlations between Pv-aCO₂ and Pv-aCO₂p (r = 0.637, p = 0.006; r = 0.625, p = 0.004; r = 0.620, p = 0.032, respectively). The ability of peripheral Pv-aCO₂p measurements to align with central values across these diverse clinical markers reinforces its potential role as a practical alternative, particularly in situations where central venous access may be risky or infeasible.

Furthermore, our research revealed a statistically significant moderate positive correlation between capillary refill time and mottling score (p < 0.001), which are both simple bedside measurements of peripheral perfusion^{13,14}. The correlation between these two non-invasive indicators enhances the significance of their combined utilization in evaluating the degree of circulatory failure in critically ill patients.

The results of our study show that peripheral venous measures can be reliably accurate under a variety of clinical situations, including those marked by increased lactate levels, decreased hemoglobin levels, impaired cardiac function, and the administration of low doses of vasopressors. This emphasizes the broader relevance

of peripheral venous sampling as a less invasive, but accurate option for central venous access in critically ill patients. Our investigation offers new insights into the correlations between Pv-aCO₂ and Pv-aCO_{2p} in different conditions. This gives a basis for future research to assess and expand upon these findings, potentially leading to modifications in the approach to venous sampling and hemodynamic monitoring in critical care.

Our study presents several limitations. First, it is a single-center study, so the results might not universally apply. Second, we did not measure cardiac index, and we excluded patients with a prior diagnosis of heart failure from our study to avoid potential confusion with cardiac dysfunction that could develop secondary to sepsis. Finally, our patient group received only single-agent vasopressor therapy with noradrenaline. Therefore, our results should be validated in future studies with a larger sample size.

Conclusion

The significant linear correlation observed between Pv-aCO₂ levels from central and peripheral venous catheters in patients with septic shock underscores the potential benefits of using peripheral venous access. This approach offers a less invasive alternative for Pv-aCO₂ measurement, simplifying patient management while still providing valuable physiological insights. Further controlled clinical trials are necessary to explore whether Pv-aCO_{2p} could be utilized in broader clinical settings beyond ICUs, aiming to assess whether CO adequately meets tissue metabolic demands during early sepsis stages.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgments

The authors acknowledge and thank the American Thoracic Society's Methods in Epidemiologic, Clinical and Operations Research (MECOR) Program and MECOR Turkey Level 2 Faculty; Zuhal Karakurt, Sinem Gungor, Ozge Yilmaz.

Ethical considerations

Protection of humans and animals. The authors declare that no experiments involving humans or animals were conducted for this research.

Confidentiality, informed consent, and ethical approval. The authors have followed their institution's confidentiality protocols, obtained informed consent from patients, and received approval from the Ethics Committee. The SAGER guidelines were followed according to the nature of the study. The protocol for this study was approved by the Marmara University School of Medicine Ethics Committee (Approval No: 09.2023.566, date: 07.04.2023). The study was carried out in strict adherence to the principles outlined in the Declaration of Helsinki.

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing of this manuscript.

References

- Cecconi M, De Backer D, Antonelli M, Beale R, Bakker J, Hofer C, et al. Consensus on circulatory shock and hemodynamic monitoring. Task force of the European Society of Intensive Care Medicine. *Intensive Care Med.* 2014;40:1795-815.
- Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R, et al. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock: 2016. *Intensive Care Med.* 2017;43:304-77.
- Bitar ZI, Maadarani OS, El-Shably AM, Elshabasy RD, Zaalouk TM. The forgotten hemodynamic (PCO₂ gap) in severe sepsis. *Crit Care Res Pract.* 2020;2020:9281623.
- Johnson BA, Weil MH. Redefining ischemia due to circulatory failure as dual defects of oxygen deficits and of carbon dioxide excesses. *Crit Care Med.* 1991;19:1432-8.
- Ho K, Harding R, Chamberlain J. A comparison of central venous-arterial and mixed venous-arterial carbon dioxide tension gradient in circulatory failure. *Anaesth Intensive Care.* 2007;35:695-701.
- Mallat J, Lazkani A, Lemyze M, Pepy F, Meddour M, Gasan G, et al. Repeatability of blood gas parameters, PCO₂ gap, and PCO₂ gap to arterial-to-venous oxygen content difference in critically ill adult patients. *Medicine (Baltimore).* 2015;94:e415.
- Evans L, Rhodes A, Alhazzani W, Antonelli M, Coopersmith CM, French C, et al. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock 2021. *Intensive Care Med.* 2021;47:1181-247.
- Orso D, Molinari C, Bacchetti G, Zanini V, Montanar V, Copetti R, et al. Correlation and agreement between the CO₂ Gap obtained from peripheral venous blood and from mixed venous blood in mechanically ventilated septic patients. *Semin Cardiothorac Vasc Anesth.* 2024;28:18-27.
- Chua CK, Le Guen M, Lim R, Udy A. Using central venous pressure waveform to confirm the placement of an internal jugular central venous catheter in the intensive care unit. *Aust Crit Care.* 2024;37:628-31.
- Waxman MJ, Schechter-Perkins EM. Peripheral intravenous catheters are preferable for emergency department patients with septic shock. *Ann Emerg Med.* 2024;83:607-8.
- Gao W, Zhang Y, Ni H, Zhang J, Zhou D, Yin L, et al. Prognostic value of difference between peripheral venous and arterial partial pressure of carbon dioxide in patients with septic shock: a pilot study. *Nan Fang Yi Ke Da Xue Xue Bao.* 2018;38:1312-7.
- Tran QK, Mester G, Bzhilyanskaya V, Afridi LZ, Andhavarapu S, Alam Z, et al. Complication of vasopressor infusion through peripheral venous catheter: a systematic review and meta-analysis. *Am J Emerg Med.* 2020;38:2434-43.
- Liu S, Yang T, Jiang Q, Zhang L, Shi X, Liu X, et al. Lactate and lactylation in sepsis: a comprehensive review. *J Inflamm Res.* 2024;17:440517.
- Gutierrez-Zarate D, Rosas-Sanchez K, Zaragoza JJ. Clinical evaluation of peripheral tissue perfusion as a predictor of mortality in sepsis and septic shock in the intensive care unit: systematic review and meta-analysis. *Med Intensiva (Engl Ed).* 2023;47:697-707.

Perirenal fat Hounsfield units: not only for renal stones but also for renal cell carcinoma

Unidades Hounsfield de grasa perirrenal: no solo para los cálculos renales, sino también para el carcinoma de células renales

Serkan Akan^{1*}, Caner Ediz², Esin Derin-Çiçek³, Ozgun Ete¹, Yunus E. Kizilkan⁴, Aytac Sahin¹, and Ayhan Verit¹

¹Department of Urology, University of Health Sciences, Fatih Sultan Mehmet Training and Research Hospital, Istanbul; ²Department of Urology, University of Health Sciences, Sultan 2. Abdulhamid Han Training and Research Hospital, Istanbul; ³Department of Radiology, University of Health Sciences, Fatih Sultan Mehmet Training and Research Hospital, Istanbul; ⁴Department of Urology, Bingol State Hospital, Bingol. Turkey

Abstract

Objective: We aimed to evaluate the role of attenuation values of perirenal fat calculated by computed tomography (CT) in differentiating aggressive renal masses. **Methods:** The data of 83 patients with histologically confirmed local stage renal cell carcinoma (RCC), who were treated with nephrectomy in our center, and 78 control group cases were analyzed. All measurements including renal mass volume, abdominal subcutaneous fat area and subcutaneous fat area pelvic subcutaneous fat area, and perirenal fat thickness and Hounsfield unit (perirenal fat tissue thickness [PFT]-Hounsfield unit of perirenal fat [PFHU]) were performed on CT by the same radiologist. **Results:** In the statistical analysis of PFT and PFHU measurements between the groups, fat thickness was found to be lower in the patient group compared with the control group. However, hounsfield unit measurements were statistically significantly higher in the patient group ($p < 0.0001$). A cutoff value of -98.1 for the PFHU can identify the patient group with a sensitivity of 89.2% and a specificity of 84.6% (area under the curve = 0.9; 95% confidence interval: 0.86-0.95 $p < 0.0001$). In the multivariate logistic regression analysis, a PFHU ≥ -98.1 was independently associated with RCC after accounting for clinical and radiologic parameters. **Conclusion:** We believe that measurement of the HU values of perirenal adipose tissue with the CT may be useful in differentiating aggressive renal masses and therefore in determining the appropriate treatment selection.

Keywords: Computed tomography. Hounsfield. Kidney. Perirenal fat. Renal cell carcinoma.

Resumen

Objetivo: Evaluar el papel de los valores de atenuación de la grasa perirrenal calculados por tomografía computarizada (TC) en la diferenciación de masas renales agresivas. **Métodos:** Se analizaron los datos de 83 pacientes con carcinoma de células renales en estadio local confirmado histológicamente, que fueron tratados con nefrectomía en nuestro centro, y de 78 casos del grupo control. Todas las mediciones, incluido el volumen de masa renal, el área de grasa subcutánea abdominal y pélvica (SFA_A - SFA_P), el espesor de la grasa perirrenal y las unidades Hounsfield (PFT-PFHU), fueron realizadas en TC por el mismo radiólogo. **Resultados:** En el análisis estadístico de las mediciones de PFT y PFHU entre los grupos se encontró que el espesor de la grasa era menor en los pacientes en comparación con los controles. Sin embargo, las mediciones de las unidades Hounsfield fueron mayores, con significación estadística, en el grupo de pacientes ($p < 0.0001$). Un valor de corte de -98.1 para la PFHU puede identificar el grupo de pacientes con una sensibilidad del 89.2% y una especificidad del 84.6% (AUC = 0.9; IC 95%: 0.86-0.95; $p < 0.0001$). En el análisis de regresión logística multivariado, una PFHU ≥ -98.1 se asoció de manera independiente con el carcinoma de células renales después de tener en cuenta los parámetros clínicos y radiológicos.

*Correspondence:

Serkan Akan

E-mail: drserkanakan@hotmail.com

2444-0507/© 2024 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 23-05-2024

Date of acceptance: 28-10-2024

DOI: 10.24875/CIRUE.M24000899

Cir Cir (Eng). 2025;93(6):612-619

Contents available at PubMed

www.cirurgiaycirujanos.com

Conclusiones: Creemos que la medición de los valores de unidades Hounsfield del tejido adiposo perirrenal con TC puede ser útil para diferenciar masas renales agresivas y, por tanto, para determinar la selección del tratamiento adecuado.

Palabras clave: Tomografía computarizada. Unidades Hounsfield. Riñón. Grasa perirrenal. Carcinoma de células renales.

Introduction

Renal cell carcinoma (RCC) is the most common solid lesion in the kidney and accounts for approximately 90% of all renal malignancies and is associated with the highest mortality^{1,2}. The three main subtypes with specific histopathologic and genetic features are clear cell RCC (ccRCC), papillary RCC (pRCC types I and II), and chromophobe cell RCC (chRCC)³. The diagnosis of early stage RCCs is increased with the widespread use of imaging methods.

There is numerous epidemiologic evidence showing that long-term obesity is one of the major causes of some types of cancers, such as breast, colon, esophageal, pancreatic, and renal cancers⁴. Although the exact etiology of RCC remains unclear, the association of obesity with RCC development was reported in some previous studies^{5,6}. In addition, perirenal fat may promote migration of the ccRCC cells⁷. A recent study showed that renal neoplasms were more common in patients with a high perirenal fat tissue thickness (PFT) than in those with a low PFT⁸. It is also reported that the density of the perirenal fat is a negative factor for the progression-free survival of localized renal cancer⁹.

The hounsfield unit (HU) measurement on computed tomography (CT) is routinely performed in clinical practice to define the hardness and composition of kidney stones and to predict the outcome of the stone treatment¹⁰⁻¹². It is known that CT has high sensitivity and specificity in the diagnosis and extension of renal tumors¹³. In patients with suspected RCC undergoing routine abdominal CT imaging, the HU can be easily measured and reported without the need for extra procedures. Detection of the radiologic and histopathologic association of the high PFT and high perirenal fat density (hounsfield unit of perirenal fat [PFHU]) with renal masses may help in planning appropriate treatment at the right time.

Methods

Patients and data collection

Patients, who were treated with partial or radical nephrectomy between January 2020 and December

2023 in two centers of University of Health Sciences, were included in the study. Those, whose pre-operative CT images were unavailable, who underwent the surgery in another hospital, and whose tumor location may have affected the subcutaneous fat thickness (SFT) measurement (posterior in the renal hilus) were excluded. Individuals, who were under 45 or over 80 years of age, had experienced a previous intra-abdominal surgery, had a history of kidney stones, a malignancy other than RCC, and infection around the kidney, and had incomplete data were excluded. Numbers of patients with pRCC and chRCC subtypes were insufficient for statistical analysis, and therefore, they were also excluded. A total of 100 cases were picked randomly and included as the control group among the participants who underwent a CT imaging for screening in the above-mentioned time period at our center and had similar descriptive characteristics such as age, gender, body mass index (BMI), and comorbidities. A total of 22 individuals, who had a history of intra-abdominal surgery or kidney stones, and had kidney stones, malignancy or infection around the kidney in the admission were excluded. The study was completed with a total of 83 patients with clinically and histologically confirmed local stage ccRCC and 78 patients are the control group (Fig. 1).

Age at the diagnosis, BMI, tumor location (right and left kidney), and the presence of systemic diseases (hypertension and diabetes mellitus) were retrospectively obtained from medical records. T stage based on the American Joint Committee on Cancer (T1vs. T2), Fuhrman grade, RCC subtype, histological necrosis, and presence of sarcomatoid component were reported by our uropathologist¹⁴. All measurements including the renal mass volume, abdominal subcutaneous fat area (SFA_A), pelvic subcutaneous fat area (SFA_P), PFT, and PFHU were performed on CT by the same radiologist Dr. Esin Derin-Çiçek (third author).

Regional Ethic Committee (Hamidiye-BAEK 20/268) reviewed and approved the study protocol and the study's protocol was in accordance with by the 1964 Declaration of Helsinki and its later changes. All participants included in the study obtained informed consent form.




Figure 1. Flowchart of patients who met inclusion/exclusion criteria for the study.

The measurement of the SFA, PFT, and PFHU

A non-contrast CT scan of abdomen (Optima 660 SE, GE Healthcare/Aquilion Lightning Ultra Helical CT, Canon Medical) was performed for all individuals in the preoperative period. The SFA_A and SFA_P (Fig. 2) calculations were consistent with those reported in the previous studies^{15,16}. First, the border between the visceral and subcutaneous compartment was defined manually along with the contour of the abdominal muscular wall at the umbilical level. Then, to identify the fat containing pixels, an image display window width between -195 and -45 HUs was utilized. A software analysis program (RAD-X[®] PACS software, Simplex, Ankara, Turkey) evaluated the values of the SFA_A and SFA_P automatically.

The PFT and PFHU were also calculated with the same method as in the previous studies^{8,16}. In the CT images, the linear ruler tool of the software was used to measure the distance from the posterior surface of the kidney to the external edge of the iliopsoas at the renal hilum level (Fig. 3). The same radiologist Dr. Esin Derin-Çiçek (third author), who was unaware of




Figure 2. A: evaluation of the subcutaneous fat area of the abdomen. B: evaluation of the subcutaneous fat area of the pelvis.

the clinical and pathological data, assessed the all preoperative scans.

Statistical analysis

IBM Statistical Package for the Social Sciences Statistics 22 program was used for statistical analysis of the findings. The conformity of the parameters to normal distribution was evaluated by Kolmogorov-Smirnov test. In addition to descriptive statistical methods (minimum, maximum, mean, standard deviation, median, and frequency), t-test was used for comparisons of quantitative data between two groups



Figure 3. The evaluation of the perirenal fat thickness and perirenal fat hounsfield unit.

for normally distributed parameters and Mann-Whitney U test was used for comparisons of non-normally distributed parameters between two group receiver operating characteristics (ROC). Chi-square and Fisher's Exact test were used to compare qualitative data. Spearman's rho correlation analysis, ROC for discrimination, and logistic regression analysis for effect were used to analyze the relationships. Analysis of covariance was performed to control/refine the SFA values. Significance was evaluated at $p < 0.05$ level.

Results

The study was conducted with a total of 161 patients, 59 (36.7%) women and 102 (63.3%) men, aged between 45 and 79 years. The mean age was 58.5 ± 8.56 years. The cases were evaluated in two groups as "Patient" ($n = 83$) and "Control" ($n = 78$). There was no statistically significant difference between the two groups in terms of descriptive characteristics ($p > 0.05$; Table 1).

The pre-operative SFA_A and SFA_P measurements were not statistically significantly different between

the groups ($p > 0.05$). In the pre-operative PFT measurement analysis in both groups, fat thickness was found to be lower in the patient group compared with the control group; however, this difference was not statistically significant ($p > 0.05$). In separate comparisons of both kidneys, fat thickness was found to be lower in the patient group than in the control group according to the PFT (PFT_R) measured in the right kidney and the PFT (PFT_L) measured in the left kidney; however, these differences were not statistically significant ($p > 0.05$). After eliminating the effect of the SFA_A and SFA_P values in our study, the PFT measurements were re-evaluated between the two groups; however, no statistically significant difference was found ($p > 0.05$; Table 1).

The analysis of the pre-operative PFHU measurements revealed higher perirenal fat densities in the patient group compared with the control group; this difference was statistically significant ($p < 0.0001$). In separate comparisons for both kidneys, perirenal fat densities were found to be higher in the patient group than in the control group according to the analysis of the PFHU (PFHU_R) measured in the right kidney and the PFHU (PFHU_L) measured in the left kidney; these

Table 1. Demographic data and the radiological measurements of the participants among the groups

Data and measurements	Patient (mean) (%)	Control (mean) (%)	Total (mean) (%)	p
Age	59.10 ± 9.29	57.87 ± 7.71	58.50 ± 8.56	0.451 ¹
Sex				
Male	54 (65)	102 (63)	48 (61)	0.643 ¹
Female	29 (35)	59 (37)	30 (39)	
BMI				
< 25	37 (45)	72 (45)	35 (45)	0.97 ¹
≥ 25	46 (55)	89 (55)	43 (55)	
HT				
+	46 (55)	80 (50)	34 (44)	0.133 ¹
-	37 (45)	81 (50)	44 (56)	
DM				
+	24 (29)	40 (25)	16 (20)	0.218 ¹
-	59 (71)	121 (75)	62 (80)	
SFA _A	25.65 ± 8.79	25.10 ± 7.49	25.38 ± 8.17	0.671 ²
SFA _P	61.96 ± 16.28	59.61 ± 13.64	60.82 ± 15.06	0.325 ²
PFT	16.31 ± 9.12	17.78 ± 7.07	17.02 ± 8.2	0.256 ²
PFT _R	15.53 ± 9.41	16.58 ± 7.01	16.04 ± 8.33	0.423 ²
PFT _L	17.09 ± 10.05	18.99 ± 8.07	18.01 ± 9.17	0.187 ²
PFHU	-92.63 ± 6.43	-103.63 ± 7.03	-97.96 ± 8.68	0.0001 ^{2*}
PFHU _R	-92.88 ± 8.55	-103.03 ± 8.06	-97.8 ± 9.73	0.0001 ^{2*}
PFHU _L	-92.38 ± 8.47	-104.24 ± 7.99	-98.13 ± 10.14	0.0001 ^{2*}

¹Mann-Whitney U-test.

²t-test.

*p < 0.05.

BMI: body mass index; HT: hypertension; DM: diabetes mellitus; SFA: subcutaneous fat area; SFA_A: subcutaneous fat area of abdomen; SFA_P: subcutaneous fat area of pelvis; PFT: perirenal fat thickness; PFT_R: perirenal fat thickness of right kidney; PFT_L: perirenal fat thickness of left kidney; PFHU: kounsfeld unit of perirenal Fat; PFHU_R: kounsfeld unit of right kidney perirenal fat; PFHU_L: kounsfeld unit of left kidney perirenal fat.

Table 2. Histopathologic data of the patient group

Characteristics	Tumor Location		p
	L (%)	R (%)	
T Stage			
1	36 (64)	20 (36)	0.059 ¹
2	5 (42)	7 (58)	
3	5 (33)	10 (67)	
Fuhrman Grade			
1	7 (64)	4 (36)	0.587 ¹
2	26 (51)	25 (49)	
3	13 (62)	8 (38)	
Tumor necrosis			
+	15 (71)	6 (29)	0.088 ¹
-	31 (50)	31 (50)	
Sarcomatoid component			
+	1 (50)	1 (50)	0.99 ²
-	45 (56)	36 (44)	

¹Ki-Kare test.

²Fisher's exact test.

differences were statistically significant (p < 0.0001; Table 1).

The histopathologic data of the patient group are shown in table 2. According to T staging, the rates of stage 1, 2, and 3 were 67.5%, 14.5%, and 18%, respectively. Fuhrman grades 1, 2, and 3 were 11%, 51%, and 21%, respectively. Tumor necrosis was positive in 21% of patients and sarcomatoid component was positive in 2.5%. T stages, Fuhrman grades, tumor necrosis, and presence or absence of the sarcomatoid component were not significantly different in the analysis of the tumor formation sides (left/right) (p > 0.05) (Table 2).

In the analysis of radiologic measurements (Table 3), there was no statistically significant difference between the right and left kidney for the SFA_A, SFA_P, and PFHU values in the patient group (p > 0.05). There was a statistically significant difference between the PFT_L and PFT_R in both patient and control groups (p < 0.0001). In the patient group, the measurement of the tumor side and in the control group and the mean of the measurements of the right and left sides were taken for further comparison. While the difference in the PFT measurements between the two groups was not statistically significant, the difference in the PFHU measurements was statistically significant (p = 0.0001).

The PFHU measurement was found to be highly differentiative for the patient group, the cutoff point was set as -98.1 and the participants were divided into two groups. Those with a PFHU value < -98.1 had 6.7 times more risk than those with a PFHU value > -98.1 (Exp(B) = 6.725; 95% confidence interval [CI] 4.2-10.6 p < 0.0001). A cutoff value of -98.1 for the PFHU can identify the patient group with a sensitivity of 89.2% and a specificity of 84.6% (area under the curve = 0.9; 95% CI: 0.86-0.95 p < 0.0001) (Fig. 4). In multivariate logistic regression analysis, a PFHU ≥ -98.1 was independently associated with RCC after clinical and radiological parameters were added to the analysis.

Discussion


The relationship between ccRCC, the most common histologic subtype of RCC, and obesity has been emphasized in previous studies^{16,17}. Obesity is highly associated with an increased risk of RCC development; therefore, it is considered to be one of the best known etiologic factors of RCC^{5,6}. A high BMI appears to be associated with poor survival outcomes in both

Table 3. Data on radiological measurement

Outcomes	L	R	p
SFA _A	26.05 ± 9.14	25.15 ± 8.45	0.647 ¹
SFA _P	61.71 ± 16.64	62.27 ± 16.05	0.877 ¹
PFT			
Patient	17.09 ± 10.05	15.53 ± 9.41	0.042*
Control	18.99 ± 8.07	16.58 ± 7.01	0.0012*
PFHU			
Patient	-92.38 ± 8.47	-92.88 ± 8.55	0.685 ²
Control	-104.24 ± 7.99	-103.03 ± 8.06	0.172 ²

¹t-test.²paired sample t-test.

*p < 0.05.

SFA_A: subcutaneous fat area of abdomen; SFA_P: subcutaneous fat area of pelvis; PFT: perirenal fat thickness; PFHU: hounsfield unit of perirenal fat.**Figure 4.** Cutoff point determination for perirenal fat thickness hounsfield unit in radiologic imaging of renal cell carcinoma.

non-metastatic and metastatic RCC^{18,19}. There are also studies on the role of body composition indices measured on cross-sectional imaging, such as sarcopenia and fat accumulation, in the prognosis of RCC^{20,21}.

Visceral and subcutaneous fat thickness measurements are quantitative methods calculated on CT. These measurements, which have been emphasized recently, may provide more objective results than BMI. The visceral fat, which increases with age, has been shown to increase the risk of metabolic diseases as well as prostate cancer (PCa) in men²². Another study showed that PCa patients with higher periprostatic fat levels had more aggressive PCa²³. In studies on RCC, Zhu et al. reported a positive correlation between the

percentage of visceral adipose tissue and the Fuhrman tumor grade²⁴. Huang et al. revealed that high visceral fat accumulation played a positive role on the progression of ccRCC¹⁶. On the contrary, there are also studies reporting that high visceral fat deposition is associated with relatively long cancer-specific survival in RCC patients¹⁵.

Perirenal fat accumulation is an ectopic fat accumulation and has been reported to be associated with high diastolic blood pressure, diabetes mellitus, metabolic syndrome, and chronic renal failure^{25,26}. In 2012, Okhunov et al. revealed that perirenal fat accumulation was an important predictor of ccRCC histopathology⁸. They showed that WNT signaling pathway-related factors secreted from perirenal adipose tissue affected the oncogenic pathways critical for ccRCC (migration of ACHN and CaKi-2 cells) and promoted the progression of local ccRCC to locally advanced (T3 stage) disease⁷. Huang et al. reported that the high PFT was associated with lower progression-free survival for ccRCC in both sexes¹⁶. In our study, we compared the images of histologically confirmed ccRCC patients with those of control group participants who were equivalent to them in terms of descriptive characteristics such as age, gender, BMI, and comorbidity (we measured the PFT using the technique described by Okhunov et al. and Huang et al., due to the lack of any standardized measurement method). The PFT was lower in the patient group compared with the control group; however, this difference was not statistically significant. The PFT was again compared by separating the right and left kidneys according to the side of the cancerous kidney, and again the PFT was found to be lower in the patient group. The PFT measurements were re-evaluated by eliminating the effect of the SFA_A and SFA_P values; however, again, no statistically significant difference was found. The PFT measurements for both genders were compared again separately; no significant difference was found between the PFTs for male and female genders in all cases and in the control group. However, the mean PFT measurement of women in the patient group was statistically significantly lower (13.2 ± 7.9, 17.9 ± 9.3; p = 0.023).

So far, most studies on the perirenal fat have focused on the thickness and volume of the adipose tissue^{16,27}. However, several experimental studies suggest that the fat quality (HU) measurements may also be important^{28,29}. They have shown that a lower HU is associated with an adipose tissue containing higher levels of lipid content and the range of HU typically

attributed to fat is in the range of -195 to -45^{27} . The fat depots with higher lipid content and lipolytic activity can cause endothelial dysfunction and high insulin resistance³⁰. These studies also suggest that adipocyte hypertrophy is associated with vascularity; decreased perfusion may lead to hypoxia and inflammation in adipose tissue³¹.

In characterization of the renal masses, the CT still remains as the first-line imaging modality³². Based on the above mentioned studies, we think that the perirenal fat tissue density (PFHU), which can be easily measured by non-contrast CT, may be associated with RCC independently of the abdominal adipose tissue. In our study, the mean PFHU value of the RCC group was significantly higher than that of the control group (-92.6 ± 6.4 vs. -103.6 ± 7 ; $p = 0.0001$). Right and left kidney measurements were also compared and similar results were obtained. In our study, gender-specific PFHU was also evaluated; the mean PFHU value for males was found to be higher (denser) in both the patient group and the control group, but this difference was not significant. According to the results of our study, those with a PFHU value < -98.1 were 6.7 times more likely to have RCC than those with a PFHU value > -98.1 . In multivariate logistic regression analysis, a PFHU ≥ -98.1 was independently associated with RCC after accounting for clinical and radiologic parameters.

The main limitations of our study are its retrospective nature, relatively small sample size and the lack of survival analysis. Second, only histologically confirmed localized ccRCC patients were included in this study, as the numbers of patients with other histological subtypes were not sufficient for statistical study. Third, our study may not include different ethnic groups. However, to our knowledge, our study is the first to examine the association between the PFHU and ccRCC, in English literature.

Conclusion

Although potential mechanisms for the association of the perirenal adipose tissue thickness with RCC are known, its exact role remains unclear. We suggest that the measurement of the perirenal adipose tissue HU by CT may be useful in differentiating the aggressive renal masses and therefore in determining the appropriate treatment selection. The effects of the PFT and PFHU need to be clarified by further research.

Acknowledgments

The authors would like to thank all the Cirugía family who worked hard to write the article.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that they have followed the protocols of their work center on the publication of patient data.

Right to privacy and informed consent. The authors declare that no patient data appear in this article.

Use of artificial intelligence for generating text. The authors declare that they have not used any type of generative artificial intelligence for the writing of this manuscript, nor for the creation of images, graphics, tables, or their corresponding captions.

References

1. Ferlay J, Colombet M, Soerjomataram I, Dyba T, Randi G, Bettio M, et al. Cancer incidence and mortality patterns in Europe: estimates for 40 countries and 25 major cancers in 2018. *Eur J Cancer*. 2018;103:356-87.
2. Sánchez-Morales GE, Osorio-Serrano JL, Guerrero-Gómez A, Chan C, Domínguez-Rosado I. Resección quirúrgica y sobrevida de metástasis de cáncer renal de células claras a páncreas [Surgical resection and survival of clear cell renal cancer metastases to the pancreas]. *Cir Cir*. 2023;92:451-5.
3. Moch H, Cubilla AL, Humphrey PA, Reuter VE, Ulbright TM. The 2016 WHO classification of tumours of the urinary system and male genital organs-part a: renal, penile, and testicular tumours. *Eur Urol*. 2016;70:93-105.
4. Calle EE, Kaaks R. Overweight, obesity and cancer: epidemiological evidence and proposed mechanisms. *Nat Rev Cancer*. 2004;4:579-91.
5. Leiba A, Kark JD, Afek A, Derazne E, Barchana M, Tzur D, et al. Adolescent obesity and paternal country of origin predict renal cell carcinoma: a cohort study of 1.1 million 16 to 19-year-old males. *J Urol*. 2013;189:25-9.
6. Gati A, Kouidhi S, Marrakchi R, El Gaaied A, Kourda N, Derouiche A, et al. Obesity and renal cancer: role of adipokines in the tumor-immune system conflict. *Oncoimmunology*. 2014;3:e27810.
7. Zi X, Lusch A, Blair CA, Okhunov Z, Yokoyama NN, Liu S, et al. Effect of perineoplasm perinephric adipose tissues on migration of clear cell renal cell carcinoma cells: a potential role of WNT signaling. *Oncotarget*. 2016;7:53277-88.
8. Okhunov Z, Mues AC, Kline M, Haramis G, Xu B, Mirabile G, et al. Evaluation of perirenal fat as a predictor of cT 1a renal cortical neoplasm histopathology and surgical outcomes. *J Endourol*. 2012;26:911-6.
9. Thiel DD, Davidiuk AJ, Meschia C, Serie D, Custer K, Petrou SP, et al. Mayo adhesive probability score is associated with localized renal cell carcinoma progression-free survival. *Urology*. 2016;89:54-62.

10. Cohen A, Anderson B, Gerber G. Hounsfield Units for nephrolithiasis: predictive power for the clinical urologist. *Can J Urol.* 2017;24:8832-7.
11. Gallioi A, De Lorenzis E, Boeri L, Delor M, Zanetti SP, Longo F, et al. Clinical utility of computed tomography Hounsfield characterization for percutaneous nephrolithotomy: a cross-sectional study. *BMC Urol.* 2017;17:104.
12. Sugino Y, Kato T, Furuya S, Sasaki T, Arima K, Sugimura Y. The usefulness of the maximum Hounsfield units (HU) in predicting the shockwave lithotripsy outcome for ureteral stones and the proposal of novel indicators using the maximum HU. *Urolithiasis.* 2020;48:85-91.
13. Ndiaye M, Sine B, Sarr A, Akpo G, Gaye AM, Kadiaké H, et al. Evaluation of the correlation between computed tomography and anatomopathological findings in adult renal tumors. *Acad J Health Sci Med Balear.* 2023;38:125-9.
14. Amin MB, Edge SB. In: American Joint Committee on Cancer, editor. *AJCC Cancer Staging Manual.* Chicago: Springer; 2017.
15. Lee HW, Jeong BC, Seo SI, Jeon SS, Lee HM, Choi HY, et al. Prognostic significance of visceral obesity in patients with advanced renal cell carcinoma undergoing nephrectomy. *Int J Urol.* 2015;22:455-61.
16. Huang H, Chen S, Li W, Wu X, Xing J. High perirenal fat thickness predicts a poor progression-free survival in patients with localized clear cell renal cell carcinoma. *Urol Oncol.* 2018;36:157.e1-6.
17. Lowrance WT, Thompson RH, Yee DS, Kaag M, Donat SM, Russo P. Obesity is associated with a higher risk of clear-cell renal cell carcinoma than with other histologies. *BJU Int.* 2010;105:16-20.
18. Donin NM, Pantuck A, Klöpfer P, Bevan P, Fall B, Said J, et al. Body mass index and survival in a prospective randomized trial of localized high-risk renal cell carcinoma. *Cancer Epidemiol Biomarkers Prev.* 2016;25:1326-32.
19. Petrelli F, Cortellini A, Indini A, Tomasello G, Ghidini M, Nigro O, et al. Association of obesity with survival outcomes in patients with cancer: a systematic review and meta-analysis. *JAMA Netw Open.* 2021;4:e213520.
20. Dai J, Zhang X, Liu Z, Song T, Zhu X, Zhang H, et al. The prognostic value of body fat components in metastasis renal cell carcinoma patients treated with TKIs. *Cancer Manag Res.* 2020;12:891-903.
21. Tsutsumi T, Komura K, Hashimoto T, Muraoka R, Satake N, Matsunaga T, et al. Distinct effect of body mass index by sex as a prognostic factor in localized renal cell carcinoma treated with nephrectomy – data from a multi-institutional study in Japan –. *BMC Cancer.* 2021;21:201.
22. Von Hafe P, Pina F, Pérez A, Tavares M, Barros H. Visceral fat accumulation as a risk factor for prostate cancer. *Obes Res.* 2004;12:1930-5.
23. Van Roermund JG, Hinnen KA, Tolman CJ, Bol GH, Witjes JA, Bosch J, et al. Periprostatic fat correlates with tumour aggressiveness in prostate cancer patients. *BJU Int.* 2011;107:1775-9.
24. Zhu Y, Wang HK, Zhang HL, Yao XD, Zhang SL, Dai B, et al. Visceral obesity and risk of high grade disease in clinical t1a renal cell carcinoma. *J Urol.* 2013;189:447-53.
25. De Pergola G, Campobasso N, Nardecchia A, Triggiani V, Caccavo D, Gesualdo L, et al. Para- and perirenal ultrasonographic fat thickness is associated with 24-hours mean diastolic blood pressure levels in overweight and obese subjects. *BMC Cardiovasc Disord.* 2015;15:108.
26. Roever L, Tse G, Biondi-Zoccai G. Perirenal fat and chronic kidney disease in patients with diabetes. *Diabetes.* 2021;70:2190-1.
27. Maurovich-Horvat P, Massaro J, Fox CS, Moselewski F, O'Donnell CJ, Hoffmann U. Comparison of anthropometric, area- and volume-based assessment of abdominal subcutaneous and visceral adipose tissue volumes using multi-detector computed tomography. *Int J Obes (Lond).* 2007;31:500-6.
28. Baba S, Jacene HA, Engles JM, Honda H, Wahl RL. CT Hounsfield units of brown adipose tissue increase with activation: preclinical and clinical studies. *J Nucl Med.* 2010;51:246-50.
29. Hu HH, Chung SA, Nayak KS, Jackson HA, Gilsanz V. Differential computed tomographic attenuation of metabolically active and inactive adipose tissues: preliminary findings. *J Comput Assist Tomogr.* 2011;35:65-71.
30. Rosenquist KJ, Pedley A, Massaro JM, Therkelsen KE, Murabito JM, Hoffmann U, et al. Visceral and subcutaneous fat quality and cardio-metabolic risk. *JACC Cardiovasc Imaging.* 2013;6:762-71.
31. Gealekman O, Guseva N, Hartigan C, Apotheker S, Gorgoglione M, Gurav K, et al. Depot-specific differences and insufficient subcutaneous adipose tissue angiogenesis in human obesity. *Circulation.* 2011;123:186-94.
32. Krishna S, Murray CA, McInnes MD, Chatelain R, Siddaiah M, Al-Dandan O, et al. CT imaging of solid renal masses: pitfalls and solutions. *Clin Radiol.* 2017;72:708-21.

Neoadjuvant chemo(radio)therapy and anastomotic leakage risk in patients with esophageal squamous cell carcinoma

Quimio(radio)terapia neoadyuvante y riesgo de fuga anastomótica en pacientes con carcinoma de células escamosas esofágico

Yuzhen Xie^{1,#}, Hongni Qin^{1,#*}, Fan Wu¹, Yong Zhang¹, and Yang Yang²

¹Department of Life Sciences, College of Biotechnology, Suzhou Industrial Park Institute of Services Outsourcing, Jiangsu Province, Suzhou;

²Department of Thoracic Oncology, Zhejiang Cancer Hospital, Zhejiang Province, Hangzhou, China

#These authors contributed equally and share first authorship.

Abstract

Objective: To investigate the different impacts of neoadjuvant chemotherapy (NCT) and neoadjuvant chemoradiotherapy (NCRT) on anastomotic leakage after esophageal cancer resection. **Methods:** The data of three clinical studies (222 patients) launched were retrospectively collected, including 113 patients in the NCRT group and 109 patients in the NCT group. The NCT group received platinum-based doublet chemotherapy once every 3-4 weeks, whereas the NCRT group received platinum-based doublet chemotherapy weekly (taxane-containing regimen) or once every 3-4 weeks (5-Fu-containing regimen) concurrently with thoracic radiotherapy. After neoadjuvant therapy, all patients underwent Ivor-Lewis or McKeown esophagectomy. The study compared the pathologic complete remission (pCR) rate, 30-day post-operative mortality rate, incidence of anastomotic leakage, post-operative complications, and survival rates between the two groups. **Results:** The pCR rate, 30-day post-operative mortality rate, and the incidence of anastomotic leakage were 39.8% versus 5.5% ($p < 0.001$), 2.7% versus 0% ($p = 0.043$), and 28.3% versus 5.5% ($p < 0.001$) in the NCRT group and NCT group, respectively. Multivariate analysis showed that only NCRT significantly increased the incidence of anastomotic leakage (odds ratio 7.28, 95% confidence interval 2.44-21.78, $p < 0.001$). **Conclusions:** The results show that NCRT can improve the pathological response rate but will increase the incidence of anastomotic leakage. Radiation dose-volume parameters were not associated with anastomotic leakage risk.

Keywords: Malignant esophageal tumor. Neoadjuvant therapy. Anastomotic leak. Risk factors. Chemoradiation.

Resumen

Objetivo: Investigar el diferente impacto de la quimioterapia neoadyuvante (QTN) y la quimiorradioterapia neoadyuvante (QRTN) en la fuga anastomótica después de la resección de cáncer de esófago. **Métodos:** Se recopilaron retrospectivamente los datos de tres estudios clínicos (222 pacientes), incluyendo 113 pacientes en el grupo de QRTN y 109 pacientes en el grupo de QTN. El grupo de QTN recibió quimioterapia de doblete a base de platino una vez cada 3-4 semanas, mientras que el grupo de QRTN recibió quimioterapia de doblete a base de platino semanal (régimen que contiene taxano) o una vez cada 3-4 semanas (régimen que contiene 5-fluorouracilo) de forma concurrente con radioterapia torácica. Después del tratamiento neoadyuvante, todos los pacientes se sometieron a esofagectomía de Ivor-Lewis o McKeown. El estudio comparó las tasas de remisión completa patológica, mortalidad posoperatoria a 30 días, incidencia de fuga de anastomosis, complicaciones posoperatorias y tasas de supervivencia entre los dos grupos. **Resultados:** Las tasas de remisión completa patológica, mortalidad posoperatoria a 30 días e incidencia de fuga de anastomosis fueron del 39.8% frente al 5.5% ($p < 0.001$), del 2.7% frente al 0% ($p = 0.043$) y del 28.3% frente al 5.5% ($p < 0.001$) en el grupo de QRTN y en el grupo de QTN, respectivamente.

*Correspondence:

Hongni Qin

E-mail: xieyuezheng245@163.com

2444-0507/© 2025 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 09-07-2024

Date of acceptance: 18-02-2025

DOI: 10.24875/CIRUE.M25000900

Cir Cir (Eng). 2025;93(6):620-627

Contents available at PubMed

www.cirurgiaycirujanos.com

*El análisis multivariado mostró que solo la QRTN aumentó significativamente la incidencia de fuga anastomótica (OR: 7.28; IC 95%: 2.44-21.78; $p < 0.001$). **Conclusiones:** Los resultados muestran que la QRTN puede mejorar la tasa de respuesta patológica, pero aumentando la incidencia de fuga anastomótica. Los parámetros de dosis-volumen de radiación no estuvieron asociados con el riesgo de fuga anastomótica.*

Palabras clave: Tumor maligno de esófago. Tratamiento neoadyuvante. Fuga de anastomosis. Factores de riesgo. Quimiorradioterapia.

Introduction

Esophageal carcinoma has a poor prognosis after surgery alone, neoadjuvant chemoradiotherapy (NCRT) is currently considered the standard care for locally advanced esophageal carcinoma^{1,2}. Neoadjuvant chemotherapy (NCT) is also a common practice in multimodality treatment due to the fear of excessive toxicity associated with NCRT, especially after the JCOG1109 NExT study³ and the CMISG1701 trial⁴. Neoadjuvant therapy may increase the incidence of post-operative complications and even mortality⁵. Anastomotic leakage refers to the rupture of the anastomosis after esophageal cancer (EC) surgery, resulting in the leakage of gastric contents⁶. It is one of the most common complications after EC surgery and can lead to serious infection and even death. Radiation therapy in NCRT can damage tissue, affect wound healing, and increase the risk of anastomotic leakage. In addition, NCRT may lead to malnutrition, affect the patient's immune function, increase the risk of infection, and thereby increase the risk of anastomotic leakage⁷. Studies have shown that anastomotic leakage is closely related to both tumor recurrence and poor survival^{8,9}. However, the impact of different neoadjuvant treatment patterns on anastomotic leakage is not well studied.

Radiation therapy can cause damage to microvessels and has a negative effect on irradiated tissue healing. Theoretically, the incidence of anastomotic leakage may be increased when radiotherapy is added to NCT, but rare data were reported from past studies. Most of the studies involve the comparison of anastomotic complications between NCRT and surgery alone, and the conclusions are inconsistent. Some studies have reported a higher incidence of anastomotic leakage in the NCRT group, whereas others have found no significant difference compared with surgery alone. This inconsistency may arise from variations in study design, patient populations, surgical techniques, and definitions of anastomotic leakage¹⁰⁻¹². Unlike EC in

Western countries, which has a high prevalence of esophagogastric junction adenocarcinoma, esophageal squamous cell carcinoma (ESCC) is the most common pathological type in East Asia, and most of the lesions are located in the upper middle portion of the chest. As a result, the radiation field is more closely related to the anastomotic region. In addition, the optimal radiation dose of NCRT has not yet been determined. In recent years, great progress has also been made in the technology of radiotherapy. Based on the above reasons, the risk factors for anastomotic leakage in multimodality treatment should be re-analyzed.

Therefore, clinical data from three prospective trials in neoadjuvant therapy were collected to evaluate whether adding radiotherapy to NCT would increase the incidence of anastomotic leakage, and the effect of radiation dose-volume parameters on rates of anastomotic leakage was also analyzed.

Methods

Clinical data

This study is a retrospective study. A total of 222 cases of clinical data from three prospective clinical studies (NCT01258192, NCT01463501, and NCT03381651) from February 2013 to December 2017 were selected. According to the choice of treatment, the NCT group formed the control group ($n = 109$), and the NRCT group formed the observation group ($n = 113$). The pre-treatment evaluation included chest and upper abdominal enhanced computed tomography (CT) scan and cervical B-ultrasound. Informed consent was waived due to the retrospective nature of the study.

Main inclusion criteria: (1) ECOG PS 0-1; (2) operable or potentially operable (cT2-4N0 or cN1) thoracic ESCC; (3) number of NCT cycles ≥ 2 ; (4) Ivor-Lewis esophagectomy or McKeown esophagectomy (open or laparoscopic) was mandatory; (5) intensity-modulated radiation therapy (IMRT) was mandatory. Exclusion

criteria: (1) salvage esophagectomy after chemoradiotherapy; (2) prophylactic cervical lymph node dissection; (3) jejunal or colon interposition for esophageal replacement; (4) exclude patients with chronic diseases such as malnutrition, diabetes, hypertension, coronary heart disease; (5) exclude patients who have received other anti-tumor treatments.

Data collection

The following data were collected: age, gender (male/female), smoking history (yes/no), drinking (yes/no), lesion location (upper/middle lower of the chest), lesion length, body mass index (BMI) before neoadjuvant treatment, types of neoadjuvant therapy (NCRT/NCT), time from the end of neoadjuvant therapy to surgery, pre-operative BMI, surgical approach (laparoscopic/open), intraoperative blood loss, anastomotic location (cervical/thoracic), residual esophageal length, degree of pathological remission (complete/incomplete), 30-day post-operative mortality rate, radiation prescribed dose, residual esophageal dose (maximum dose and average dose), and anastomotic region dose-volume parameters (V10, V20, V30, V40, V50, average dose, and maximum dose).

Treatment

NEOADJUVANT THERAPY

The platinum-based doublet chemotherapy regimen was delivered once every 3-4 weeks in the NCT group, and few patients received a platinum-based triplet regimen^{13,14}. In the NCRT group, platinum-containing doublet chemotherapy was given weekly (taxane-containing regimen) or once 3-4 weeks (5-Fu-containing regimen) concurrently with thoracic radiotherapy. IMRT was delivered in all patients, GTV included primary tumor and metastatic lymph node (> 5-10 mm in maximum diameter), which was contoured on the planning CT scans. The PTV was defined as GTV plus a 3-4 cm margin superior and inferior to the primary tumor and a 1-1.5 cm radial margin. For patients with lesions in the upper chest or lymph node metastasis in the superior mediastinum, prophylactic irradiation of para-recurrent laryngeal nerve lymph chain and lower neck lymph node was preferred, but the cervical esophagus and the bilateral lower neck should be protected from radiation

as much as possible. The radiation dose was 40-50.4 Gy at 1.8-2 Gy per fraction.

Delineation of the residual esophagus and anastomotic region, and evaluation of dosimetry: a diagnostic CT scan which was performed within half a year after surgery with the same body position and layer thickness as planning CT scan was selected. Anastomosis was recognized in diagnostic CT slices through anastomosis staples and was uniformly expanded by 5 mm to create a so-called "anastomotic region," which was then copied to the corresponding planning CT image. Delineation of the residual esophagus: the upper boundary was at the level of the cricothyroid membrane, and the lower boundary was at the level of the anastomotic region. To reduce the contour error, the delineation of the anastomotic region and residual esophagus were all completed by the first author. The evaluation of anastomotic region dose-volume parameters included V10, V20, V30, V40, V50, mean dose, and maximum dose. The evaluation of residual esophageal dose included the maximum dose and mean dose. The length of the residual esophagus was measured on the CT slice.

SURGERY

Ivor-Lewis esophagectomy or McKeown esophagectomy was performed around 1.5 months after the completion of neoadjuvant therapy. Laparoscopic resection was also allowed. Jejunal or colon interposition for esophageal replacement was not allowed.

Definition of anastomotic leakage

Esophageal-gastric anastomotic leakage was diagnosed through clinical and/or adjuvant examination, including esophagography, digestive endoscopy, and diagnostic CT. The incidence of anastomotic leakage is usually highest within 5-7 days after surgery, but it also occurs in approximately 12% of patients 30 days after surgery. Therefore, we followed up with patients after surgery for EC for 60 days to monitor the occurrence of anastomotic leakage¹⁵.

Definition of pathologic complete remission (pCR) rate

In patients with EC, the pCR rate usually refers to surgical resection after NCT or chemoradiotherapy, and pathological examination finds no residual tumor, including carcinoma *in situ* and invasive carcinoma.

PCR is one of the important indicators to evaluate the effect of neoadjuvant therapy for EC.

Statistical analysis

Detailed statistical analysis of the collected data was conducted using Statistical Package for Social Sciences 18.0 statistical software. Continuous numerical variables that conform to normal distribution are statistically described by mean standard deviation (\bar{x} s), and a two-sample t-test is used for comparison between groups. Numerical variables that do not conform to the normal distribution are described by median (minimum–maximum), and Mann-Whitney U test is used for comparison between groups. Categorical data were statistically described using the number of cases (%), and χ^2 tests were used for comparisons between groups. Univariable logistic regression model and multivariable Cox regression model analysis were used to evaluate risk factors associated with anastomotic leakage. $P < 0.05$ indicates that the difference is statistically significant.

Results

General information description

About 113 patients received NCRT, and 109 received NCT. The median age was 60 years (42–77 years old). There were 60 cases in stage IIA, 26 cases in stage IIB, 122 cases in stage III, and 14 cases in stage IVA. Among them, 38 (17.1%) patients had lesions in the upper thoracic region, 144 (64.9%) in the middle thoracic region, and 40 (18.0%) in the lower thoracic region. Compared with the NCT group, patients in the NCRT group had younger ages, longer time from the end of neoadjuvant therapy to surgery, shorter residual esophagus, less blood loss, and more laparoscopic surgery (Table 1). The dose of radiation was between 3,400 to 5,220 cGy (median radiation dose was 4,500 cGy). 89% of patients received taxane + platinum chemotherapy regimen, and other regimens included 5-Fu+platinum, NVB+ platinum.

Post-operative results between NCRT group and NCT group

The pathological complete remission rate (pCR) in the NCRT group and NCT group was 39.8% and 5.5%, respectively ($p < 0.001$), and the 30-day

post-operative mortality was 2.7% and 0%, respectively, the log-rank test showed that there was a significant difference in survival between the two groups ($p = 0.019$). One patient in the NCRT group died from major bleeding caused by anastomotic leakage (Table 2).

About 38 patients had a post-operative anastomotic leakage, 27 of which were diagnosed by clinical plus adjuvant examination, nine were diagnosed according to clinical examination, and two were confirmed by adjuvant examination. The median time of leakage occurrence after surgery was 13 days (4 days–57 days). The incidence of anastomotic leakage in the NCRT group and NCT group was 28.3% and 5.5%, respectively, with a statistically significant difference ($p < 0.001$) (Table 2).

Univariate and multivariate analysis of clinicopathologic factors associated with anastomotic leakage incidence in whole group

Univariate analysis showed that NCRT ($p < 0.001$), cervical anastomosis ($p = 0.002$), laparoscopic surgery ($p = 0.017$), longer time from the end of neoadjuvant therapy to surgery ($p = 0.010$), and shorter residual esophageal length ($p = 0.001$) were significantly associated with the risk of anastomotic leakage. The multivariate analysis showed that NCRT was an independent risk factor for anastomotic leakage (odds ratio 7.28, 95% confidence interval 2.44–21.78, $p < 0.001$). However, this study only included patients in the NCT and NCRT groups, and the impact of other factors on anastomotic leakage requires further study (Table 3).

Univariate and multivariate analysis of risk factors associated with anastomotic leakage in NCRT group

Further analysis of the dosimetry parameters related to anastomotic leakage in the NCRT group found no significant association between anastomotic leakage incidence and the radiation prescribed dose, anastomotic region dose-volume parameters (V10, V20, V30, V40, V50, mean dose, and maximum dose), and residual esophageal dose (mean dose and maximum dose) based on univariate and multivariate analysis (Table 4).

Table 1. Baseline characteristics between NCRT group and NCT group

Characteristics	NCRT (n = 113)	NCT (n = 109)	u/X ²	p
Gender, n (%)			0.005	0.944
Male	106 (93.8)	102 (93.6)		
Female	7 (6.2)	7 (6.4)		
Smoking, n (%)			0.059	0.808
Yes	84 (74.3)	84 (77.1)		
No	29 (25.7)	25 (22.9)		
Drinking, n (%)			0.073	0.797
Yes	89 (78.8)	85 (78.0)		
No	24 (21.2)	23 (21.1)		
Lesion location, n (%)			0.055	0.815
Upper chest	20 (17.7)	18 (16.5)		
Middle-lower chest	93 (82.3)	91 (83.5)		
cTNM stage, n (%)			1.348	0.717
IIA	27 (23.9)	33 (30.3)		
IIB	13 (11.5)	13 (11.9)		
III	66 (58.4)	56 (51.4)		
IVA	7 (6.2)	7 (6.4)		
Surgery approach, n (%)			51.061	0.000
Open	45 (39.8)	94 (86.2)		
Laparoscopic	68 (60.2)	15 (13.8)		
Anastomotic location, n (%)			110.47	0.000
Cervical	113 (100)	37 (33.9)		
Intrathoracic	0 (0)	72 (66.1)		
Median age (years)	59 (42-77)	61 (43-73)	8.371	0.009
Lesion length (cm), mean ± SE	6.05 ± 0.18	5.94 ± 0.23	0.427	0.191
BMI before neoadjuvant therapies (kg/m ²), mean ± SE	21.82 ± 0.27	22.23 ± 0.27	0.401	0.204
Median time from the end of neoadjuvant therapy to surgery (days)	35 (15-80)	29 (11-55)	11.745	0.000
Pre-operative BMI (kg/m ²), mean ± SE	21.38 ± 0.29	22.00 ± 0.23	2.355	0.060
Intraoperative blood loss (mL), mean ± SE	219.42 ± 13.61	240.54 ± 17.03	3.597	0.034
Residual esophageal length (cm), mean ± SE	3.76 ± 0.09	6.25 ± 0.23	9.471	0.000
TNM stage			0.821	0.365
II	43	48		
III	70	61		
T staging of tumor			0.427	0.514
T2	18	21		
T3	95	88		

cTNM: clinical tumor node metastasis; TNM: tumor node metastasis; BMI: body mass index; SE: standard error.

Table 2. Post-operative results between NCRT group and NCT group

Characteristics	NCRT (n = 113)	NCT (n = 109)	X ²	p
R0, n (%)	106 (93.8)	105 (96.3)	0.751	0.386
pCR, n (%)	45 (39.8)	6 (5.5)	36.927	0.000
pT0, n (%)	61 (54.0)	7 (6.4)	59.063	0.000
pN0, n (%)	66 (58.4)	47 (43.1)	5.189	0.022
Anastomotic leakage, n (%)	32 (28.3)	6 (5.5)	20.354	0.000
30-day post-operative mortality rate, n (%)	3 (2.7)	0 (0)	-	-

NCT: neoadjuvant chemotherapy; NCRT: neoadjuvant chemoradiotherapy; R0: macroscopic complete resection; pCR: pathologic complete remission; pT0: pathologic T0; pN0: pathologic n0.

Discussion

This study compared the impact of NCT and NCRT on anastomotic leakage after EC resection. The results showed that compared with NCT, NCRT significantly increased the rate of pCR but also increased the incidence of anastomotic leakage. Multivariate analysis confirmed that NCRT was an independent risk factor for anastomotic leakage. Further dosimetry analysis showed no significant association between anastomotic leakage risk and radiation dose to the residual esophagus and anastomotic region.

Our findings are similar to those of previous studies^{2,9,15-23}. For example, the CROSS study reported

Table 3. Univariate and multivariate analysis of clinicopathologic factors associated with anastomotic leakage incidence in whole group

Characteristics	Univariate analysis			Multivariate analysis		
	OR	95% CI	p	OR	95% CI	p
Gender, male/female	0.29	0.04-2.28	0.241			
Smoking, yes/no	1.56	0.64-3.76	0.325			
Drinking, yes/no	1.08	0.46-2.53	0.866			
Tumor location, proximal third/middle-lower third	2.23	1.00-5.00	0.050			
Type of neoadjuvant therapy, NCRT/NCT	7.51	3.01-18.75	< 0.001	7.28	2.44-21.78	< 0.001
Surgery approach, open/laparoscopic	0.43	0.21-0.86	0.017	0.47	0.33-1.17	0.144
Anastomosis location, cervical/thoracic	25.07	3.37-186.54	0.002	10.381	0.84-65.41	0.082
Age (years)	0.98	0.94-1.04	0.532			
Lesion length (cm)	1.15	0.99 ± 1.33	0.068			
BMI before neoadjuvant therapies (kg/m ²)	0.94	0.83-1.06	0.301			
Time from the end of neoadjuvant therapy to surgery (days)	1.04	1.01-1.07	0.010	1.28	0.87-1.36	0.085
Pre-operative BMI (kg/m ²)	0.98	0.87-1.11	0.767			
Intraoperative blood loss (mL)	1.00	1.00-1.00	0.288			
Residual esophageal length (cm)	0.71	0.58-0.87	0.001	0.62	0.64-1.22	0.064

OR: odds ratio; CI: confidence interval; BMI: body mass index.

an anastomotic leakage rate of 22% in the NCRT group². A retrospective study of 686 ESCC patients in Japan showed that the incidences of anastomotic leakage were 27.9% in the NCRT group (376 patients with a radiation dose of 30-42 Gy) and 16.5% in the surgery group (310 patients)⁹. The incidence of anastomotic leakage in the NCT group was similar to that reported in the surgery-alone groups in other studies¹⁷⁻¹⁹, suggesting that NCT does not significantly increase the incidence of anastomotic leakage.

The current results suggest that factors other than NCRT and tumor size may contribute to anastomotic leakage. Although the dose parameters V20, V30, and V40 have been shown to be associated with anastomotic leakage, the underlying mechanisms are unclear. Patient-specific factors such as comorbidities or nutritional status may play a role. The surgical technique itself, such as the type of anastomosis or the use of drains, may also have an impact.

The relationship between radiation dose and anastomotic leakage is more complicated. Despite the high pre-operative radiation dose given (50.4 Gy/28 F), the CALGB9781 study reported a low leakage incidence of 8%²², which was similar to that in the NEOCRTEC5010

study (8.6%)¹⁶, but lower than that in the CROSS study (22%)². A lower irradiation dose was prescribed in both NEOCRTEC 5010 (40Gy) and CROSS study (41.4Gy). Our study also found a non-significant relationship between radiation dose and leakage. When the influence of radiation dose on leakage was investigated, the relationship between radiation fields and the location of anastomosis should also be considered. A retrospective study from MD Anderson revealed that the incidence of anastomotic leakage was 11% in NCRT (radiation dose: 50.4Gy/28F), and the incidence of leakage was increased significantly (39% vs. 2.6%) when the anastomosis was placed within the pre-operative radiation field²⁴. Another study from France showed that although the anastomosis was within the radiation field, the leakage incidence did not rise significantly when thoracic anastomosis was performed¹⁷. Since a wide range of irradiation doses was delivered in NCRT and the spatial relationship between the radiation field and the anastomosis location was sometimes difficult to define accurately, it is more reasonable to explore the relationship between radiation dose of the anastomotic region and leakage incidence. Our results indicated

Table 4. Univariate analysis of risk factors associated with anastomotic leakage in NCRT group

Characteristics	Univariate analysis		
	OR	95% CI	p
Gender, male/female	2.56	0.30-22.13	0.393
Smoking, yes/no	1.19	0.45-3.17	0.727
Drinking, yes/no	1.44	0.54-3.84	0.461
Tumor location, proximal third/ middle-lower third	1.84	0.67-5.03	0.234
Surgery approach, open/laparoscopic	1.13	0.50-2.57	0.773
Age (years)	1.02	1.00-1.09	0.427
Lesion length (cm)	1.15	0.94-1.42	0.181
BMI before neoadjuvant therapies (kg/m ²)	0.99	0.86-1.13	0.846
Time from the end of neoadjuvant therapy to surgery (days)	1.02	0.98-1.05	0.349
Pre-operative BMI (kg/m ²)	1.04	0.91-1.18	0.610
Intraoperative blood loss (mL)	1.00	1.00-1.01	0.038
Residual esophageal length (cm)	1.12	0.73-1.74	0.601
Anastomotic region			
V10	1.00	0.99-1.01	0.893
V20	1.00	0.99-1.01	0.891
V30	1.00	0.99-1.01	0.592
V40	1.00	0.99-1.01	0.432
V50	0.99	0.99-1.01	0.298
Average dose	1.00	1.00-1.00	0.565
Maximum dose	1.00	1.00-1.00	0.586
Residual esophagus			
Average dose	1.00	1.00-1.00	0.586
Maximum dose	1.00	1.00-1.00	0.537
Radiation dose	1.00	1.00-1.00	0.968

NCRT: neoadjuvant chemoradiotherapy; OR: odds ratio; CI: confidence interval; BMI: body mass index.

that there was no significant relationship between the dose-volume parameters of the anastomotic region and leakage incidence. In MD Anderson's study, the anastomotic region should have been exposed to a dose of $\geq 50\text{Gy}$ in those patients whose anastomosis was completely placed within the irradiation field. In our study, we tried to protect the lower neck or cervical esophagus from being irradiated when a higher dose was prescribed, and neck anastomosis was performed in all patients receiving NCRT. Thus, the average dose in the anastomosis was only 37Gy, and V40, V45, and V50 were 64%, 41%, and 22%, respectively, suggesting that good communication between the

surgeon and radiotherapist is indispensable in irradiation field design.

Our study had some limitations, as all retrospective studies did. Although the anastomotic region was delineated by the same investigator, there still might be some deviations in identifying the location of the anastomosis and delineating it on the corresponding planning CT slices, especially for patients with manual anastomosis. Moreover, mixed factors were thought to work together in the development of leakage, posing some potential confounding factors that could not be included in this retrospective analysis. Furthermore, the relatively few studies included in our analysis may limit the generalizability of our findings. Future studies of a larger sample size and a more diverse patient population are needed to further validate our results.

Conclusion

Our results confirmed that NCRT could improve the pathologic response rate but also increase the incidence of anastomotic leakage simultaneously. However, neither the prescribed radiation dose nor the doses to the anastomotic site and residual esophageal tissue were significantly associated with anastomotic leakage risk.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of Humans and Animals. The authors declare that the procedures followed complied with the ethical standards of the responsible human experimentation committee and adhered to the World Medical Association and the Declaration of Helsinki. The procedures were approved by the institutional Ethics Committee.

Confidentiality, Informed Consent, and Ethical Approval. The authors have followed their institution's confidentiality protocols and obtained informed consent from patients. The authors have obtained approval from the Ethics Committee for the analysis of routinely

obtained and anonymized clinical data, so informed consent was not necessary. Relevant guidelines were followed.

Declaration on the Use of Artificial Intelligence.

The authors declare that no generative artificial intelligence was used in the writing of this manuscript.

References

1. Yang H, Liu H, Chen Y, Zhu C, Fang W, Yu Z, et al. Long-term efficacy of neoadjuvant chemoradiotherapy plus surgery for the treatment of locally advanced esophageal squamous cell carcinoma: the NEOCR-TEC₅₀₁₀ randomized clinical trial. *JAMA Surg.* 2021;156:721-9.
2. Van Hagen P, Hulshof MC, Van Lanschot JJ, Steyerberg EW, Van Berge Henegouwen MI, Wijnhoven BP, et al. Preoperative chemoradiotherapy for esophageal or junctional cancer. *N Engl J Med.* 2012;366:2074-84.
3. Kato K, Machida R, Ito Y, Daiko H, Ozawa S, Ogata T, et al. A randomized controlled phase III trial comparing two chemotherapy regimens and chemoradiotherapy regimen as neoadjuvant treatment for locally advanced esophageal cancer, JCOG1109 NExT study. *J Clin Oncol.* 2022;40:238.
4. Tang H, Wang H, Fang Y, Zhu JY, Yin J, Shen YX, et al. Neoadjuvant chemoradiotherapy versus neoadjuvant chemotherapy followed by minimally invasive esophagectomy for locally advanced esophageal squamous cell carcinoma: a prospective multicenter randomized clinical trial. *Ann Oncol.* 2023;34:163-72.
5. Kumagai K, Rouvelas I, Tsai JA, Mariosa D, Klevebro F, Lindblad M, et al. Meta-analysis of postoperative morbidity and perioperative mortality in patients receiving neoadjuvant chemotherapy or chemoradiotherapy for resectable oesophageal and gastro-oesophageal junctional cancers. *Br J Surg.* 2014;101:321-38.
6. Papadakos SP, Argyrou A, Katsaros I, Lekakis V, Mpouga G, Vergadis C, et al. The impact of EndoVAC in addressing post-esophagectomy anastomotic leak in esophageal cancer management. *J Clin Med.* 2024;13:7113.
7. Zhao Y, Dai Z, Min W, Sui X, Kang H, Zhang Y, et al. Perioperative versus preoperative chemotherapy with surgery in patients with resectable squamous cell carcinoma of esophagus: a phase III randomized trial. *J Thorac Oncol.* 2015;10:1349-56.
8. Kofoed SC, Calatayud D, Jensen LS, Helgstrand F, Achiam MP, De Heer P, et al. Intrathoracic anastomotic leakage after gastroesophageal cancer resection is associated with increased risk of recurrence. *J Thorac Cardiovasc Surg.* 2015;150:42-8.
9. Andreou A, Biebl M, Dadras M, Struecker B, Sauer IM, Thuss-Patience PC, et al. Anastomotic leak predicts diminished long-term survival after resection for gastric and esophageal cancer. *Surgery.* 2016;160:191-203.
10. Morita M, Masuda T, Okada S, Yoshinaga K, Saeki H, Tokunaga E, et al. Preoperative chemoradiotherapy for esophageal cancer: factors associated with clinical response and postoperative complications. *Anticancer Res.* 2009;29:2555-62.
11. Koëter M, Van Der Sangen MJ, Hurkmans CW, Luyer MD, Rutten HJ, Nieuwenhuijzen GA. Radiation dose does not influence anastomotic complications in patients with esophageal cancer treated with neoadjuvant chemoradiation and transhiatal esophagectomy. *Radiat Oncol (London, England).* 2015;10:59.
12. Morse RT, Mouw TJ, Moreno M, Erwin JT, Cao Y, DiPasco P, et al. Neoadjuvant radiotherapy facility type affects anastomotic complications after esophagectomy. *J Gastrointest Surg.* 2023;27:1313-20.
13. Zhao X, Ren Y, Hu Y, Cui N, Wang X, Cui Y. Neoadjuvant chemotherapy versus neoadjuvant chemoradiotherapy for cancer of the esophagus or the gastroesophageal junction: a meta-analysis based on clinical trials. *PLoS One.* 2018;13:e0202185.
14. Makary MS, Khandpur U, Cloyd JM, Mumtaz K, Dowell JD. Locoregional therapy approaches for hepatocellular carcinoma: recent advances and management strategies. *Cancers (Basel).* 2020;12:1914.
15. Li H, Zhuang S, Yan H, Wei W, Su Q. Risk factors of anastomotic leakage after esophagectomy with intrathoracic anastomosis. *Front Surg.* 2021;8:743266.
16. Yang H, Liu H, Chen Y, Zhu C, Fang W, Yu Z, et al. Neoadjuvant chemoradiotherapy followed by surgery versus surgery alone for locally advanced squamous cell carcinoma of the esophagus (NEOCR-TEC5010): a phase III multicenter, randomized, open-label clinical trial. *J Clin Oncol.* 2018;36:2796-803.
17. Gronnier C, Tréchet B, Duhamel A, Mabrut JY, Bail JP, Carrere N, et al. Impact of neoadjuvant chemoradiotherapy on postoperative outcomes after esophageal cancer resection: results of a European multicenter study. *Ann Surg.* 2014;260:764-70; discussion 770-1.
18. Medical Research Council Oesophageal Cancer Working Group. Surgical resection with or without preoperative chemotherapy in oesophageal cancer: a randomised controlled trial. *Lancet.* 2002;359:1727-33.
19. Ma S, Yan T, Liu D, Wang K, Wang J, Song J, et al. Neoadjuvant chemotherapy followed by minimally invasive esophagectomy is safe and feasible for treatment of esophageal squamous cell carcinoma. *Thorac Cancer.* 2018;9:310-5.
20. Briel JW, Tamhankar AP, Hagen JA, DeMeester SR, Johansson J, Choustoulakis E, et al. Prevalence and risk factors for ischemia, leak, and stricture of esophageal anastomosis: gastric pull-up versus colon interposition. *J Am Coll Surg.* 2004;198:536-41; discussion 541-2.
21. Chao YK, Chan SC, Chang HK, Liu YH, Wu YC, Hsieh MJ, et al. Salvage surgery after failed chemoradiotherapy in squamous cell carcinoma of the esophagus. *Eur J Surg Oncol.* 2009;35:289-94.
22. Tepper J, Krasna MJ, Niedzwiecki D, Hollis D, Reed CE, Goldberg R, et al. Phase III trial of trimodality therapy with cisplatin, fluorouracil, radiotherapy, and surgery compared with surgery alone for esophageal cancer: CALGB 9781. *J Clin Oncol.* 2008;26:1086-92.
23. Wang H, Tang H, Fang Y, Tan L, Yin J, Shen Y, et al. Morbidity and mortality of patients who underwent minimally invasive esophagectomy after neoadjuvant chemoradiotherapy vs neoadjuvant chemotherapy for locally advanced esophageal squamous cell carcinoma: a randomized clinical trial. *JAMA Surg.* 2021;156:444-51.
24. Juloori A, Tucker SL, Komaki R, Liao Z, Correa AM, Swisher SG, et al. Influence of preoperative radiation field on postoperative leak rates in esophageal cancer patients after trimodality therapy. *J Thorac Oncol.* 2014;9:534-40.

Evaluation of mediastinal lymphadenopathy from long-term radiological findings in COVID-19

Evaluación de la linfadenopatía mediastínica a partir de hallazgos radiológicos a largo plazo en COVID-19

Gulhan Ayhan-Albayrak*¹ and Mustafa I. Bardakci¹

Department of Chest Diseases, Sisli Hamidiye Etfal Training and Research Hospital, University of Health Sciences, Istanbul, Turkey

Abstract

Objective: Mediastinal lymphadenopathy (MLAP) has been reported in post-COVID-19 patients. In this study, the relationship between post-COVID-19 infection and MLAP was investigated in patients who had been diagnosed with MLAP and decided to undergo surgery. **Methods:** The study included the records of 140 patients who had been diagnosed with MLAP and were decided for surgical treatment. Demographic findings, reverse transcription-polymerase chain reaction (PCR) test results, chest X-ray, thorax computed tomography (CT) findings, positron emission tomography (PET)-CT findings, and histopathological results were recorded. **Results:** SUV_{max} value above 2.5 was 15 times more common in patients with positive PCR test results than in patients with negative results. Abnormal chest X-ray results were associated with a 9.3-fold increase in the number of patients, and the number of patients with abnormal pathology results was 33.9 times higher than those with normal results. **Conclusions:** Post-COVID-19 and MLAP (SUV_{max} 3-5) were shown to be associated independently of age, gender, comorbidities, and disease outcomes. MLAP lesions in patients with COVID-19 demonstrated SUV_{max} values that were 10-fold higher compared to patients without COVID-19. Determining reliable SUV_{max} values in patients with severe COVID-19 may help guide clinical decisions, tailor therapeutic approaches, and avoid unnecessary surgical indications.

Keywords: Mediastinal lymphadenopathy. SUV_{max} . Surgical indication. Positron emission tomography. Post-COVID-19.

Resumen

Objetivo: Se ha informado linfadenopatía mediastínica en pacientes que se recuperan de COVID-19. Nuestro objetivo fue investigar si la linfadenopatía mediastínica encontrada en pacientes con neumonía grave por COVID-19 al menos 6 meses después de la infección está asociada con la enfermedad. **Métodos:** El estudio incluyó los registros de 140 pacientes a quienes se les había diagnosticado MLAP y se decidió que recibirían tratamiento quirúrgico. Los hallazgos demográficos se compararon mediante los resultados de RT-PCR, TC de tórax y PET-CT. **Resultados:** Un valor de SUV_{max} superior a 2,5 fue 15 veces más frecuente en pacientes con resultados positivos en la prueba PCR que en aquellos con resultados negativos. Los resultados anormales en la radiografía de tórax se asociaron con un aumento de 9,3 veces en el número de pacientes, y el número de pacientes con resultados anormales en las pruebas de anatomía patológica fue 33,9 veces mayor que el de aquellos con resultados normales. **Conclusiones:** Se demostró que la presencia de MLAP (SUV_{max} 3-5) tras la COVID-19 se asocia independientemente de la edad, el sexo, las comorbilidades y la evolución de la enfermedad. Las lesiones de MLAP en pacientes con COVID-19 mostraron valores de SUV_{max} 10 veces superiores a los de los pacientes sin COVID-19. Determinar valores fiables de SUV_{max} en pacientes con COVID-19 grave puede ayudar a orientar las decisiones clínicas, personalizar los enfoques terapéuticos y evitar indicaciones quirúrgicas innecesarias.

Palabras clave: Linfadenopatía mediastínica. SUV_{max} . Indicación quirúrgica. Tomografía por emisión de positrones. Post-COVID-19.

*Correspondence:

Gulhan Ayhan-Albayrak
E-mail: dr.gulhanayhan@hotmail.com

Date of reception: 09-01-2025
Date of acceptance: 20-08-2025
DOI: 10.24875/CIRUE.M25000901

Cir Cir (Eng). 2025;93(6):628-635
Contents available at PubMed
www.cirugiaycirujanos.com

2444-0507/© 2025 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Mediastinal lymph node enlargement is a critical computed tomography (CT) finding that can range from benign to malignant in patients with respiratory complaints and may require further examination. Since the presence of mediastinal lymphadenopathy (MLAP) is related to the severity of the disease in post-COVID patients who have had and recovered from severe COVID-19 infection, the presence of MLAP should be carefully examined in subsequent examinations, even in patients without lung involvement on admission CT¹. The detection of MLAP in patients with idiopathic pulmonary fibrosis (IPF) is thought to be a result of high-grade chronic inflammation, and a correlation between the severity of IPF disease and MLAP has been shown in the literature².

When the current literature on MLAP and COVID-19 were examined, studies were conducted in the acute and active period of COVID-19 pneumonia. However, in our study, MLAP in the controls was performed 6-12 months after the patients had COVID-19 pneumonia. After suffering severe COVID-19 pneumonia, the mediastinal lymph nodes begin to enlarge during acute inflammation, which is considered a poor-prognosis marker at that time. After 6 months, this turns into chronic inflammation, and MLAP occurs. It is thought that there may be a correlation between the severity of chronic inflammation and mediastinal lymph node enlargement in patients with severe COVID-19 pneumonia, just as in IPF.

MLAP can be caused by infectious or non-infectious causes and is defined as a measurement of the lymph node short axis ≥ 10 mm. SUV_{max} value ≥ 2.5 was defined as pathological MLAP³. The most common atypical CT features of COVID-19 pneumonia are MLAP, linear opacities, tree bud sign, interlobular and intralobular septal thickening, cavitation, and pleural effusion⁴. MLAP is frequently associated with benign diseases such as heart failure, sarcoidosis, and diseases accompanying malignant diseases. Although MLAP is not the typical chest finding in COVID-19 pneumonia, it is associated with the prognosis of COVID-19 patients. The mortality rate in hospitalized COVID-19 patients with MLAP was found to be higher than in those without MLAP, and it has been reported in the literature that it should be investigated as a prognostic factor for severe disease. The literature shows a correlation between the severity of IPF disease and the presence of MLAP, suggesting that

MLAP results from severe chronic inflammation in these patients^{1,5,6}.

For MLAP, the size and number of lymph nodes are essential for surgical treatment and follow-up. MLAP, usually over 2 cm in size, should be closely monitored regardless of the cause⁶. The radiological findings of one of our patients are shown in figure 1.

COVID-19 is a respiratory disease caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection. In this study, we aimed to determine whether MLAP, which was detected in patients with severe COVID-19 pneumonia in the controls at least 6 months after the disease (post-COVID-19 period), was related to having the disease. If the detected MLAP formation is determined to be secondary to COVID-19 pneumonia, then immediate referral to the surgery clinic for further examination may be unnecessary. At the same time, by closely monitoring the patients in whom we detect MLAP, we can protect them from surgical complications and not be late in making the diagnosis.

Methods

This study was approved by the Ethical Committee of the University of Health Sciences Şişli Hamidiye Etfal Training and Research Hospital. Ethics Committee approval was granted by our institution on September 06, 2022, protocol number 2137. It was conducted according to the Declaration of Helsinki and Good Clinical Practice. The authors have obtained the approval of the Ethics Committee for the analysis and publication of clinical data obtained routinely. The informed consent of the patients was not required because it was a retrospective observational study. All procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1975, as revised in 2008.

Study population and data collection

Our study was a single-center, retrospective, case-control study. Our study analyzed the data of 140 patients who applied to our hospital's Şişli Hamidiye Etfal Training and Research Hospital Chest Diseases Outpatient Clinic between January 01, 2022, and August 01, 2022, and were reviewed retrospectively. During etiological investigations, individuals over 18 years old, regardless of gender, who applied to our hospital's chest diseases outpatient clinics with the MLAP, who had not been diagnosed with lymphoma and sarcoidosis, immunodeficiency, cancer,




Figure 1. A view from the chest radiograph of the patients*. **A:** a section with Hilar fullness from the chest X-ray findings of the patients. **B:** a section from the mediastinal window view with mediastinal lymphadenopathy. **C:** a section with positive fluorodeoxyglucose from positron emission tomography-computed tomography findings. *It was taken from the hospital file archives of the patients in our study, and their permission was obtained.

tuberculosis, asthma, chronic obstructive pulmonary disease, and had no previous mediastinal pathology. Patients with MLAP detected in thorax CT examinations were included in the study. Patients with lung and other malignancies, under 18 years of age, pregnant women, and patients who had image artifacts on their CT scans were excluded from the study. A total of 140 cases who were followed up in our clinic and diagnosed with chest MLAP by CT were included in the study. These cases were divided into two categories: 83 cases with COVID-19 who had COVID-19 pneumonia and recovered, and 57 cases without COVID-19. The two categories were compared using data obtained from demographic characteristics, clinical findings, reverse transcription-polymerase chain reaction (RT-PCR) results, chest radiography findings, thorax CT findings, positron emission tomography (PET)-CT findings, histopathological results, and treatment approaches. Data were collected from the hospital automation system and analyzed with Statistical Package for the Social Sciences (SPSS) 24.

Sample size

The sample of the study was determined as a total of 140 patients, at least 51 in both groups, with an effect size of 0.40, $\alpha = 0.05$, and power $(1-\beta) = 0.85$, using the G-power program. There were two groups: 57 patients without covid-19 and 83 patients with RT-PCR (+)

Image analysis

Demographic information, clinical findings, comorbidities, thorax CT, and PET-CT images were obtained

from hospital computer records and Picture Archiving and Communication System. RT-PCR testing was performed using nasopharyngeal or oropharyngeal swabs using the Biospeedy SARS-CoV-2 Dual Gene RT-qPCR Kit (Bioeksen). Chest CT scans were performed with patients in the supine position, during last inspiration, and without intravenous contrast administration. The protocol included a peak voltage of 100 kV and an effective milliampere-second setting of 20 mA-s, resulting in unenhanced chest CT scans. These scans used a 512×512 pixel image matrix with 2 mm thin sections taken in the axial plane.

In this study, adjustments were applied to the mediastinal windows (width: 400 HU; level: 100 HU) and lung windows (width: 1500 HU; level: -500 HU). Tomography findings were evaluated by two experienced radiologists who agreed on the CO-RADS value. In routine axial CT images, the short axis of the mediastinal lymph node was measured, and if the short axis was 10 mm or more, it was considered MLAP. When the SUVmax value was ≥ 2.5 in the PET-CT examination of patients diagnosed with MLAP, fluorodeoxyglucose (FDG) uptake was considered pathological and was called "FDG positive".

Statistical analysis

Patient data collected within the scope of the study were analyzed with the IBM SPSS for Windows 23.0 (IBM Corp., Armonk, NY) package program. Frequency and percentage were given for categorical data, and median, minimum, and maximum descriptive values for continuous data. The normality test of the data was conducted with the Kolmogorov-Smirnov Test. In the comparisons between the groups, the "Independent

Table 1. Distribution of demographic characteristics of all patients according to MLPA (lymph node short axis, SUV_{max} value)

Characteristics	Lymph node short axis			p	SUV _{max}			p
	10-20 (n = 103)	21-30 (n = 27)	> 30 (n = 10)		< 2.5 (n = 22)	2.5-5.0 (n = 51)	> 5.0 (n = 49)	
	n (%)	n (%)	n (%)		n (%)	n (%)	n (%)	
Age (years)				0.553				0.432
18-64	62 (60.2)	19 (70.4)	7 (70)		16 (72.7)	33 (64.7)	28 (57.1)	
65-94	41 (39.8)	8 (29.6)	3 (30)		6 (27.3)	18 (35.3)	21 (42.9)	
Gender				0.696				0.428
Male	64 (62.1)	15 (55.6)	7 (70)		16 (72.7)	29 (56.9)	29 (59.2)	
Female	39 (37.9)	12 (44.4)	3 (30)		6 (27.3)	22 (43.1)	20 (40.8)	
Smoking status				0.988				0.276
Smoker	44 (42.7)	13 (48.1)	4 (40)		13 (59.1)	22 (43.1)	20 (40.8)	
Non-smoker	38 (36.9)	9 (33.3)	4 (40)		8 (36.4)	21 (41.2)	17 (34.7)	
Ex-smoker	21 (20.4)	5 (18.5)	2 (20)		1 (4.5)	8 (15.7)	12 (24.5)	
Symptoms				0.058				0.275
Dyspnea	17 (16.5)	1 (3.7)	5 (50)		4 (18.2)	5 (9.8)	12 (24.5)	
Chest pain	14 (13.6)	5 (18.5)	1 (10)		1 (4.5)	9 (17.6)	7 (14.3)	
Cough	61 (59.2)	17 (63)	4 (40)		13 (59.1)	33 (64.7)	26 (53.1)	
Back pain	11 (10.7)	4 (14.8)	0 (0)		4 (18.2)	4 (7.8)	4 (8.2)	
Concomitant disease				0.118				0.950
Asthma	36 (38.7)	4 (15.4)	2 (20)		5 (27.8)	13 (26)	15 (34.1)	
DM	7 (7.5)	3 (11.5)	0 (0)		1 (5.6)	5 (10)	2 (4.5)	
GIS	5 (5.4)	3 (11.5)	2 (20)		2 (11.1)	6 (12)	2 (4.5)	
HT	13 (14)	4 (15.4)	1 (10)		3 (16.7)	7 (14)	8 (18.2)	
HD	6 (6.5)	2 (7.7)	3 (30)		2 (11.1)	5 (10)	3 (6.8)	
COPD	26 (28)	10 (38.5)	2 (20)		5 (27.8)	14 (28)	14 (31.8)	

DM: diabetes mellitus; GISD: gastrointestinal system diseases; HT: hypertension; COPD: chronic obstructive pulmonary disease; HD: heart disease; MLPA: mediastinal lymphadenopathy.

Sample T-Test” was used for those with normal distribution for the two groups, the “Mann-Whitney U Test” for those who did not show normal distribution, and the “Fisher’s Exact Test or Chi-Square Test” was used for the comparison of categorical variables. Logistic regression analysis was used to examine the risk factors affecting the SUV_{max} value of ± 2.5 . The results were considered statistically significant when $p < 0.05$.

Results

A total of 140 patients, 38.6% (n = 54) female and 61.4% (n = 86) male, were included in the evaluation. 40.7% (n = 57) of the patients were diagnosed without COVID-19, and 59.3% (n = 83) had positive RT-PCR test results, and it was determined that they had COVID-19. The distribution of demographic characteristics of all patients according to MLAP (lymph node short axis, SUV_{max} value) is shown in table 1. When table 1 was examined, no statistically significant difference was detected between demographic characteristics and mediastinal lymph node short axis and SUV_{max} values ($p > 0.05$).

The distribution of demographic characteristics of the patients with COVID-19 and without COVID-19 is denoted in table 2. When the table was examined, there was only a statistically significant difference between the two groups regarding age ($p < 0.05$).

Significant differences were observed between the two groups in all clinical findings ($p < 0.05$) (Table 3). Peripheral lymphadenopathy was in the intra-abdominal region in 24.1% (n = 20) of RT-PCR-positive with COVID-19 patients and 3.5% (n = 2) of those without COVID-19 patients ($p < 0.001$). Mediastinal width and lymph node size were higher in COVID-19 patients than in non-COVID-19 patients ($p = 0.008$; $p < 0.001$). The pathology result of the majority of COVID-positive patients with COVID-19 was benign (65%), whereas this rate was 22.8% in patients without COVID-19 ($p < 0.001$). As a result, 77.2% of the patients without COVID-19 PCR results regressed in the follow-up, whereas this rate was 24.1% in the patients with COVID-19 ($p < 0.001$).

Table 4 elaborates on the logistic regression analysis results, examining the factors causing the SUV_{max}

Table 2. Distribution of demographic characteristics of patients with COVID-19 and without COVID-19

Characteristics	Total (n = 140)	without COVID-19 (n = 57)	with COVID-19 (n = 83)	p
	n (%) or Median (Min-Max)	n (%) or Median (Min-Max)	n (%) or Median (Min-Max)	
Age (years)	61 (20-94)	55 (22-78)	62 (20-94)	0.006
Gender				0.477
Male	86 (61.4)	33 (57.9)	53 (63.9)	
Female	54 (38.6)	24 (42.1)	30 (36.1)	
Smoking status				0.115
Smoker	61 (43.6)	25 (43.9)	36 (43.4)	
Non-smoker	51 (36.4)	25 (43.9)	26 (31.3)	
Exsmoker	28 (20)	7 (12.3)	21 (25.3)	
Symptoms				0.656
Dyspnea	23 (16.4)	8 (14)	15 (18.1)	
Chest pain	20 (14.3)	7 (12.3)	13 (15.7)	
Cough	82 (58.6)	34 (59.6)	48 (57.8)	
Back pain	15 (10.7)	8 (14)	7 (8.4)	
Concomitant disease	129 (92.1)	50 (87.7)	79 (95.2)	0.122
Asthma	42 (32.6)	19 (38)	23 (29.1)	
DM	10 (7.8)	5 (10)	5 (6.3)	
GIS	10 (7.8)	4 (8)	6 (7.6)	
HT	18 (14)	8 (16)	10 (12.7)	
HD	11 (8.5)	2 (4)	9 (11.4)	
COPD	38 (29.5)	12 (24)	26 (32.9)	

DM: diabetes mellitus; GISD: gastrointestinal system diseases; HT: hypertension; COPD: chronic obstructive pulmonary disease; HD: heart disease.

value to be ≥ 2.5 in the patients included in the study. COVID-19 patients caused a 15-fold SUV_{max} value of ≥ 2.5 . In addition, peripheral lymphadenopathy caused a 3.9-fold SUV_{max} value of ≥ 2.5 , abnormal chest X-rays caused a 9.3-fold SUV_{max} value of ≥ 2.5 , and abnormal pathology results led to a 33.9-fold SUV_{max} value of ≥ 2.5 .

The statistically significant variables in the univariate model were re-evaluated in the multivariate model. RT-PCR positivity with COVID-19 and abnormal pathology findings were statistically significant. Accordingly, it was determined that positive RT-PCR test results with COVID-19 were 5.9 times higher than without COVID-19 outcomes, and abnormal pathology results were found to be 13.4 times more frequently in SUV_{max} value being ≥ 2.5 compared to normal samples.

Discussion

In the study, MLAP was detected in the 6-month follow-up of patients with severe COVID-19, and the

aim was to evaluate the relationship between these MLAP SUV_{max} values and having COVID-19 disease. While the frequency of MLAP in patients with COVID-19 pneumonia is 0-66% in the literature, the frequency of MLAP in this study (965 patients with COVID-19 pneumonia were scanned, and MLAP was detected in 83 and included in the study) was found to be 11.6%, similar to the literature^{1,6,7}.

Thorax tomography and PET-CT scans were performed on our patients who applied to us with respiratory complaints at least 6 months to 1 year after COVID-19, that is, during the post-COVID-19 patient controls. Diagnostic PET-CT was performed in patients with MLAP detected in thorax CT examination. In addition, we evaluated that MLAP size (SUV_{max} value) and FDG uptake are not associated with malignancy in patients with severe COVID-19, but may be associated with the severity of chronic inflammation.

Our results showed that mediastinal width and lymph node size were higher in patients with COVID-19 than in those without COVID-19. The pathology result of most COVID-19-positive patients was benign (65%), whereas this rate was 22.8% in patients without COVID-19. As a result, 77.2% of the patients without COVID-19 regressed in the follow-up, whereas this rate was 24.1% in the patients with COVID-19.

COVID-19-positive patients included in the study increased the incidence of $MLAP \geq 2.5$ above the SUV_{max} value 15 times more than those without COVID-19. In addition, detecting peripheral lymphadenopathy in PET-CT increased the incidence of ≥ 2.5 times above the SUV_{max} value, 3.9 times more. Abnormal chest X-rays of the patients increased the incidence of ≥ 2.5 above the SUV_{max} value 9.3 times, and abnormal pathology results increased the incidence of ≥ 2.5 above the SUV_{max} value 33.9 times.

In our study, the chest X-ray findings of the patients without COVID-19 were mostly expected. In contrast, the chest X-ray of the patients with COVID-19-positive patients was found to be more abnormal, resulting in mediastinal enlargement and hilar lymphadenopathies⁸⁻¹⁰. The most common thorax CT finding detected in our study was M4R right lower paratracheal lymph node enlargement. Pulmonary parenchyma findings were deficient, and sequelae were reported as changes¹¹.

Regarding the results of our study, the presence of aggravated COVID-19 is linked with the occurrence of MLAP. COVID-19-positive patients and abnormal pathology results have indicated higher SUV_{max} values, thus the presence of lymphadenopathy. To date, MLAP has been observed widely in COVID-19 (0-66%).

Table 3. Distribution of clinical characteristics of patients with COVID-19 and without COVID-19

Characteristics	Total (n = 140)	Without COVID-19 (n = 57)	With COVID-19 (n = 83)	p
	n (%) or Median (Min-Max)	n (%) or Median (Min-Max)	n (%) or Median (Min-Max)	
Peripheral LAP				< 0.001
None	55 (39.3)	35 (61.4)	20 (24.1)	< 0.001
Cervical	43 (30.7)	15 (26.3)	28 (33.7)	0.454
Intraabdominal	22 (15.7)	2 (3.5)	20 (24.1)	< 0.001
Pelvic	4 (2.9)	1 (1.8)	3 (3.6)	0.646
Diffuse	16 (11.4)	4 (7)	12 (14.5)	0.276
Chest X-ray				0.002
Normal	78 (55.7)	42 (73.7)	36 (43.4)	< 0.001
Enlarged mediastinum	50 (35.7)	13 (22.8)	37 (44.6)	0.008
Hilar fullness	12 (8.6)	2 (3.5)	10 (12)	0.122
Computerize tomografi				0.024
M2R right upper paratracheal	39 (27.9)	14 (24.6)	25 (30.1)	0.597
M3 prevascular	20 (14.3)	12 (21.1)	8 (9.6)	0.099
M4R right lower paratracheal	39 (27.9)	21 (36.8)	18 (21.7)	0.049
M4L lower left paratracheal	14 (10)	3 (5.3)	11 (13.3)	0.207
M7 subcarinal	28 (20)	7 (12.3)	21 (25.3)	0.093
Lymph node size (mm)	17 (10-51)	15 (10-44)	18 (10-51)	< 0.001
SUV _{max}				< 0.001
FDG (-)	18 (12.9)	15 (26.3)	3 (3.6)	< 0.001
< 2.5	22 (15.7)	18 (31.6)	4 (4.8)	< 0.001
2.5-5.0	51 (36.4)	15 (26.3)	36 (43.4)	0.039
> 5.0	49 (35)	9 (15.8)	40 (48.2)	< 0.001
Pathology				< 0.001
None	57 (40.7)	40 (70.2)	17 (20.5)	< 0.001
Benign	67 (47.9)	13 (22.8)	54 (65.1)	< 0.001
Malign	16 (11.4)	4 (7)	12 (14.5)	0.276
Pathology result				< 0.001
Granulomatous lymphadenitis (benign)	20 (14.3)	4 (7)	16 (19.3)	0.073
Regressed at follow-up (benign)	64 (45.7)	44 (77.2)	20 (24.1)	< 0.001
Reactive lymph node (benign)	37 (26.4)	5 (8.8)	32 (38.6)	< 0.001
Anthracosis (benign)	3 (2.1)	0 (0)	3 (3.6)	0.271
Cancer metastasis (malignant)	16 (11.4)	4 (7)	12 (14.5)	0.276

LAP: lymphadenopathy; FDG: fluorodeoxyglucose.

However, it should be emphasized that not all lymphadenopathies are located in the mediastinal region. In published case reports, two cases (both female) are reported to have MLAP at the initial and repeated CT scan on the 6th day (1/6/10R and 2R/4R/4L)^{12,13}.

However, there are controversial data on this subject. Bayramoglu et al. indicated the presence of MLAP as 0-8.1% in their retrospective analysis¹⁴. In another retrospective research, no MLAP was found during pregnancy¹⁵. Grassi et al. found the incidence of MLAP as 1.3% in a cohort of 80 patients from Italy¹⁶. In two different studies (n = 418 and n = 134), the rate of MLAP was reported as 18.2% and 54.8%¹⁶⁻¹⁸. Fang et al. did not detect any association between

MLAP and gender, age, cancer history, intensive care unit (ICU) admission, length of hospital stays, and laboratory parameters, but with cobblestone imaging findings¹⁹. In a French study, Valette et al. published that MLAP existed in six of nine individuals with severe COVID-19 in the ICU²⁰. Studies from Chinese patients indicated the presence of MLAP at 43.5%, 41.7%, and 19.8%^{18,19,21}.

Kaya and Akman investigated the relationship between MLAP and ICU hospitalization and mortality in COVID-19 patients and found a statistical significance between the three parameters⁶. They claimed that the more severe findings on CT scans predict the prognosis. In addition, they stated that increased bronchial wall

Table 4. Laboratory parameters affecting $SUV_{max} \geq 2.5$

Characteristics	Univariate		Multivariate	
	Odds ratio (95% CI)	p	Odds ratio (95% CI)	p
Age	1.013 (0.990-1.037)	0.271		
Gender				
Male	Reference	-		
Female	1.690 (0.771-3.702)	0.190		
RT-PCR				
Negative	Reference	-	Reference	-
Positive	15.000 (5.856-38.058)	< 0.001	5.964 (1.937-18.370)	0.002
Peripheral LAP				
No	Reference	-	Reference	-
Yes	3.890 (1.801-8.398)	< 0.001	1.451 (0.491-4.289)	0.500
Chest X-Ray				
Normal	Reference	-	Reference	-
Abnormal	9.280 (3.355-25.662)	< 0.001	2.777 (0.758-10.165)	0.123
Pathology				
Normal	Reference	-	Reference	-
Abnormal	33.857 (10.833-105.818)	< 0.001	13.452 (3.864-46.831)	< 0.001

RT-PCR: reverse transcription-polymerase chain reaction; LAP: lymphadenopathy.

thickness was more common in patients with MLAP. The mortality rate was higher in patients with MLAP¹.

In another study, Satici et al.²² reported that mediastinal lymphadenopathies were detected more in elderly patients with comorbid diseases and were significantly associated with mortality. The rate of MLAP was 9.2% in the whole study population but 19.65% in deceased individuals²¹. Both studies declared that the presence of MLAP led to increased mortality. Similar to our findings, mediastinal lymph node involvement has been an essential factor in COVID-19.

In the study of Bhatti et al.,²³ MLAP was detected in 131 (62.4%) of 210 patients included in the study. Covid-19 patients with MLAP had a higher mean and median severity score than those without MLAP. This study documents the high prevalence of MLAP in hospitalized COVID-19 patients and shows that the severity score, which represents the more severe course of the disease, is higher²².

Limitations of our study

Since our study was retrospective, some laboratory findings and respiratory function findings could not be

obtained. In this regard, a study can be organized to investigate the laboratory findings showing the severity of chronic inflammation in patients with severe COVID-19 pneumonia, respiratory functional characteristics, and the relationship between radiological imaging and MLAP features. Although this study is a retrospective observational study and provides valuable information about the potential etiology of the disease, it typically cannot definitively determine causality.

Conclusions

This study showed that severe COVID-19 and mediastinal lymph node involvement (SUV_{max} : 3-5) were associated regardless of age, gender, comorbidities, and disease outcomes. Establishing reliable SUVMAX values in severe post-COVID-19 patients will help guide clinical decisions, adapt therapeutic approaches, and avoid unnecessary surgical interventions for patients with MLAP who require surgical diagnosis and treatment, and will help these patients avoid surgery complications.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of humans and animals. The authors declare that no experiments involving humans or animals were conducted for this research.

Confidentiality, informed consent, and ethical approval. The authors have obtained approval from the Ethics Committee for the analysis of routinely obtained and anonymized clinical data, so informed consent was not necessary. Relevant guidelines were followed. The study was carried out with the permission of Health Sciences University Şişli Hamidiye Etfal Training and Research Hospital Ethics Committee (Date: September 06, 2022, Decision No:2137).

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing of this manuscript.

References

1. Zhang HP, Sun YL, Wang YF, Yazici D, Azkur D, Ogulur I, et al. Recent developments in the immunopathology of COVID-19. *Allergy*. 2023; 78:369-88.
2. Filchakova O, Dossym D, Ilyas A, Kuanysheva T, Abdizhamil A, Bukasov R. Review of COVID-19 testing and diagnostic methods. *Talanta*. 2022;244:123409.
3. Jeong YJ, Wi YM, Park H, Lee JE, Kim SH, Lee KS. Current and emerging knowledge in COVID-19. *Radiology*. 2023;306:e222462.
4. Valentin B, Steuwe A, Wienemann T, Andree M, Keitel V, Ljimini A, et al. CT findings in patients with COVID-19-compatible symptoms but initially negative qPCR Test. *Rofo*. 2022;194:1110-18.
5. Munden RF, Carter BW, Chiles C, MacMahon H, Black WC, Ko JP, et al. Managing incidental findings on thoracic CT: mediastinal and cardiovascular findings. A white paper of the ACR incidental findings committee. *J Am Coll Radiol*. 2018;15:1087-96.
6. Kaya AT, Akman B. Mediastinal lymph node enlargement in COVID-19: relationships with mortality and CT findings. *Heart Lung*. 2022;54:19-26.
7. T.R. Ministry of Health, General Directorate of Public Health. COVID-19 (SARS-CoV-2 Infection) Guide, Scientific Board Study. Ankara: T.R. Ministry of Health; 2020.
8. Lee JH, Koh J, Jeon YK, Goo JM, Yoon SH. An integrated radiologic-pathologic understanding of COVID-19 pneumonia. *Radiology*. 2023;306:e222600.
9. Abbasi B, Pezeshki-Rad M, Soleimani H, Mozdourian M, Akhavan R, Maftouh M. Temporal changes of lung computed tomography findings pulmonary COVID-19 infection. *J Med Imaging Radiat Sci*. 2022;53:564-70.
10. Landini N, Colzani G, Ciet P, Tessarin G, Dorigo A, Bertana L, et al. Chest radiography findings of COVID-19 pneumonia: a specific pattern for a confident differential diagnosis. *Acta Radiol*. 2022;63:1619-26.
11. Ravaglia C, Doglioni C, Chilosi M, Piciocchi S, Dubini A, Rossi G, et al. Clinical, radiological and pathological findings in patients with persistent lung disease following SARS-CoV-2 infection. *Eur Respir J*. 2022;60:2102411.
12. Taweeseedt PT, Surani S. Mediastinal lymphadenopathy in COVID-19: a review of literature. *World J Clin Cases*. 2021;9:2703-10.
13. Peng X, Guo Y, Xiao H, Xia W, Zhai A, Zhu B, et al. Overview of chest involvement at computed tomography in children with coronavirus disease 2019 (COVID-19). *Pediatr Radiol*. 2021;51:222-30.
14. Bayramoglu Z, Canipek E, Comert RG, Gasimli N, Kaba O, Sari Yanartaş M, et al. Imaging features of pediatric COVID-19 on chest radiography and chest CT: a retrospective, single-center study. *Acad Radiol*. 2021;28:18-27.
15. Gong X, Song L, Li H, Li L, Jin W, Yu K, et al. CT characteristics and diagnostic value of COVID-19 in pregnancy. *PLoS One*. 2020;15:e0235134.
16. Grassi R, Fusco R, Belfiore MP, Montanelli A, Patelli G, Urraro F, et al. Coronavirus disease 2019 (COVID-19) in Italy: features on chest computed tomography using a structured report system. *Sci Rep*. 2020;10:17236.
17. Sardanelli F, Cozzi A, Monfardini L, Bnà C, Foà RA, Spinazzola A, et al. Association of mediastinal lymphadenopathy with COVID-19 prognosis. *Lancet Infect Dis*. 2020;20:1230-1.
18. Fang X, Li X, Bian Y, Ji X, Lu J. Relationship between clinical types and radiological subgroups defined by latent class analysis in 2019 novel coronavirus pneumonia caused by SARS-CoV-2. *Eur Radiol*. 2020;30:6139-50.
19. Fang X, Li X, Bian Y, Ji X, Lu J. Radiomics nomogram for the prediction of 2019 novel coronavirus pneumonia caused by SARS-CoV-2. *Eur Radiol*. 2020;30:6888-901.
20. Valette X, du Cheyron D, Goursaud S. Mediastinal lymphadenopathy in patients with severe COVID-19. *Lancet Infect Dis*. 2020;20:1230.
21. Li X, Fang X, Bian Y, Lu J. Comparison of chest CT findings between COVID-19 pneumonia and other types of viral pneumonia: a two-center retrospective study. *Eur Radiol*. 2020;30:5470-8.
22. Satici C, Cengel F, Gurkan O, Demirkol MA, Altunok ES, Esatoglu SN. Mediastinal lymphadenopathy may predict 30-day mortality in patients with COVID-19. *Clin Imaging*. 2021;75:119-24.
23. Bhatti FS, Malik AA, Malik AA. Presence of mediastinal lymphadenopathy in hospitalized Covid-19 patients in a tertiary care hospital in Pakistan-A cross-sectional study. *PLoS One*. 2023 May 25;18(5).

An evaluation of factors which affect recurrence and malignancy of sinonasal papillomas

Evaluación de los factores que afectan la recurrencia y la malignidad de los papilomas sinasales

Zülküf Küçüktağ^{1*}, Bülent Öcal¹, Cem Saka¹, Esmâ Altan¹, Mehmet H. Korkmaz^{1,2}, and Emel Çadallı-Tatar¹

¹Department of Otolaryngology Head and Neck Surgery, University of Health Sciences Dışkapı Yıldırım Beyazıt Training and Research Hospital;

²Department of Otolaryngology Head and Neck Surgery, Ankara Yıldırım Beyazıt University, Faculty of Medicine. Ankara, Turkey

Abstract

Objective: Sinonasal papillomas are rare benign tumors. They challenge otorhinolaryngologists due to their high recurrence rate and malignant transformation. The aim of this study was to investigate the clinical and demographic factors associated with recurrence and/or malignancy of sinonasal papillomas. **Methods:** This retrospective study included 49 patients who were operated on between 2014 and 2020. Clinical and demographic characteristics of the patients were retrieved from the patients' records. **Results:** The recurrence rate was determined to be 12.2% and the malignancy rate was 8.2%. Recurrence was significantly higher in females, the advanced stage (T3 and T4), and papillomas located in the maxillary sinus. The recurrence rate was significantly lower in patients treated with the endoscopic surgery. Factors affecting malignancy were the presence of dysplasia and extension out of the paranasal sinuses. **Conclusions:** The results of this study showed that the factors affecting recurrence were gender, papilloma stage, type of surgery performed, and whether the tumor extended to the paranasal sinuses.

Keywords: Sinonasal papilloma. Inverted papilloma. Paranasal sinus.

Resumen

Objetivo: Los papilomas sinasales son tumores benignos poco frecuentes. Suponen un reto para los otorrinolaringólogos por su elevada tasa de recurrencia y transformación maligna. Investigar los factores clínicos y demográficos asociados con la recurrencia y con la malignidad de los papilomas sinasales. **Métodos:** Estudio retrospectivo que incluyó 49 pacientes operados entre 2014 y 2020. Las características clínicas y demográficas se recuperaron de los registros de los pacientes. **Resultados:** Se determinó una tasa de recurrencia del 12.2% y una tasa de malignidad del 8.2%. La recurrencia fue significativamente mayor en las mujeres, en los estadios avanzados (T3 y T4) y en los papilomas localizados en el seno maxilar. La tasa de recidiva fue significativamente menor en los pacientes tratados con cirugía endoscópica. Los factores que afectaron a la malignidad fueron la presencia de displasia y la extensión fuera de los senos paranasales. **Conclusiones:** Los resultados de este estudio muestran que los factores que afectaron la recidiva fueron el sexo, el estadio del papiloma, el tipo de cirugía realizada y si el tumor se extendía a los senos paranasales.

Palabras clave: Papiloma sinasal. Papiloma invertido. Seno paranasal.

*Correspondence:

Zülküf Küçüktağ

E-mail: zkucuk79@yahoo.com

2444-0507/© 2024 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 05-11-2023

Date of acceptance: 30-01-2024

DOI: 10.24875/CIRUE.M24000902

Cir Cir (Eng). 2025;93(6):636-643

Contents available at PubMed

www.cirugiaycirujanos.com

Introduction

Sinonasal papillomas are benign tumors originating from the Schneiderian membrane of the nasal cavity and paranasal sinuses. There are three types of sinonasal papillomas; exophytic, oncocytic, and inverted, with the latter being the most common type. The most important features that distinguish inverted papilloma (IP) from other benign sinonasal tumors are; high recurrence, malignancy potential, and significant local bone erosion¹. The recurrence rate has been reported to be between 15% and 20%²⁻⁷. Recurrences usually occur within the first 3 years after incomplete excisions. IP may be histopathologically associated with varying degrees of dysplasia, carcinoma *in situ*, and squamous cell carcinoma (SCC)⁸. The malignancy rate has been reported to be approximately 1.9-27%⁹⁻¹³. Malignancy may coexist with the IP in the same lesion or may develop years after excision of the papilloma.

Many factors, including smoking, size of the papilloma, type of surgery, the presence of squamous hyperplasia, high hyperkeratosis, increased mitosis, HPV positivity, bilateral papilloma and localization of IP have been associated with recurrence or malignancy in the literature^{9,10,14}. In the majority of case series, the limited number of patients makes it difficult to draw reliable conclusions from these data, as sinonasal papillomas are rare and it can take many years to gather a large number of cases. Each institution should share their experiences to increase the number of the patients to be able to make an additional contribution to the literature on this subject. The aim of this study was to investigate the clinical and demographic factors associated with recurrence and/or malignancy in sinonasal papillomas. Furthermore, with the results obtained from this study, it was aimed to contribute to the pool of knowledge on this subject, which has many controversies and uncertainties.

Methods

The study was initiated following the approval of the University of Health Sciences, Dışkapı Yıldırım Beyazıt Training and Research Hospital Clinical Research Ethics Committee (Date: September 21, 2020, Number: 96/6). The study cohort included 61 patients who underwent surgery for sinonasal papilloma between 2014 and 2020. After the exclusion of 12 patients who were first operated on another center or who did not

have complete follow-up data, the analyses were completed with 49 patients. The parameters evaluated in this study were patient age, gender, smoking and alcohol consumption, follow-up time, type of surgery, papilloma side and localization of origin, the presence extranasal extension, the type and stage of the papilloma, and recurrence and malignancy rates. The Krouse Staging System¹⁵ was used for staging. The stage, localisation, origin, and invasion site of papillomas were determined according to the intraoperative findings and paranasal sinus tomography images.

The post-operative regular follow-up examinations included nasal endoscopic examination every 3 months in the 1st year, every 6 months in the 2nd year, and annually thereafter for up to 7 years. In the follow-up examinations, all cavities and sinuses were checked with straight and angled telescopes. In cases with suspicious lesions in the frontal sinus or in the anterior wall of the maxillary sinus, imaging was performed to view the areas that could not be seen with endoscopy. Patients with suspicious lesions identified in endoscopic examination underwent biopsies.

The Chi-square test was used in the statistical analysis. Parameters with a $p < 0.05$ were considered statistically significant.

Results

Demographic and clinical parameters of patients

The patients comprised 43 (87.8%) males and 6 (12.2%) females with a mean age of 48.96 ± 15.9 years (Range, 15-78 years) and a mean follow-up period of 39.16 ± 20.4 months (Range, 8-88 months). The demographic and clinical parameters of the patients are summarized in table 1.

Factors associated with recurrence

The overall recurrence rate of this cohort was 12.2%. The earliest time to recurrence was 9 months, and the latest was 72 months (Mean, 39.5 ± 21.6 months). All recurrences occurred in the same localisation as the original tumor. The relationships between the demographic and clinical parameters and recurrence are shown in table 2. Recurrence was observed in 7% of male and 50% of female patients. The difference between the gender in terms of recurrence rates was statistically significant ($p = 0.003$).

Table 1. Demographic and clinical parameters of patients

Parameters	Number of patients (%)
Age	
Mean ± standard deviation (SD)	48.96 ± 15.9
≤ 45	16 (32.7)
> 45	33 (67.3)
Gender	
Male	43 (87.8)
Female	6 (12.2)
Follow-up time (months)	
Min-Max	8-88
Mean ± SD	39.16 ± 20.4
Side of papilloma	
Right	34 (69.4)
Left	15 (30.6)
Type of papilloma	
Inverted	44 (89.8)
Non-inverted	5 (10.2)
Exophytic	5 (10.2)
Oncocytic	0
Originated site	
Paranasal sinus	27 (55.1)
Maxillary sinus	23 (46.9)
Frontal sinus	2 (4.1)
Sphenoid sinus	2 (4.1)
Ethmoid sinus	0
Nasal cavity	22 (44.9)
Lateral nasal wall	12 (24.5)
Middle turbinate	9 (18.4)
Inferior turbinate	1 (2)
Stage of papilloma	
Early (T1-T2)	18 (36.7)
T1	8 (16.3)
T2	10 (20.4)
Advanced (T3-T4)	31 (63.3)
T3	26 (53.1)
T4	5 (10.2)
Malign transformation	
Yes	4 (8.2)
No	45 (91.8)
Displasia	
Yes	5 (10.2)
No	44 (89.8)
Smoking	
Yes	20 (40.8)
No	29 (59.2)
Alcoholism	
Yes	3 (6.1)
No	46 (93.8)
Type of surgery	
Endoscopic	42 (85.7)
Open	0
Combine	7 (14.3)

Endoscopic surgery was performed on 42 (85.7%) patients, and 7 patients (14.3%) underwent combined surgery (Maxillary trocar, Caldwell Luc, and Bicoronal Forehead flap in 4, 2, and 1 patients, respectively). The recurrence rate in patients who underwent only endoscopic surgery was found to be significantly lower than those who underwent combined surgery ($p = 0.008$).

The sinonasal papillomas were separated into two groups according to localisation in the paranasal sinus or nasal cavity. The distribution of the sinonasal papillomas according to localisations is summarized in table 3. Sinonasal papillomas most commonly originated from the maxillary sinus in this cohort. Recurrence occurred in 6 of 27 patients (22.2%) with papilloma located in the paranasal sinus, and no recurrence was observed in any of the papillomas located in the nasal cavity. The difference between the two groups was statistically significant ($p = 0.018$). All the recurrences originated from the maxillary sinus.

In this study, the patients were evaluated in two groups. Those with Krouse stage T1 and T2, in which the papilloma is limited to the nasal cavity, ethmoid sinuses, and medial wall of maxillary sinus, were defined as early stage (Group 1). Patients with papilloma, which are more difficult to reach surgically, such as in the other walls of the maxillary sinus, sphenoid, and/or frontal sinuses (Krouse T3 and T4), were accepted as advanced stage (Group 2). Patients with malignancy and/or with extension out of the nasal cavity and paranasal sinus were also included in Group 2. The distribution of papilloma types according to the Krouse Staging System is shown in table 4. More than half of the patients were at the advanced stage (53.1% T3 and 10.2 % T4). All patients with recurrence were at T3 stage. There was a statistically significant difference between the two groups in terms of recurrence ($p = 0.046$). In five patients, the papilloma extended out of the nasal cavity and paranasal sinus. IP extended to the orbit in two of these patients, to the orbit and intracranial region in one patient, and to the parapharyngeal region in one patient. Another patient had an oroantral fistula stemming from IP invasion.

The study results showed that patient age, smoking and alcohol consumption, papilloma side and type, and the extension of sinonasal papilloma out of the nasal cavity and paranasal sinuses had no significant effect on recurrence ($p > 0.05$).

Table 2. The effects of demographic and clinical parameters of patients on recurrence

Parameters	Patients with Recurrence (number of patients [%])	Patients without recurrence (number of patients [%])	Total (number of patients [%])	p
Age Min-Max (mean±standard deviation)	16-59 (44.83 ± 15.2)	15-78 (49.53 ± 16)	15-78 (48.96 ± 15.9)	
≤ 45	2 (12.5)	14 (87.5)	16 (32.7)	0.97
> 45	4 (12.1)	29 (87.9)	33 (67.3)	
Side of papilloma				
Right	5 (14.7)	29 (85.3)	34 (69.4)	0.43
Left	1 (6.7)	14 (93.3)	15 (30.6)	
Gender				
Female	3 (50)	3 (50)	6 (12.2)	0.003*
Male	3 (7)	40 (93)	43 (87.8)	
Type of papilloma				
Inverted	6 (13.6)	38 (86.4)	44 (89.8)	0.38
Non-inverted	-	5 (100)	5 (10.2)	
Papilloma's originated site				
Intra-paranasal sinus	6 (22.2)	21 (77.8)	27 (55.1)	0.018*
Nasal cavity	-	22 (100)	22 (44.9)	
Extension out of the paranasal sinus				
Yes	-	5 (100)	5 (10.2)	0.38
No	6 (13.6)	38 (86.4)	44 (89.8)	
Stage of papilloma				
Early (T1-T2)	-	18 (100)	18 (36.7)	0.046*
Advanced (T3-T4)	6 (19.3)	25 (80.7)	31 (63.3)	
Smoking				
Yes	3 (15)	17 (85)	20 (40.8)	0.63
No	3 (10.3)	26 (89.7)	29 (59.2)	
Alcohol use				
Yes	-	3 (100)	3 (6.1)	0.71
No	6 (13)	40 (87)	46 (93.9)	
Type of surgery				
Endoscopic surgery	3 (7.1)	39 (92.9)	42 (85.7)	0.008*
Combined surgeries	3 (42.9)	4 (57.1)	7 (14.3)	

*p < 0.05, statistically significant values.

Table 3. Distribution of sinonasal papillomas according to originated site

Originated site	Number of patients (%)
Paranasal sinus	27 (55.1)
Maxillary sinus	23 (46.9)
Ethmoid sinus	-
Frontal sinus	2 (4.1)
Sphenoid sinus	2 (4.1)
Nasal cavity	22 (44.9)
Lateral nasal wall	12 (24.5)
Middle turbinate	9 (18.4)
Inferior turbinate	1 (2)

Factors associated with malignancy

The malignancy rate of this study was found to be 8.2% (4/49). All the patients who developed malignancy had inverted type and synchronous SCC. All patients with malignancy were male with a mean age of 54.75 years. All four patients with malignancy were treated with endoscopic surgery, and the follow-up time of these patients was 8, 22, 23, and 37 months, respectively. Recurrence was observed in only one patient with malignancy.

Of the 5 patients (40%) with dysplasia, malignancy was determined in 2 (40%). The malignancy rate of IP with dysplasia was significantly higher than that of IP without dysplasia (p = 0.006). The rate of malignancy

in IP extending out of the nasal cavity and paranasal sinuses was statistically significantly higher than in those without extension (3 of 5 patients, $p < 0.001$). In this cohort, no other parameters had any effect on malignancy.

Discussion

The overall recurrence rate in this study was 12.2% (6 of 49 patients), which was consistent with the literature. Some studies have reported significantly higher rates of recurrence¹⁶⁻¹⁸. The high rate of recurrence in one of these studies was explained by the fact that their hospital was a tertiary health-care center which treated more complex patients¹⁸. However, the development of recurrence in sinonasal papillomas is a multifactorial event and should not be attributed to a single factor. Factors such as the type of surgery applied, the surgeon's experience, the localization and stage of the papilloma, and the presence of residue may also affect recurrence. Many studies have been conducted on the recurrence and malignancy of sinonasal papillomas, but no consensus has been reached. In the current study, the factors with significant effect on the recurrence of sinonasal papillomas were determined to be; female gender, the stage and localization of the papilloma, and the type of surgery applied. Recurrence rates were found to be increased by a younger patient age and smoking according to Jardine et al.¹⁹, whereas Roh et al.²⁰ and Moon et al.²¹ reported that smoking, Suh et al.²² stated that female gender, and Katori et al.²³ that some histopathological features of papilloma (increased mitotic activity, hyperkeratosis, and the presence of squamous hyperplasia) increased recurrence. In other studies, residual papilloma tissue following surgery has also been blamed for recurrence^{10,24}.

In many studies, the recurrence rate in males has been reported to be higher than in females^{18,24}. In contrast, the recurrence rate in the current study was significantly higher in females than in males. Suh et al.²² also reported the relationship between recurrence and gender, with a recurrence rate in female patients of 50% (4 of 8 patients), similar to the current study. However, to be able to more clearly state that female gender increases recurrence, there is a need for further studies to be conducted with more female patients.

It has been reported that recurrence is more common in IP located in the frontal sinus, which is due to the difficulty in visualization of the sinus^{6,18,25-27}.

Recurrence in IP with frontal sinus localization has been reported to be 22.4%²⁸. In the two patients with frontal sinus localization in the current study, no recurrence developed. In some studies, it has been reported that maxillary sinus localisation increases recurrence^{7,29} and the results obtained in the current study were similar to those studies. In all the patients with recurrence, the IP was located in the maxillary sinus, and there was invasion of the anterior and lateral walls of the maxillary sinus. Just like the lateral wall of the frontal sinus, the anterior wall of the maxillary sinus is an area where papilloma can be overlooked, and residual tissue may remain. Although Caldwell-Luc and/or gingival trocar surgery was added to the endoscopic surgery in these cases; unfortunately, recurrence was still observed in some patients. In these cases, the inside of the sinus should be carefully checked with straight and angled telescopes to ensure that no residual tissue remains. The mucosa invaded by the papilloma should be curetted, and the bone tissue should be removed by drilling. When there is frontal sinus lateral wall invasion, it is difficult to reach this area with endoscopic surgery alone, so adding open surgeries such as bicoronal forehead flap and/or osteoplastic flap to the endoscopic surgery allows the papilloma to be removed without residue.

There are different opinions about the relationship between the stage of the IP and recurrence. Some studies have highlighted an association between recurrence and Krouse classification³⁰, while others have not^{4,5,7,17,24-26}. A meta-analysis revealed an increased risk of recurrence between Krouse T2 and T3 stages,³¹ and another study showed a significant relationship only between Krouse T4 stage and recurrence²¹. In most studies, the four stages have been compared with each other in terms of recurrence. In the current study, T1 and T2 stage papillomas, which are easier to reach surgically, were defined as early stage, and T3 and T4 stage papillomas, which are more difficult to reach, as advanced stage. Thus, it was aimed to compare 2 more homogeneous groups. The results showed that advanced-stage papillomas significantly increased recurrence.

In this cohort, the malignancy rate was observed to be 8.2% (4/49), which was consistent with the literature. All four patients had synchronous SCC, and no metachronous malignancy was observed during follow-up. Metachronous malignancy may develop many years after excision of the IP. Lesperance et al.³² described an average interval of 63 months between the onset of IP and the development of malignancy.



Figure 1. In the coronal paranasal sinus tomography, inverted papilloma + squamous cell carcinoma extending to the orbit by eroding the frontal sinus wall on the right side (the arrow shows erosion in frontal sinus wall).




Figure 2. In the coronal paranasal sinus magnetic resonance T1 image, an inverted papilloma + squamous cell carcinoma that fills both nasal passages and all sinuses and extends to the orbit and intracranial area.

Another study showed that the development of metachronous malignancy occurred approximately 10 years after undergoing IP surgery³³. These data reveal the importance of long-term follow-up of patients with IP. Therefore, these patients should be checked at regular intervals, their complaints should be questioned, and endoscopic nasal examinations should be performed. If a suspicious mass is detected, radiological imaging should be performed and a biopsy should be taken.

Malignancy of IPs may present with different clinical scenarios. They may present with limited eye movements without a mass in the nasal passage and complaints of nasal congestion, as in one of the current study patients (Fig. 1). Or, as in the other patient, they may present with visual loss and loss of consciousness due to bilateral nasal mass that has invaded the orbit and intracranial area (Fig. 2). Clinicians must be alert to such cases.

Many factors affecting the malignancy of IPs have been reported. Hong et al. reported that smoking significantly affects malignancy³⁴. In a literature review, some biochemical and immunohistochemical markers

were found to play a role in malignant transformation¹³. In this cohort, the presence of dysplasia and invasion of papilloma outside the nasal cavity and paranasal sinus had a significant effect on malignancy in IPs. The rate of malignancy in IPs extending out of the paranasal sinus was significantly higher than those that did not. This result can be explained in 2 ways. First, papillomas that tend to extension outside the nasal cavity and the paranasal sinus are perhaps more aggressive and have higher malignant potential. The second reason may be the malignant and aggressive characteristics of the malignant tumor accompanying the IP. More research is needed of IPs that tend to invade to be able to predict which papillomas may behave more aggressively or become malignant.

For treatment of IPs, more experienced surgeons prefer the classical open surgery, whereas younger surgeons tend to perform endoscopic surgery²². In 2019, a meta-analysis stated that open surgeries are no longer the “gold standard” for surgery of IPs⁶. Goudakos et al.³⁵ stated that endoscopic surgery is the first choice for IPs, regardless of the stage of

Table 4. Distribution of papilloma types according to krouse staging system

Stage	Inverted (number of patients [%])	Exophytic (number of patients [%])	Oncocytic (number of patients [%])	Total (number of patients [%])
Early stage T1	8 (18.2%)	-	-	8 (16.3%)
T2	9 (20.4%)	1 (20%)	-	10 (20.4%)
Advanced stage T3	22 (50%)	4 (80%)	-	26 (53.1%)
T4	5 (11.4%)	-	-	5 (10.2%)
Total	44 (100%)	5 (100%)	0	49

papilloma. Endoscopic surgery provides better surgical results than open and combined surgeries³. In a meta-analysis by Peng et al., it was stated that the recurrence rate of the endoscopic surgery group was significantly lower compared to the open surgery group⁶. Bugter et al.¹⁷ also confirmed the superiority of endoscopic surgery over open and combine surgery. In the current cohort, 85.71% of the patients were treated with the endoscopic approach and 14.29% with combined surgery. Open surgery alone was not performed in any patient. It can be considered that every surgeon experienced in endoscopic sinus surgery can treat most patients with sinonasal papilloma, even malignant cases, with an endoscopic approach. Open surgeries can be added to the endoscopic approach, especially in cases that are difficult to reach endoscopically, such as localisation in the lateral wall of the frontal sinus, or anterior and lateral wall of the maxillary sinus. In appropriately selected patients, endoscopic surgery allows removal of the tumor without residue. In summary, endoscopic surgery; facilitates the treatment of sinonasal papillomas, decreases the complication and recurrence rates, shortens the hospitalization time, and accelerates post-operative recovery.

The present study had some limitations primarily that the retrospective nature of the study had a negative impact on the quantity and quality of the information, and this should be taken into account when interpreting the results. The number of patients was not very high. Because sinonasal papilloma is a rare disease and it takes a long time to gather a large case series. Further limitations could be said to be that the study was conducted in a single center and patients who had previously undergone surgery in another center were excluded. Therefore, there is a need for further studies with more patients to support the values that were found to be significant.

Conclusions

Although many factors affecting the recurrence of sinonasal papillomas have been identified, there is no definite consensus on this issue. In this study, the factors with a significant effect on recurrence were determined to be female gender, localization and stage of papilloma, and type of surgery. Factors affecting malignancy were the presence of dysplasia and extension out of the paranasal sinuses. Most sinonasal papillomas, including malignant cases, can be treated with endoscopic surgery regardless of stage. When necessary, open surgical approaches can be added to the endoscopic approach. Nevertheless, there is a need for more extensive research, especially molecular research, to further elucidate this complex issue.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of humans and animals. The authors declare that no experiments involving humans or animals were conducted for this research.

Confidentiality, informed consent, and ethical approval. The authors have followed their institution's confidentiality protocols. The authors have obtained approval from the Ethics Committee for the analysis of routinely obtained and anonymized clinical data, so informed consent was not necessary.

Declaration on the use of artificial intelligence.

The authors declare that no generative artificial intelligence was used in the writing of this manuscript.

References

- Wang MJ, Noel JE. Etiology of sinonasal inverted papilloma: a narrative review. *World J Otorhinolaryngol Head Neck Surg.* 2017;3:54-8.
- Peng P, Har-El G. Management of inverted papillomas of the nose and paranasal sinuses. *Am J Otolaryngol.* 2006;27:233-7.
- Busquets JM, Hwang PH. Endoscopic resection of sinonasal inverted papilloma: a meta-analysis. *Otolaryngol Head Neck Surg.* 2006;134:476-82.
- Marino-Sanchez F, De Los Santos G, Giribet A, Aguirre A, Alonso J, Wansley DL, et al. Incidental histopathologic finding of sinonasal inverted papilloma among surgically excised polyps increases the risk of tumor recurrence. *Indian J Otolaryngol Head Neck Surg.* 2019;71 Suppl 3:1910-7.
- Jiang XD, Dong QZ, Li SL, Huang TQ, Zhang NK. Endoscopic surgery of a sinonasal inverted papilloma: surgical strategy, follow-up, and recurrence rate. *Am J Rhinol Allergy.* 2017;31:51-5.
- Peng R, Thamboo A, Choby G, Ma Y, Zhou B, Hwang PH. Outcomes of sinonasal inverted papilloma resection by surgical approach: an updated systematic review and meta-analysis. *Int Forum Allergy Rhinol.* 2019;9:573-81.
- Healy DY Jr., Chahabra N, Metson R, Holbrook EH, Gray ST. Surgical risk factors for recurrence of inverted papilloma. *Laryngoscope.* 2016;126:796-801.
- Li W, Lu H, Zhang H, Sun X, Hu L, Wang D. Squamous cell carcinoma associated with inverted papilloma: recurrence and prognostic factors. *Oncol Lett.* 2020;19:1082-8.
- Vorasubin N, Vira D, Suh JD, Bhuta S, Wang MB. Schneiderian papillomas: comparative review of exophytic, oncocytic, and inverted types. *Am J Rhinol Allergy.* 2013;27:287-2.
- Xiao-Ting W, Peng L, Xiu-Quing W, Hai-Bo W, Wen-Hui P, Bing L, et al. Factors affecting recurrence of sinonasal inverted papilloma. *Eur Arch Otorhinolaryngol.* 2013;270:1349-53.
- Yang YJ, Abraham JL. Undifferentiated carcinoma arising in oncocytic Schneiderian (cylindrical cell) papilloma. *J Oral Maxillofac Surg.* 1997;55:289-94.
- Nudell J, Chiosea S, Thompson LD. Carcinoma ex-Schneiderian papilloma (malignant transformation): a clinicopathologic and immunophenotypic study of 20 cases combined with a comprehensive review of the literature. *Head Neck Pathol.* 2014;8:269-86.
- Re M, Gioacchini FM, Bajraktari A, Tomasetti M, Kaleci S, Rubini C, et al. Malignant transformation of sinonasal inverted papilloma and related genetic alterations: a systematic review. *Eur Arch Otorhinolaryngol.* 2017;274:2991-3000.
- Govindaraj S, Wang H. Does human papilloma virus play a role in sinonasal inverted papilloma? *Curr Opin Otolaryngol Head Neck Surg.* 2014;22:47-51.
- Krouse JH. Development of a staging system for inverted papilloma. *Laryngoscope.* 2000;110:965-8.
- Sbrana MF, Borges RF, Pinna FR, Neto DB, Voegels RL. Sinonasal inverted papilloma: rate of recurrence and malignant transformation in 44 operated patients. *Braz J Otorhinolaryngol.* 2021;87:80-4.
- Bugter O, Monserez DA, Van Zijl FV, De Jong JB, Hardillo JA. Surgical management of inverted papilloma; a single-center analysis of 247 patients with long follow-up. *J Otolaryngol Head Neck Surg.* 2017;46:67.
- Lee JJ, Roland LT, Licata JJ, Orlowski HL, Jiramongkolchai P, Piccirillo JF, et al. Morphologic, intraoperative, and histologic risk factors for sinonasal inverted papilloma recurrence. *Laryngoscope.* 2020;130:590-6.
- Jardine AH, Davies GR, Birchall MA. Recurrence and malignant degeneration of 89 cases of inverted papilloma diagnosed in a non-tertiary referral population between 1975 and 1995: clinical predictors and p53 studies. *Clin Otolaryngol Allied Sci.* 2000;25:363-9.
- Roh HJ, Mun SJ, Cho KS, Hong SL. Smoking, not human papilloma virus infection, is a risk factor for recurrence of sinonasal inverted papilloma. *Am J Rhinol Allergy.* 2016;30:79-82.
- Moon IJ, Lee DY, Suh MW, Han DH, Kim ST, Min YG, et al. Cigarette smoking increases risk of recurrence for sinonasal inverted papilloma. *Am J Rhinol Allergy.* 2010;24:325-9.
- Suh KW, Facer GW, Devine KD, Weiland LH, Zujko RD. Inverting papilloma of the nose and paranasal sinuses. *Laryngoscope.* 1997;87:35-46.
- Katori H, Nozawa A, Tsukuda M. Histopathological parameters of recurrence and malignant transformation in sinonasal inverted papilloma. *Acta Otolaryngol.* 2006;126:214-8.
- Lisan Q, Laccourreye O, Bonfils P. Sinonasal inverted papilloma: risk factors for local recurrence after surgical resection. *Ann Otol Rhinol Laryngol.* 2017;126:498-504.
- Kim DY, Hong SL, Lee CH, Jin HR, Kang JM, Lee BJ, et al. Inverted papilloma of the nasal cavity and paranasal sinuses: a Korean multicenter study. *Laryngoscope.* 2012;122:487-94.
- Lilja M, Viitasalo S, Hytönen M, Haapaniemi A, Hagström J, Mäkitie A. Sinonasal oncocytic papilloma-A series of 20 cases with special emphasis on recurrences. *Laryngoscope Investig Otolaryngol.* 2019;4:567-72.
- Gamrot-Wrzol M, Sowa P, Lisowska G, Scierski W, Misiulek M. Risk factors of recurrence and malignant transformation of sinonasal inverted papilloma. *Biomed Res Int.* 2017;2017:9195163.
- Walgama E, Ahn C, Batra PS. Surgical management of frontal sinus inverted papilloma: a systematic review. *Laryngoscope.* 2012;122:1205-9.
- Minni A, Gera R, Bulgheroni C, Ralli M, Cialente F, Candelori F, et al. Endoscopic resection of sinonasal inverted papilloma: a multivariate retrospective analysis of factors affecting recurrence and persistence. *Ear Nose Throat J.* 2021;100 Suppl 5:542S-8.
- Satter NB, Cannady SB, Citardi MJ, Roh HJ, Batra PS. Comparison of open versus endoscopic resection of inverted papilloma. *Am J Rhinol.* 2007;21:320-3.
- Lisan Q, Moya-Plana A, Bonfils P. Association of krouse classification for sinonasal inverted papilloma with recurrence. *JAMA Otolaryngol Head Neck Surg.* 2017;143:1104-10.
- Lesperance MM, Esclamado RM. Squamous cell carcinoma arising in inverted papilloma. *Laryngoscope.* 1995;105:178-83.
- Choi CW, Kim SG, Kim YM, Yoon YH, Kim AY, Rha KS. Clinical and histologic features of inverted papilloma-associated malignancy. *Eur Arch Otorhinolaryngol.* 2012;269:2349-54.
- Hong SY, Kim BH, Lee JH, Cho KS, Roh HJ. Smoking and malignancy in sinonasal inverted papilloma. *Laryngoscope.* 2013;123:1087-91.
- Goudakos JK, Blioskas S, Nikolaou A, Vlachtsis K, Karkos P, Markou KD. Endoscopic resection of sinonasal inverted papilloma: systematic review and meta-analysis. *Am J Rhinol Allergy.* 2018;32:167-74.

Cellular indices and acute phase reactants as predictors of abscess after appendectomy

Índices celulares y reactantes de fase aguda como predictores de absceso tras apendicectomía

Julio C. Moreno-Alfonso^{1,2*}, Laura C. Lozano-Eslava³, Sharom Barbosa-Velásquez⁴, Ada Molina-Caballero¹, M. Concepción Yáñez-Irazábal⁵, and Alberto Pérez-Martínez¹

¹Pediatric Surgery Service, Hospital Universitario de Navarra, Pamplona, España; ²Doctoral School of Health Sciences, Universidad Pública de Navarra, Pamplona, España; ³Fundación Universitaria Sanitas, Bogotá, Colombia; ⁴Neurology Service, Hospital Universitario de Araba, Vitoria-Gasteiz, España; ⁵General and Digestive System Surgery Service, Hospital Universitario de Navarra, Pamplona, España

Abstract

Objective: To compare the accuracy of four biomarkers as predictors of post-appendectomy abdominal abscess (PAA). **Methods:** Diagnostic study of patients under 15 years of age operated for appendicitis in a pediatric hospital between 2010 and 2022 was analyzed. Neutrophil-lymphocyte ratio (NLR), platelet-lymphocyte ratio (PLR), C-reactive protein (CRP) and procalcitonin (PCT) were compared between patients with PAA and those without such complication (NPAA). **Results:** 86 patients with PAA (64% men; mean age: 9.4 ± 3.5 years) and 91 patients NPAA (65% men; mean age: 10.2 ± 2.9 years) were included. NLR, PLR, CRP and PCT values were higher in the PAA group than in the NPAA group ($p < 0.0001$). PCT had the highest area under the ROC curve (0.761), a cutoff point of 0.37, sensitivity 74%, specificity 75%, and positive predictive value of 74% for predicting the occurrence of PAA. **Conclusions:** PCT is a good predictor of abdominal abscess after appendectomy. Considering preoperative PCT values to optimize postoperative management and antibiotics could be an additional tool in the prevention of this complication.

Keywords: Appendicitis. Abdominal abscess. Procalcitonin. Neutrophil-lymphocyte ratio. C-reactive protein. Platelet-lymphocyte ratio.

Resumen

Objetivo: Comparar la precisión de cuatro biomarcadores como predictores de absceso abdominal posapendicectomía (APA). **Métodos:** Estudio diagnóstico de los pacientes menores de 15 años intervenidos por apendicitis en un hospital pediátrico entre 2010 y 2022. Se compararon el índice neutrófilo-linfocito (INL), el índice plaqueta-linfocito (IPL), la proteína C reactiva (PCR) y la procalcitonina (PCT) entre los pacientes con APA y los que no presentaron dicha complicación (NAPA). **Resultados:** Se incluyeron 86 pacientes con APA (64% hombres; edad media: 9.4 ± 3.5 años) y 91 pacientes NAPA (65% hombres; edad media: 10.2 ± 2.9 años). Los valores de INL, IPL, PCR y PCT fueron significativamente mayores en el grupo APA que en el grupo NAPA ($p < 0.0001$). La PCT tuvo el mayor área bajo la curva ROC (0.761), un punto de corte de 0.37, sensibilidad del 74%, especificidad del 75% y valor predictivo positivo del 74% para predecir la aparición de APA. **Conclusiones:** La PCT es un buen predictor de absceso abdominal tras una apendicectomía. Tener en cuenta los valores preoperatorios de PCT para optimizar el manejo y la antibioticoterapia en el posoperatorio podría ser una herramienta adicional en la prevención de esta complicación.

Palabras clave: Apendicitis. Absceso abdominal. Procalcitonina. Índice neutrófilo-linfocito. Proteína C reactiva. Índice plaqueta-linfocito.

*Correspondence:

Julio C. Moreno-Alfonso
E-mail: juliomorenosmd@gmail.com

Date of reception: 11-11-2023
Date of acceptance: 04-02-2025
DOI: 10.24875/CIRUE.M25000903

Cir Cir (Eng). 2025;93(6):644-649
Contents available at PubMed
www.cirugiaycirujanos.com

2444-0507/© 2025 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Acute appendicitis is the most frequent surgical emergency in pediatrics, with a lifetime cumulative risk of 9%¹. Its diagnosis fundamentally depends on clinical abdominal and general evaluation, although in pediatric age, early diagnosis is often difficult due to difficulties in anamnesis and the absence of classic symptoms². Appendicitis requires timely diagnosis and treatment, as its delay can lead to appendicular perforation and the development of complications.

Patients who undergo surgery for advanced appendicitis are at higher risk of developing complications, such as post-appendectomy abdominal abscess (APA). This complication occurs in approximately 4% of non-perforated appendicitis cases and in 6-28% of perforated cases, and is associated with a significant increase in morbidity and mortality^{3,4}. Although the value of different laboratory parameters as possible predictors of APA has been investigated, the results have been diverse⁵.

Recently, the usefulness of various biomarkers, such as the direct bilirubin-lymphocyte ratio, the neutrophil-lymphocyte ratio (NLR), and the platelet-lymphocyte ratio (PLR), has been evaluated as predictors of APA and in other surgical conditions, with disparate results⁶⁻⁸. However, scientific evidence is still limited, and there are no comparative studies on the usefulness of these ratios and other inflammatory parameters as predictors of this complication. The objective of this study is to compare the accuracy of four biomarkers as predictors of APA.

Methods

A retrospective diagnostic study was conducted on a consecutive series of patients under 15 years of age who developed APA, comparing them with a random sample of subjects undergoing surgery for acute appendicitis who did not present this complication (NAPA), in a second-level maternal-child hospital, between January 2010 and December 2022. Patients in the control group (NAPA) were sequentially selected from the same study population and during the same time period as patients with APA, regardless of the type of appendicitis, with the only mandatory characteristic being the absence of APA, in order to control for possible selection biases and make the groups comparable. Different variables were collected from the computerized medical records, including age, sex,

weight in kilograms, histological type of appendicitis, days of hospital stay, C-reactive protein (CRP) and procalcitonin (PCT) values, NLR and PLR at the time of hospital admission (these ratios are presented as continuous numerical values and have no units of measurement). Cell counts were performed with the Beckman Coulter hematological analyzer, and in all cases, the pathologist was unaware of the analytical results of the subjects and the occurrence of APA. Patients undergoing incidental appendectomy, with unorganized free fluid, or with an incomplete clinical record, were excluded from the study.

NLR and PLR were calculated by dividing the absolute number of neutrophils and platelets by the number of lymphocytes, respectively. The clinical diagnosis of acute appendicitis was based on anamnesis, physical examination, and ultrasound and analytical findings; the definitive diagnosis was based on histopathological analysis, classified as congestive, phlegmonous, gangrenous, perforated appendicitis, or peritonitis. The diagnosis of appendicular phlegmon was established during surgical intervention by identifying the cecal appendix with primary inflammatory signs and covered by a mass of intestinal loops with adjacent fibrin or edematous omentum. Localized peritonitis was defined as purulent or fecaloid contamination of one or two quadrants of the abdominal cavity, and generalized if there was contamination in three or more quadrants. Abscess was defined as a delimited collection of echogenic or purulent fluid within the abdominal cavity during the first month after an appendectomy, radiologically confirmed and accompanied by clinical signs of infection. All patients received intravenous antibiotic treatment with cefotaxime (150 mg/kg per day) and metronidazole (30 mg/kg per day) from the time of acute appendicitis diagnosis and during the postoperative period according to the type of appendicitis: three doses for phlegmonous, 3-5 days for gangrenous, and 7 days for the rest of appendicitis types.

Qualitative variables were expressed as relative and absolute frequencies, while quantitative variables were presented as mean and standard deviation or as median and interquartile range (IQR). The distribution of quantitative variables were determined using the Kolmogorov-Smirnov test to define the hypothesis testing method to be used (Mann-Whitney U or Student's t-test), and qualitative variables were analyzed using the χ^2 test or Fisher's exact test. Statistical significance was set as a p-value < 0.05 for an allowed type I error of 5%. CRP, PCT, NLR, and PLR were

evaluated to calculate their sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), and *odds ratio* (OR). The optimal cutoff point for predicting the occurrence of APA was obtained using the area under the receiver operating characteristic curve (AUC-ROC). According to the AUC-ROC, the accuracy of the test was considered poor (0.5-0.6), fair (0.61-0.7), acceptable (0.71-0.8), good (0.81-0.9), or excellent (> 0.9). Although reference values for some of these ratios have been described in the pediatric population, in this study, a specific cutoff point was calculated for each of them, as it is a particular pathological situation (APA)⁹.

The collected data were anonymized and tabulated in Microsoft Excel (Version 16.66.1) and analyzed using the SPSS statistical software (IBM Corp., Armonk, NY, USA). This work was approved by the Clinical Research Ethics Committee (Registration No. 3318-0000200) and followed the recommendations of the STARD guidelines (*Standards for Reporting Diagnostic Accuracy Studies*)¹⁰.

Results

During the period from 2010 to 2022, 1766 appendectomies were performed in patients under 15 years of age at the hospital under study. The APA rate was 5% ($n = 89$). This analysis included 86 children with APA who met the inclusion criteria and 91 appendectomized patients without postoperative collection as a comparative sample (Fig. 1). Of the total subjects with APA, 64% were male, similar to 64.8% in the group of appendectomized patients without complications ($p = 0.903$). The mean age of patients with APA was 9.4 ± 3.5 years, while in the NAPA group it was 10.2 ± 2.9 years ($p = 0.166$). Participants with APA had a weight of 35.8 ± 14.9 kg, while in the NAPA group, the average weight was 41.4 ± 17.1 kg ($p = 0.021$). No patient had a surgical history or relevant comorbidity, and all were operated on within the first 24 hours of admission (Table 1).

The main type of appendicitis in patients with APA versus the NAPA group was generalized peritonitis, at 32.6% ($n = 28/86$) vs. 2.2% ($n = 2/91$), followed by localized peritonitis at 31.4% ($n = 27/86$) vs. 8.8% ($n = 8/91$), perforation at 18.6% ($n = 16/86$) vs. 6.6% ($n = 6/91$), appendicular phlegmon at 10.5% ($n = 9/86$) vs. 2.2% ($n = 2/91$), gangrene at 3.5% ($n = 3/86$) vs. 13.2% ($n = 12/91$), and phlegmonous inflammation of the appendix at 3.5% ($n = 3/86$) vs. 61.5% ($n = 56/91$); only in the NAPA group were there congestive

appendicitis cases (5.5%, $n = 5/91$). These differences were statistically significant ($p < 0.0001$).

The most used surgical approach was trans umbilical video-assisted in 72% of NAPA patients and 73% of APA patients, followed by laparotomy access (9% vs. 8%) and laparoscopic (5% vs. 2%), with a similar rate of conversion to laparotomy from the laparoscopic or video-assisted approach (14% in NAPA patients vs. 17% in APA patients; $p > 0.05$). In no case was surgical drainage placed. The hospital stay for APA patients was 14 days (IQR: 5 days), while in the NAPA group it was 2 days (IQR: 4 days) ($p < 0.0001$).

The NLR value in the APA group versus the NAPA cohort was 12.2 vs. 6.4, the PLR was 260.8 vs. 153.1, CRP was 74.1 vs. 11.4 mg/l, and PCT was 0.90 vs. 0.10 ng/ml, respectively. All inflammatory markers were significantly higher in patients with APA compared to those without postoperative collection ($p < 0.0001$) (Table 1).

PCT was the inflammatory marker with the best performance and highest AUC-ROC for predicting the occurrence of APA (Fig. 2), with a cutoff point of 0.37 ng/ml, sensitivity of 74%, specificity of 75%, PPV of 74%, and NPV of 76%. The predictive capacity of each of the analyzed biomarkers is shown in detail in table 2.

Discussion

Abdominal abscesses occur in up to 20% of patients after appendectomy, especially in those with complicated appendicitis^{11,12}. Although different clinical and analytical prognostic variables have been evaluated in the pediatric population, including weight, the presence of appendicolith, or intolerance to oral intake in the postoperative period, among others, there are still gaps in the current literature regarding the usefulness of some parameters inflammatory^{13,14}. This study aimed to determine and compare the accuracy of CRP, PCT, NLR, and PLR as predictors of APA in pediatrics. This is the first study to determine and compare these parameters with the objective of analyzing their prognostic value in the occurrence of APA. It was found that CRP, PCT, NLR, and PLR values at admission were significantly higher in patients with APA compared to those without this complication, all showing acceptable accuracy. However, PCT was the biomarker that demonstrated the best performance.

The incidence of APA in our cohort was 5%, lower than reported in similar series where it reaches up to 18%¹⁵. Regarding prognostic clinical variables,




Figure 1. Flow diagram showing the potentially eligible, excluded, and included population in each of the two analyzed groups. APA: post-appendectomy abdominal abscess.

Table 1. Demographic characteristics and biomarker values in the studied population

Variable	APA (n = 86)	NAPA (n = 91)	p
Woman	36%	35%	0.903
Man	64%	65%	
Age (years)	9.4 ± 3.5	10.2 ± 2.9	0.166
Weight (kg)	35.8 ± 14.9	41.4 ± 17.1	0.021
Hours from admission to surgery	7.2 ± 3.6	8.1 ± 3.1	0.178
CRP (mg/l)	74.1 ± 170.3	11.4 ± 42.5	< 0.0001
PCT (ng/ml)	0.9 ± 4.8	0.1 ± 0.26	< 0.0001
NLR	12.2 ± 12.7	6.4 ± 6.2	< 0.0001
PLR	260.8 ± 150.7	153.1 ± 260.8	< 0.0001

APA: post-appendectomy abdominal abscess; NLR: neutrophil-lymphocyte ratio; PLR: platelet-lymphocyte ratio; NAPA: no APA; CRP: C-reactive protein; PCT: procalcitonin. Values are presented as median and interquartile range.




Figure 2. Area under the receiver operating characteristic curve of the analyzed parameters. NLR: neutrophil-lymphocyte ratio; PLR: platelet-lymphocyte ratio; CRP: C-reactive protein; PCT: procalcitonin.

Table 2. Area under the ROC curve and accuracy of each inflammatory marker studied as predictors of abdominal abscess after appendectomy

Biomarker	AUC-ROC	Cutoff point	S	AND	PPV	NPV	OR	p
CRP (mg/l)	0.741	21.9	72%	67%	67%	72%	5.25	<0.0001
PCT (ng/ml)	0.761	0.37	74%	75%	74%	76%	8.6	<0.0001
NLR	0.724	7.77	73%	63%	65%	71%	4.5	<0.0001
PLR	0.731	191.8	73%	73%	72%	74%	7.2	<0.0001

ROC: area under the receiver operating characteristic curve; E: specificity; NLR: neutrophil-lymphocyte ratio; PLR: platelet-lymphocyte ratio; OR: odds ratio; CRP: C-reactive protein; PCT: procalcitonin; S: sensitivity; NPV: negative predictive value; PPV: positive predictive value.

although overweight has been described as a risk factor for the development of APA because intraperitoneal adipose tissue can inhibit the inflammatory response to contamination of the abdominal cavity¹⁵, in our population, patients with APA had a lower weight than those without postoperative collection (35.8 ± 14.9 kg vs. 41.4 ± 17.1 kg; p = 0.021). However, this finding could have biases because the height of all patients was not recorded, and therefore, we do not know the body mass index. Likewise, perforation

of the cecal appendix has been shown to be a risk factor for the development of APA¹⁶, and although this association was not directly studied in this work, our results suggest the same, as there were more patients with appendicular perforation (18.6% vs. 6.6%) and peritonitis (64% vs. 11%) among patients with APA compared to the NAPA group. On the other hand, in this study, no statistically significant differences were found between the different surgical approaches and their association with the occurrence of APA, nor in the antibiotherapy used, since, according to the service protocol, all patients received the same antibiotic regimen before and after the surgical intervention.

Delgado et al.¹⁷ reported significantly higher CRP values in patients with APA compared to those without

complications (81.8 vs. 33.1 mg/l, respectively), with an AUC-ROC of 0.73, although with low sensitivity for the diagnosis of abscess (65%) and a discretely higher specificity of 72%. These results are similar to those found in our study, where CRP demonstrated a PPV of 67% for the occurrence of postoperative collection. However, in the experience of other authors, CRP showed no differences between patients with and without APA (8.20 vs. 5.36 mg/l; $p = 0.06$), and it was concluded that postoperative laboratory analyses have limited clinical utility for evaluating the development of peritoneal abscess¹⁸. These differences could be explained because the researchers in the mentioned study evaluated the value of CRP during the postoperative period, while in our study, inflammatory parameters were analyzed at the time of admission to the emergency department. Regarding PCT, Gavela et al.¹⁹ evaluated its use as a predictor of appendicitis severity and showed that it was significantly higher in patients with perforated appendicitis and peritonitis (4.95 ng/ml) than in those with simple appendicitis (0.15 ng/ml), with a specificity and PPV similar to ours, but with much higher sensitivity and NPV, 97% and 89%, respectively, compared to the present study, where sensitivity was 74% and NPV was 76%¹⁹. It is possible that these differences are due to discrepancies in the sociodemographic characteristics of each population and the primary objective of each study, which makes the results not entirely comparable.

Lodwick et al.²⁰ described lymphocytopenia as an effective predictor of post-appendectomy collection, which indirectly suggests that NLR could be a useful inflammatory parameter. Other authors have confirmed this hypothesis, reporting that NLR has an AUC-ROC of 0.85 and a cutoff point of 10.5 with a sensitivity of 85% and a specificity of 75% for predicting the occurrence of APA¹⁷. This predictive power of NLR is superior to that documented in our study (AUC-ROC 0.724, sensitivity 73%, and specificity 63%) when establishing the cutoff point at 7.7; this disparity in results is probably explained because a higher cutoff point was used in that study than in ours, which increases the sensitivity of the test but may increase the number of false negatives. In the same work, NLR demonstrated the best performance as a predictor of APA, compared to the absolute number of leukocytes, neutrophils, fibrinogen, and CRP¹⁷; however, a possible bias is that the population under 5 years of age was excluded. Considering the age group of patients is of great clinical importance; in fact, Aydođdu et al.²¹ evaluated different cellular

indices in the diagnosis of pediatric appendicitis and concluded that, in childhood, the reference values of NLR, PLR, and lymphocyte-monocyte ratio vary according to age and sex, which should be taken into account when evaluating blood test results.

During the inflammatory process of acute appendicitis, an immune reaction that also involves platelets is triggered. Therefore, platelet changes have been studied to determine the usefulness of PLR in the diagnosis and prognosis of appendicitis²². Our analysis documented a significantly higher PLR value in patients with APA (260.8) than in those without this complication (153.1), which was associated with an increase of up to 7.2 times in the probability of developing APA (OR: 7.2; 95% confidence interval: 3.7-14). These findings are consistent with what has been described in the literature, documenting that patients with higher PLR values were more likely to present complicated appendicitis and postoperative complications⁸.

This study describes the usefulness of PCT as a predictor of APA in pediatric patients undergoing acute appendicitis surgery, with better diagnostic performance than CRP, NLR, and PLR. Due to its wide availability, healthcare personnel should be aware of its potential role in the prevention, diagnosis, and timely treatment of APA, with the aim of reducing the morbidity and mortality associated with this complication. In patients with a reasonable clinical suspicion of APA, with symptoms such as fever and intolerance to oral intake during the postoperative period, elevated preoperative PCT values support the urgent performance of an imaging test. This study presents the limitations inherent to a retrospective study in a single hospital center, in addition to the physiological heterogeneity in blood cell counts in the pediatric population. To our knowledge, this is the first study to compare these four inflammatory markers as predictors of APA, therefore, prospective studies with a larger number of patients are required to corroborate our results and seek prevention strategies for the development of APA, such as adjusting the perioperative fluid therapy and antibiotic protocol in patients at high risk of peritoneal collection.

Conclusions

PCT is a good biomarker as a predictor of APA in the pediatric population. The evaluation of PCT values before surgical intervention would allow optimizing perioperative management and postoperative antibiotic therapy, becoming a valuable tool to prevent this complication.

Funding

The authors declare that they have not received funding for this study.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Ethical considerations

Protection of people and animals. The authors declare that no experiments on humans or animals were performed for this research.

Confidentiality, informed consent, and ethical approval. The authors have followed their institution's confidentiality protocols. The authors obtained approval from the Ethics Committee for the analysis of routinely obtained and anonymized clinical data, so informed consent was not required. This work was approved by the Clinical Research Ethics Committee (Registration No. 3318-0000200) and has followed the recommendations of the STARD guidelines (*Standards for Reporting Diagnostic Accuracy Studies*).

Declaration on the use of artificial intelligence. The authors declare that they did not use any type of generative artificial intelligence for the writing of this manuscript.

References

- Prada M, Salgado A, Montero M, Fernández P, García S, Gómez J, et al. Appendicitis versus non-specific acute abdominal pain: Paediatric Appendicitis Score evaluation. *An Paediatr (Engl Ed)*. 2018;88:32-8. doi: 10.1016/j.anpedi.2017.01.006.
- Aneiros B, Cano I, García A, Yuste P, Ferrero E, Gómez A. Pediatric appendicitis: age does make a difference. *Rev Paul Pediatr*. 2019;37:318-24. doi: 10.1590/1984-0462/2019;37;3;00019.
- Rentea RM, St. Peter SD. Contemporary management of appendicitis in children. *Adv Pediatr*. 2017;64:225-51. doi: 10.1016/j.yapd.2017.03.008.
- Sarda S, Short HL, Hockenberry JM, McCarthy I, Raval MV. Regional variation in rates of pediatric perforated appendicitis. *J Pediatr Surg*. 2017;52:1488-91. doi: 10.1016/j.jpedsurg.2017.02.009.
- Dickinson CM, Coppersmith NA, Luks FI. Early predictors of abscess development after perforated pediatric appendicitis. *Surg Infect (Larchmt)*. 2017;18:886-9. doi: 10.1089/sur.2017.134.
- Jung SK, Rhee DY, Lee WJ, Woo SH, Seol SH, Kim DH, et al. Neutrophil-to-lymphocyte count ratio is associated with perforated appendicitis in elderly patients of emergency department. *Aging Clin Exp Res*. 2017;29:529-36. doi: 10.1007/s40520-016-0584-8.
- Celik B, Nalcacioglu H, Ozcatat M, Altuner Y. Role of neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio in identifying complicated appendicitis in the pediatric emergency department. *Ulus Travma Acil Cerrahi Derg*. 2019;25:222-8. doi: 10.5505/tjtes.2018.06709.
- Azizoglu M, Arslan S, Kamci TO, Basuguy E, Aydogdu B, Karabel MA, et al. Can direct bilirubin-to-lymphocyte ratio predict surgery for pediatric adhesive small bowel obstruction? *Cir Cir*. 2024;92:307-13. doi: 10.24875/CIRU.23000524.
- Moosmann J, Krusemark A, Dittrich S, Ammer T, Rauh M, Woelfle J, et al. Age- and sex-specific pediatric reference intervals for neutrophil-to-lymphocyte ratio, lymphocyte-to-monocyte ratio, and platelet-to-lymphocyte ratio. *Int J Lab Hematol*. 2022;44:296-301. doi: 10.1111/ijlh.13768.
- Bossuyt PM, Reitsma JB, Bruns DE, Gatsonis CA, Glasziou PP, Irwig L, et al. STARD 2015: an updated list of essential items for reporting diagnostic accuracy studies. *BMJ*. 2015;351:h5527. doi: 10.1136/bmj.h5527.
- Morrow SE, Newman KD. Current management of appendicitis. *Semin Pediatr Surg*. 2007;16:34-40. doi: 10.1053/j.sempedsurg.2006.10.005.
- Singh M, Kadian YS, Rattan KN, Jangra B. Complicated appendicitis: analysis of risk factors in children. *Afr J Paediatr Surg*. 2014;11:109-13. doi: 10.4103/0189-6725.132796.
- Hughes MJ, Harrison E, Paterson-Brown S. Post-operative antibiotics after appendectomy and post-operative abscess development: a retrospective analysis. *Surg Infect*. 2013;14:56-61. doi: 10.1089/sur.2011.100.
- Henry MC, Walker A, Silverman BL, Gollin G, Islam S, Sylvester K, et al. Risk factors for the development of abdominal abscess following operation for perforated appendicitis in children: a multicenter case-control study. *Arch Surg*. 2007;142:236-41. doi: 10.1001/archsurg.142.3.236.
- Fraser JD, Aguayo P, Sharp SW, Snyder CL, Holcomb GW 3rd, Ostlie DJ, et al. Physiologic predictors of postoperative abscess in children with perforated appendicitis: subset analysis from a prospective randomized trial. *Surgery*. 2010;147:729-32. doi: 10.1016/j.surg.2009.10.057.
- Emil S, Elkady S, Shbat L, Youssef F, Baird R, Laberge JM, et al. Determinants of postoperative abscess occurrence and percutaneous drainage in children with perforated appendicitis. *Pediatr Surg Int*. 2014;30:1265-71. doi: 10.1007/s00383-014-3617-4.
- Delgado C, Muñoz AJ, Núñez V, Estefanía K, Velayos M, Miguel M, et al. Neutrophil-to-lymphocyte ratio as a predictor of postsurgical intraabdominal abscess in children operated for acute appendicitis. *Front Pediatr*. 2019;7:424. doi: 10.3389/fped.2019.00424.
- Zheng S, Christy K, Herzak K, Kobal N, Novak M, Young J, et al. Labs do not predict postoperative intra-abdominal abscess in pediatric perforated appendicitis. *J Surg Res*. 2023;285:20-5. doi: 10.1016/j.jss.2022.12.015.
- Gavela T, Cabeza B, Serrano A, Casado J. C-reactive protein and procalcitonin are predictors of the severity of acute appendicitis in children. *Pediatr Emerg Care*. 2012;28:416-9. doi: 10.1097/PEC.0b013e318252d875.
- Lodwick DL, Cooper JN, Kenney B, Deans KJ, Minneci PC, Thakkar RK. Lymphocyte depression as a predictor of postoperative intraabdominal abscess after appendectomy in children. *J Pediatr Surg*. 2017;52:93-7. doi: 10.1016/j.jpedsurg.2016.10.028.
- Aydoğdu B, Azizoğlu M, Arslan S, Aydoğdu G, Basuguy E, Salik F, et al. A novel diagnostic scoring system for pediatric appendicitis based on age and sex-adjusted hematological parameters. *Gac Med Mex*. 2023;159:103-9. doi: 10.24875/GMM.M22000753.
- Yazar FM, Bakacak M, Emre A, Urfalioglu A, Serin S, Cengiz E, et al. Predictive role of neutrophil-to-lymphocyte and platelet-to-lymphocyte ratios for diagnosis of acute appendicitis during pregnancy. *Kaohsiung J Med Sci*. 2015;31:591-6. doi: 10.1016/j.kjms.2015.10.005.

Effects of ropivacaine alone or in combination with dexmedetomidine on cesarean section: a systematic review and meta-analysis

Efectos de la ropivacaína sola o en combinación con dexmedetomidina en la cesárea: revisión sistemática y metaanálisis

Huaping Li¹, Mingwei Kan², and Rong Jia^{3*}

¹Department of Anesthesiology, Logistic Support Forces of the Chinese People's Liberation Army No. 989 Hospital, Henna Province; ²Department of Anesthesiology, The Fifth Affiliated Hospital of Zunyi Medical University, Guangdong Province; ³Department of Anesthesiology, Ningxia Armed Police Corps Hospital, Ningxia Hui Autonomous Region, China

Abstract

Objective: Different from physiological pain, post-operative pain is caused by surgical trauma. We aimed to systematically assess the effects of ropivacaine alone or in combination with dexmedetomidine on cesarean section and to conduct a meta-analysis. **Methods:** According to the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) statement and Cochrane intervention system review manual, retrieval software and data analysis tools were used for literature retrieval, screening, exclusion and inclusion, data extraction, analysis and statistics, and risk assessment. A total of 37 literatures were retrieved, and 11 literatures were left after 26 duplicates were excluded. **Results:** A total of 37 literatures were retrieved in all databases, of which 11 literatures were left and finally 7 all published in English were obtained. They were seven randomized controlled trials on ropivacaine alone or in combination with dexmedetomidine applied in 502 parturients receiving cesarean section. The results of meta-analysis on main outcome indices showed that the number of parturients in need of rescue ($\chi^2 = 28.62, p < 0.001, I^2 = 93\%$), incidence rate of adverse reactions ($\chi^2 = 28.66, p = 0.007, I^2 = 55\%$), satisfaction ($\chi^2 = 7.97, p = 0.05, I^2 = 62.3\%$), visceral respiratory response ($\chi^2 = 19.26, p < 0.001, I^2 = 89.6\%$), satisfaction with muscle relaxation ($\chi^2 = 6.92, p = 0.03, I^2 = 71.1\%$), and spinal anesthesia grade ($\chi^2 = 25.89, p < 0.01, I^2 = 92.3\%$). **Conclusions:** Ropivacaine combined with dexmedetomidine has a better prognostic effect on cesarean section and causes fewer adverse reactions.

Keywords: Ropivacaine. Dexmedetomidine. Cesarean section. Meta-analysis.

Resumen

Objetivo: A diferencia del dolor fisiológico, el dolor posoperatorio está ocasionado por un trauma quirúrgico. El objetivo es evaluar sistemáticamente los efectos de la ropivacaína sola o en combinación con dexmedetomidina en la cesárea y realizar un metaanálisis. **Métodos:** De acuerdo con la declaración PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses Statement) y el Cochrane Intervention System Review Manual, se utilizaron software y herramientas de análisis de datos para la recuperación de la literatura, los exámenes de detección, la exclusión y la inclusión, la extracción de datos, el análisis y las estadísticas, y la evaluación del riesgo. Se obtuvieron 37 estudios, de los cuales quedaron 11 después de excluir 26 duplicados. **Resultados:** De las bases de datos se obtuvieron 37 estudios, de los cuales se seleccionaron 11 y finalmente se obtuvieron 7, todos publicados en inglés. Correspondieron a 7 ensayos controlados aleatorizados sobre ropivacaína sola o en combinación con dexmedetomidina en 502 parturientas sometidas a cesárea. Los resultados del metaanálisis

*Correspondence:

Rong Jia

E-mail: tunluke4@163.com

2444-0507/© 2024 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 27-10-2023

Date of acceptance: 09-04-2024

DOI: 10.24875/CIRUE.M24000904

Cir Cir (Eng). 2025;93(6):650-659

Contents available at PubMed

www.cirugiaycirujanos.com

sobre los principales índices de resultados mostraron que el número de parturientas con necesidad de rescate ($\chi^2 = 28.62$; $p < 0.001$; $I^2 = 93\%$), la tasa de incidencia de reacciones adversas ($\chi^2 = 28.66$; $p = 0.007$; $I^2 = 55\%$), la satisfacción ($\chi^2 = 7.97$; $p = 0.05$; $I^2 = 62.3\%$), la respuesta respiratoria visceral ($\chi^2 = 19.26$; $p < 0.001$; $I^2 = 89.6\%$), la satisfacción con la relajación muscular ($\chi^2 = 6.92$; $p = 0.03$; $I^2 = 71.1\%$), el grado de anestesia espinal ($\chi^2 = 25.89$; $p < 0.01$; $I^2 = 92.3\%$). **Conclusiones:** La ropivacaína combinada con dexmedetomidina tiene un mejor efecto pronóstico en la cesárea y provoca menos reacciones adversas.

Palabras clave: Ropivacaína. Dexmedetomidina. Cesárea. Metaanálisis.

Introduction

Different from physiological pain, post-operative pain is caused by surgical trauma. After tissue injury at the surgical incision, inflammatory mediators are released under stress, and then the sensory nerves feel the conduction of stimulus of inflammatory mediators to the center. The pain sensitivity of patients also changes postoperatively, thereby producing an unpleasant emotional expression in the cerebral cortex, which is the protective response of the body to harmful stimuli¹. Post-operative pain is one of the most common problems faced by patients, which is not conducive to the rapid post-operative recovery of patients, and the severer the pain is, the greater the patient's physiological and psychological fluctuation will be². In general, the pain is the severest within 2 days after surgery, and it mostly lasts for < 1 week. If not controlled well, post-operative pain may transform into chronic pain or persistent pain, and even neuropathic pain in severe cases.

With the opening of the two-child policy in China, the cesarean section rate has been increasing. After cesarean section, incision pain and uterine contraction pain are serious problems faced by parturients³. Postpartum pain is an important factor causing anxiety, depression and insomnia in parturients, seriously affecting the post-operative rehabilitation and lactation, and also affecting parturients' care for newborns. Therefore, multi-modal analgesia is adopted after cesarean section. Post-operative analgesia methods commonly used in clinic include patient-controlled epidural analgesia (PCEA), patient-controlled intravenous analgesia (PCIA), oral administration or intramuscular injection and even intravenous infusion of opioids or antipyretic analgesics, and ultrasound-guided plane block of transverse abdominal muscle (TAP) popular in recent years^{4,5}. Nowadays when minimally-invasive surgery is strongly advocated, cesarean section is considered as a cruel operation, after which parturients

often suffer from highly severe pain. At present, PCEA and PCIA are mainly adopted by anesthesiologists for post-operative analgesia. However, these two analgesia methods lead to serious complications such as nausea and vomiting, pruritus, respiratory depression, epidural hematoma, hemodynamic instability, and even unsatisfactory analgesia⁶. Besides, systemic administration has limitations of drugs and short duration of action, causes severe gastrointestinal discomfort, nausea, vomiting, increased urinary retention, hepatic and renal dysfunction, and increased frequency of administration, and affects the rest and sleep quality of parturients.

At present, the local anesthetics used in TAP include lidocaine, bupivacaine, and ropivacaine. Among them, ropivacaine characterized by long duration of action, low toxicity, and good sensorimotor block separation has become a preferred local anesthetic in field block for anesthesiologists⁷. Despite a good analgesic effect in clinic, ropivacaine has short duration of block and fails to meet the demand for post-operative analgesia with one-time administration⁸. Therefore, local anesthetics with longer duration of action are needed, or local anesthetic adjuvants need to be rationally used to prolong the duration of action of local anesthetics, without causing any unnecessary complications to increase the duration of sensory block. Dexmedetomidine, a research hotspot in the field of anesthesia and analgesia, is often applied. As a highly-selective α -adrenergic receptor agonist, dexmedetomidine is characterized by sedation, anti-anxiety, analgesia, stabilization of hemodynamics, inhibition on sympathetic nerve activity, and mild respiratory depression⁹. At present, dexmedetomidine has been widely used in intensive care unit sedation, non-invasive endoscopic examination, pre-operative medication, adjuvant medication during anesthesia, and post-operative analgesia¹⁰. Dexmedetomidine has been combined with bupivacaine to markedly prolong the sensory and motor block duration and to improve intraoperative and post-operative analgesia for women undergoing

cesarean section, without any significant complications for the mothers or infants¹¹. However, there are few reports on the combination of dexmedetomidine and ropivacaine for analgesia in cesarean section. The present study aims to observe the analgesic effect of dexmedetomidine combined with ropivacaine on cesarean section and related adverse reactions, so as to provide a new idea for multi-modal analgesia.

Methods

Literature retrieval method

The retrieval strategy was developed according to Cochrane international standards and the PRISMA flowchart, and the English databases included PubMed, Web of Science, and EMBase.

- English retrieval words
Dexmedetomidine, ropivacaine, cesarean delivery or cesarean section, and randomized controlled trial.
- Retrieval time period
The retrieval was restricted to human randomized controlled trials in the above databases from the establishment to April 30, 2021.

Data extraction

The basic information included the research title, first author, publication time, control group and observation group, intervention measures, and indices.

Literature quality assessment

The quality of included literatures was assessed using the Cochrane risk of bias assessment tool, criteria for judging risk of bias in the risk of bias assessment tool, and possible approach for summary assessments of the risk of bias for each important outcome within and across studies designated by Cochrane.

Statistical analysis

Meta-analysis was conducted for indices enrolled using RevMan5.4 provided by Cochrane, and the forest plot made by RevMan software was used to assess the publication bias.

Random effects model (REM) was used in the case of $I^2 > 50\%$ in heterogeneity test, while fixed effects model (FEM) was used in the case of $I^2 \leq 50\%$. The




Figure 1. Preferred reporting Items for systematic reviews and meta-analysis flowchart.

effect scale indices included two-category data and continuous data.

Results

Literature retrieval results

A total of 37 literatures were retrieved in all databases, and 11 literatures were left after 26 duplicates were excluded. Then, one literature was excluded according to the inclusion and exclusion criteria, and three literatures were excluded after reading the full text. Finally, seven literatures all published in English were obtained (Fig. 1).

Basic characteristics of included studies

The indices, that is, the number of parturients in need of rescue, adverse reactions, satisfaction, incidence rate of transient neurological syndrome (TNS), visceral respiratory response, satisfaction with muscle relaxation, spinal anesthesia grade, mean time for rescue analgesia, visual analog scale (VAS) score, 24 h mean cumulative dose for rescue analgesia, pain-free

Table 1. Basic characteristics of included studies

First author	Published year	Treatment/control group	Treatment method	Outcome index
Bi ¹²	2020	25/25	Ropivacaine/ropivacaine + dexmedetomidine	1, 5, 6, 7
Qian ¹³	2020	35/35	Ropivacaine/ropivacaine + dexmedetomidine	1,9, 11, 12
Bhardwaj ¹⁴	2017	30/30	Ropivacaine/ropivacaine + dexmedetomidine	1, 2, 3, 8, 9, 10
Kong ¹⁵	2018	51/51	Ropivacaine/ropivacaine + dexmedetomidine	2, 4, 9, 13, 14
Mo and Qiu ¹⁶	2017	40/40	Ropivacaine/ropivacaine + dexmedetomidine	2, 9, 13
Tang ¹⁷	2020	30/30	Ropivacaine/ropivacaine + dexmedetomidine	2
Zhao ¹⁸	2017	40/40	Ropivacaine/ropivacaine + dexmedetomidine	9

1: number of parturients in need of rescue; 2: adverse reactions; 3: satisfaction; 4: incidence rate of TNS; 5: visceral respiratory response; 6: satisfaction with muscle relaxation; 7: spinal anesthesia grade; 8: mean time for rescue analgesia; 9: VAS score; 10: 24 h mean cumulative dose for rescue analgesia; 11: pain-free duration; 12: time to first analgesia required; 13: Ramsay sedation score; 14: MMSE score.

duration, time to first analgesia required, Ramsay sedation score, and mini-mental state examination (MMSE) score, were covered in three studies, five studies, one study, one study, one study, one study, one study, one study, five studies, one study, one study, one study, two studies, and one study, respectively (Table 1).

Quality assessment of included studies

The randomization method was known in all of the seven included studies. The allocation concealment was known in two studies and unknown in the remaining five studies. The blind method for subjects was unknown in three studies, and the blind method for result assessment was unknown in two studies. The outcome indices were incomplete in one study, and the selective reporting was high risk in one study and also unknown in one study (Fig. 2).

Meta-analysis and publication bias

NUMBER OF PARTURIENTS IN NEED OF RESCUE

As shown in figure 3, the number of parturients in need of rescue was reported in three studies, the results belonged to two-category data, and REM was used ($I^2 > 50\%$). The results showed that the number of parturients in need of rescue in ropivacaine + dexmedetomidine group was significantly smaller than that in ropivacaine control group, and the difference was statistically significant ($\chi^2 = 28.62$, $p < 0.001$, $I^2 = 93\%$), indicating that ropivacaine combined with dexmedetomidine can significantly improve the prognosis of parturients receiving cesarean section.

INCIDENCE RATE OF ADVERSE REACTIONS

As shown in figure 4, the incidence rate of adverse reactions was reported in five studies, the results belonged to two-category data, and REM was used ($I^2 > 50\%$). The results showed that the incidence rate of adverse reactions in ropivacaine + dexmedetomidine group was significantly lower than that in ropivacaine control group, and the difference was statistically significant ($\chi^2 = 28.66$, $p = 0.007$, $I^2 = 55\%$), indicating that ropivacaine combined with dexmedetomidine can significantly reduce the incidence rate of clinical adverse reactions in parturients receiving cesarean section.

SATISFACTION OF PARTURIENTS

As shown in figure 5, the satisfaction of parturients was reported in one study, the results belonged to two-category data, and REM was used ($I^2 > 50\%$). The results showed that the number of parturients with excellent satisfaction in ropivacaine + dexmedetomidine group was significantly larger than that in ropivacaine control group, and the difference was statistically significant ($\chi^2 = 7.97$, $p = 0.05$, $I^2 = 62.3\%$), indicating that ropivacaine combined with dexmedetomidine can significantly enhance the satisfaction of parturients receiving cesarean section.

INCIDENCE RATE OF TNS

As exhibited in figure 6, the incidence rate of TNS was reported in 1 study, the results belonged to two-category data and FEM was used ($I^2 < 50\%$). The results revealed that the incidence rate of TNS in ropivacaine +




Figure 2. A and B: Risk of bias of included literatures.




Figure 3. Forest plot of meta-analysis on the number of parturients in need of rescue in the two groups.

dexmedetomidine group was significantly lower than that in ropivacaine control group, and there was no statistically significant difference ($\chi^2 = 0.16$, $p = 0.93$, $I^2 = 0\%$).

VISCERAL RESPIRATORY RESPONSE

As presented in supplementary data figure S1, the visceral respiratory response of parturients was reported in one study, the results belonged to two-category data,

and REM was used ($I^2 > 50\%$). The results revealed that the number of parturients with visceral respiratory response in grade 2 and 3 in ropivacaine + dexmedetomidine group was significantly smaller than that in ropivacaine control group, and there was a statistically significant difference ($\chi^2 = 19.26$, $p < 0.001$, $I^2 = 89.6\%$), suggesting that ropivacaine combined with dexmedetomidine can improve the visceral respiratory response of parturients receiving cesarean section.




Figure 4. Forest plot of meta-analysis on the incidence rate of adverse reactions in the two groups.

SATISFACTION WITH MUSCLE RELAXATION

As displayed in supplementary data figure S2, the satisfaction of parturients with muscle relaxation was

reported in one study, the results belonged to two-category data and REM was used (I² > 50%). The results revealed that the satisfaction of parturients with muscle relaxation in ropivacaine + dexmedetomidine




Figure 5. Forest plot of meta-analysis on the satisfaction of parturients in the two groups.




Figure 6. Forest plot of meta-analysis on the incidence rate of transient neurological syndrome in the two groups.

group was significantly higher than that in ropivacaine control group, and there was a statistically significant difference ($\chi^2 = 6.92$, $p = 0.03$, $I^2 = 71.1\%$), suggesting

that ropivacaine combined with dexmedetomidine can raise the satisfaction of parturients receiving cesarean section with muscle relaxation.

SPINAL ANESTHESIA GRADE

The spinal anesthesia grade of parturients was reported in 1 study, the results belonged to two-category data and REM was used ($I^2 > 50\%$) (Supplementary data Fig. S3). The results manifested that the number of parturients receiving spinal anesthesia in grade 2 and 3 in ropivacaine + dexmedetomidine group was significantly smaller than that in ropivacaine control group, and there was a statistically significant difference ($\chi^2 = 25.89$, $p < 0.01$, $I^2 = 92.3\%$), suggesting that ropivacaine combined with dexmedetomidine can lower the spinal anesthesia grade of parturients receiving cesarean section.

MEAN TIME FOR RESCUE ANALGESIA

The mean time for rescue analgesia of parturients was reported in one study, the results belonged to continuous data, and REM was used ($I^2 > 50\%$) (Supplementary data Fig. S4). The results manifested that the mean time for the 4th and 5th rescue analgesia was significantly shorter in ropivacaine + dexmedetomidine group than that in ropivacaine control group, while the mean time for the 1st to 3rd rescue analgesia was significantly longer in ropivacaine + dexmedetomidine group than that in ropivacaine control group, showing statistically significant differences ($\chi^2 = 15.31$, $p = 0.0005$, $I^2 = 87\%$). It can be seen that the mean time for rescue analgesia is decreased under ropivacaine combined with dexmedetomidine compared with that under ropivacaine.

VAS SCORE

The VAS score of parturients was reported in five studies, the results belonged to continuous data, and REM was used ($I^2 > 50\%$) (Supplementary data Fig. S5). The results manifested that the VAS score at T1-T4 was significantly lower in ropivacaine + dexmedetomidine group than that in ropivacaine control group, showing a statistically significant difference ($\chi^2 = 66.68$, $p < 0.001$, $I^2 = 96\%$; $\chi^2 = 54.85$, $p < 0.001$, $I^2 = 95\%$; $\chi^2 = 16.59$, $p = 0.002$, $I^2 = 88\%$; $\chi^2 = 11.86$, $p = 0.003$, $I^2 = 83\%$). It can be seen that the VAS score obviously declines under ropivacaine combined with dexmedetomidine compared with that under ropivacaine.

24 H MEAN CUMULATIVE DOSE FOR RESCUE ANALGESIA

The 24 h mean cumulative dose for rescue analgesia of parturients was reported in one study, and the results belonged to continuous data (Supplementary data Fig. S6). It was found that the 24 h mean cumulative dose for rescue analgesia was significantly lower in ropivacaine + dexmedetomidine group than that in ropivacaine control group, displaying a statistically significant difference ($p < 0.001$). It can be seen that the 24 h mean cumulative dose for rescue analgesia obviously declines under ropivacaine combined with dexmedetomidine compared with that under ropivacaine.

PAIN-FREE DURATION

The pain-free duration of parturients was reported in one study, and the results belonged to continuous data (Supplementary data Fig. S7). It was found that the pain-free duration was significantly longer in ropivacaine + dexmedetomidine group than that in ropivacaine control group, and the difference was statistically significant ($p < 0.001$). It can be seen that the pain-free duration is prolonged and the prognosis is good under ropivacaine combined with dexmedetomidine compared with those under ropivacaine.

TIME TO FIRST ANALGESIA REQUIRED

The time to first analgesia required was reported in one study, and the results belonged to continuous data (Supplementary data Fig. S8). It was found that the time to first analgesia required was significantly later in ropivacaine + dexmedetomidine group than that in ropivacaine control group, and the difference was statistically significant ($p < 0.001$). It can be seen that the time to first analgesia required is later and parturients have milder pain under ropivacaine combined with dexmedetomidine than those under ropivacaine.

RAMSAY SEDATION SCORE

The Ramsay sedation score of parturients was reported in two studies, the results belonged to continuous data, and REM was used ($I^2 > 50$) (Supplementary data Fig. S9). It was found that the Ramsay sedation score at T1-T2 was significantly better in ropivacaine + dexmedetomidine group than that in

ropivacaine control group, and the difference was statistically significant ($\chi^2 = 12.36$, $p < 0.001$, $I^2 = 92\%$; $\chi^2 = 102.53$, $p < 0.001$, $I^2 = 99\%$), indicating that the Ramsay sedation score greatly rises and the prognosis is good under ropivacaine combined with dexmedetomidine compared with those under ropivacaine.

MMSE SCORE

The MMSE score of parturients was reported in 1 study, the results belonged to continuous data, and REM was used ($I^2 > 50$) (Supplementary data Fig. S10). The results showed that the MMSE score after 1 d was significantly superior in ropivacaine + dexmedetomidine group to that in ropivacaine control group, and the difference was statistically significant ($\chi^2 = 67.60$, $p < 0.001$, $I^2 = 99\%$), indicating that the MMSE score after 1 d greatly rises and the prognosis is good under ropivacaine combined with dexmedetomidine compared with those under ropivacaine.

Discussion

All the seven included studies had a small sample size (< 200 cases), so false-positive or false-negative results caused by random errors could not be excluded.

Several outcome indices were selected from the seven literatures, including the number of parturients in need of rescue, adverse reactions, satisfaction, incidence rate of TNS, visceral respiratory response, satisfaction with muscle relaxation, spinal anesthesia grade, mean time for rescue analgesia, VAS score, 24 h mean cumulative dose for rescue analgesia, pain-free duration, time to first analgesia required, Ramsay sedation score, and MMSE score¹²⁻¹⁸. The authenticity and reliability of the above outcome indices were assessed using the RevMan software.

The randomization method was known in all of the seven included studies. The allocation concealment was known in two studies and unknown in the remaining five studies. The blind method for subjects was unknown in three studies, and the blind method for result assessment was unknown in two studies. The outcome indices were incomplete in one study, and the selective reporting was high risk in one study and also unknown in one study.

The number of parturients in need of rescue, adverse reactions, satisfaction, incidence rate of TNS, visceral respiratory response, satisfaction with muscle relaxation, and spinal anesthesia grade were all

two-category data. The results showed that ropivacaine + dexmedetomidine group had a significantly smaller number of parturients in need of rescue, those with visceral respiratory response in grade 2 and 3 and those receiving spinal anesthesia in grade 2 and 3, and significantly lower incidence rates of adverse reactions and TNS than ropivacaine control group, while ropivacaine + dexmedetomidine group had a significantly larger number of parturients with excellent satisfaction and significantly better satisfaction with muscle relaxation than ropivacaine control group. It can be seen that ropivacaine combined with dexmedetomidine can significantly improve the prognosis and satisfaction of parturients receiving cesarean section and reduce the incidence rate of clinical adverse reactions.

The mean time for rescue analgesia, VAS score, 24 h mean cumulative dose for rescue analgesia, pain-free duration, time to first analgesia required, Ramsay sedation score, and MMSE score were all continuous data. The results revealed that the mean time for the 4th and 5th rescue analgesia was significantly shorter, the VAS score at T1-T4 and 24 h mean cumulative dose for rescue analgesia were significantly lower, and the time to first analgesia required was significantly later in ropivacaine + dexmedetomidine group than those in ropivacaine control group, while the Ramsay sedation score at T1-T2 and MMSE score after 1 d were significantly superior in ropivacaine + dexmedetomidine group to those in ropivacaine control group. The above findings demonstrate that parturients in ropivacaine + dexmedetomidine group had milder pain, a better prognosis and a significantly better mental state than those in ropivacaine control group.

Nevertheless, this study is limited. The number of patients is small. In addition, the results between many studies are not compared, which may have a negative impact on the conclusion.

Conclusions

Ropivacaine combined with dexmedetomidine has a better prognostic effect and causes fewer adverse reactions in parturients receiving cesarean section under anesthesia. The meta-analysis results remain to be improved with larger-sample higher-quality studies.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of humans and animals. The authors declare that no experiments involving humans or animals were conducted for this research.

Confidentiality, informed consent, and ethical approval. The authors have followed their institution's confidentiality protocols. The study does not involve patient personal data.

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing of this manuscript.

This study approved by Ningxia Armed Police Corps Hospital.

Supplementary data

Supplementary data are available at DOI: 10.24875/CIRU.23000532. These data are provided by the corresponding author and published online for the benefit of the reader. The contents of supplementary data are the sole responsibility of the authors.

References

- Lee A, Blair HA. Dexamethasone intracanalicular insert: a review in treating post-surgical ocular pain and inflammation. *Drugs*. 2020; 80:1101-8.
- Osorio W, Ceballos C, Moyano J. Effectiveness of acute post-operative pain management by the acute pain service. *Cir Cir*. 2022;90:197-201.
- Vázquez-Rodríguez JG, Serrano-Rodríguez J, Arredondo-Andrade SA, Chable-Chan FG, García-Bello JA, Nájera-Coutiño CM. Prevalence, causes, and clinical course of patients with surgical reintervention due to complications during cesarean section. *Cir Cir*. 2023;91:446-50.
- Halpern SH, Muir H, Breen TW, Campbell DC, Barrett J, Liston R, et al. A multicenter randomized controlled trial comparing patient-controlled epidural with intravenous analgesia for pain relief in labor. *Anesth Analg*. 2004;99:1532-8.
- Blanco R, Ansari T, Riad W, Shetty N. Quadratus lumborum block versus transversus abdominis plane block for postoperative pain after cesarean delivery: a randomized controlled trial. *Reg Anesth Pain Med*. 2016; 41:757-62.
- Jin J, Min S, Chen Q, Zhang D. Patient-controlled intravenous analgesia with tramadol and lornoxicam after thoracotomy: a comparison with patient-controlled epidural analgesia. *Medicine (Baltimore)*. 2019; 98:e14538.
- Cacciamani GE, Menestrina N, Pirozzi M, Tafuri A, Corsi P, De Marchi D, et al. Impact of combination of local anesthetic wounds infiltration and ultrasound transversus abdominal plane block in patients undergoing robot-assisted radical prostatectomy: perioperative results of a double-blind randomized controlled trial. *J Endourol*. 2019;33:295-301.
- Sun N, Wang S, Ma P, Liu S, Shao A, Xiong L. Postoperative analgesia by a transversus abdominis plane block using different concentrations of ropivacaine for abdominal surgery: a meta-analysis. *Clin J Pain*. 2017; 33:853-63.
- Weerink MA, Struys MM, Hannivoort LN, Barends CR, Absalom AR, Colin P. Clinical pharmacokinetics and pharmacodynamics of dexmedetomidine. *Clin Pharmacokinet*. 2017;56:893-913.
- Lin N, Vutskits L, Bebawy JF, Gelb AW. Perspectives on dexmedetomidine use for neurosurgical patients. *J Neurosurg Anesthesiol*. 2019; 31:366-77.
- Jarahzadeh MH, Mirjalili M. Analgesic effect of bupivacaine-dexmedetomidine versus bupivacaine-sufentanil in spinal anesthesia during cesarean section. *Acad J Health Sci*. 2023;38:38-42.
- Bi YH, Wu JM, Zhang YZ, Zhang RQ. Effect of different doses of intrathecal dexmedetomidine as an adjuvant combined with hyperbaric ropivacaine in patients undergoing cesarean section. *Front Pharmacol*. 2020;11:342.
- Qian H, Zhang Q, Zhu P, Zhang X, Tian L, Feng J, et al. Ultrasound-guided transversus abdominis plane block using ropivacaine and dexmedetomidine in patients undergoing caesarian sections to relieve postoperative analgesia: a randomized controlled clinical trial. *Exp Ther Med*. 2020;20:1163-8.
- Bhardwaj S, Devgan S, Sood D, Katyal S. Comparison of local wound infiltration with ropivacaine alone or ropivacaine plus dexmedetomidine for postoperative pain relief after lower segment cesarean section. *Anesth Essays Res*. 2017;11:940-5.
- Kong D, Bai J, Ma S, Li C, Yang L, Kong X. Effects of dexmedetomidine hydrochloride on hemodynamics, postoperative analgesia and cognition in cesarean section. *Exp Ther Med*. 2018;16:1778-83.
- Mo Y, Qiu S. Effects of dexmedetomidine in reducing post-cesarean adverse reactions. *Exp Ther Med*. 2017;14:2036-9.
- Tang Y, Yang M, Fu F, Huang X, Feng Y, Chen X. Comparison of the ED50 of intrathecal hyperbaric ropivacaine co-administered with or without intrathecal dexmedetomidine for cesarean section: a prospective, double-blinded, randomized dose-response trial using up-down sequential allocation method. *J Clin Anesth*. 2020;62:109725.
- Zhao Y, Xin Y, Liu Y, Yi X, Liu Y. Effect of epidural dexmedetomidine combined with ropivacaine in labor analgesia: a randomized double-blinded controlled study. *Clin J Pain*. 2017;33:319-24.

Integrated pulmonary index in pediatric sedation for endoscopy: a prospective cohort study

Índice pulmonar integrado en sedación pediátrica para endoscopia: estudio de cohorte prospectivo

Ruslan Abdullayev^{1*}, Yavuz Kelleci², Adem Halis³, Sevket Girgin⁴, Ayten Saracoglu^{5,6}, and Tumay Umuroglu¹

¹Department of Anesthesiology and Reanimation, Marmara University School of Medicine; ²Department of Anesthesiology and Reanimation, Marmara University Training and Research Hospital; ³Department of Anesthesiology and Reanimation, Koç University Hospital; ⁴Department of Public Health, Marmara University School of Medicine. Istanbul, Turkey; ⁵Department of Anesthesiology, ICU and Perioperative Medicine, Hamad Medical Corporation; ⁶Department of Anesthesiology, Qatar University College of Medicine. Doha, Qatar

Abstract

Objective: Integrated pulmonary index (IPI™) is a device working with fuzzy logic principle that analyzes patient's end-tidal carbon dioxide (ETCO₂), respiratory rate (RR), peripheral oxygen saturation (SpO₂), and pulse rate and provides a number between 1 and 10. We aimed to investigate the usefulness of IPI monitor in pediatric patients. **Methods:** After the Investigational Review Board approval pediatric patients undergoing gastrointestinal endoscopy under sedation were recruited. Propofol (Group P) and ketamine (Group K) were used for sedation. The primary outcome measure was average periprocedural IPI values. Secondary outcome measures were recovery time, endoscopist, and anesthetist satisfactions. Correlation of IPI values with physiological parameters was examined as well. **Results:** Periprocedural IPI scores were comparable between the groups (6.3 [4.9-7] vs. 6.8 [5.3-7.6] in Group P and Group K, respectively, $p = 0.153$). Recovery time was significantly longer in Group K ($p < 0.001$). Endoscopist and anesthetist satisfaction scores were comparable. Low IPI scores were significantly associated with low ETCO₂, RR, and SpO₂ values ($p < 0.001$). **Conclusions:** IPI monitor is a valuable tool in the monitorization of the pediatric patients undergoing sedation with propofol and ketamine. Both drugs are associated with comparable IPI scores. ETCO₂, RR, and SpO₂ values are measured lower in patients with low IPI scores (1-3 points).

Keywords: Gastrointestinal. IPI. Ketamine. Non-operating room. Propofol.

Resumen

Objetivo: El índice pulmonar integrado (IPI™) es un dispositivo que funciona con el principio de lógica difusa para analizar el dióxido de carbono al final de la espiración (ETCO₂), la frecuencia respiratoria (FR), la saturación periférica de oxígeno (SpO₂) y la frecuencia cardíaca (FC), y proporciona un número entre 1 y 10. Nuestro objetivo fue investigar la utilidad del monitor IPI en pacientes pediátricos. **Métodos:** Tras la aprobación por la Junta de Revisión de Investigaciones, se reclutaron pacientes pediátricos sometidos a endoscopia gastrointestinal bajo sedación. Para la sedación se utilizó propofol (grupo P) o ketamina (grupo K). La medida de resultado primaria fueron los valores medios del IPI periprocedimiento. Las medidas de resultado secundarias fueron el tiempo de recuperación y la satisfacción del endoscopista y del anestesista. También se examinó la correlación de los valores del IPI con los parámetros fisiológicos. **Resultados:** Las puntuaciones del IPI periprocedimiento fueron comparables entre los grupos (6.3 [4.9-7] en el grupo P frente a 6.8 [5.3-7.6] en el grupo K; $p = 0.153$). El tiempo de recuperación fue significativamente mayor en el grupo K ($p < 0.001$). Las puntuaciones de satisfacción del endoscopista y del

*Correspondence:

Ruslan Abdullayev
E-mail: ruslan_jnr@hotmail.com

Date of reception: 25-11-2024
Date of acceptance: 15-04-2025
DOI: 10.24875/CIRUE.M25000905

Cir Cir (Eng). 2025;93(6):660-668

Contents available at PubMed

www.cirugiaycirujanos.com

2444-0507/© 2025 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

anestésista fueron comparables. Unas puntuaciones bajas del IPI se asociaron significativamente con valores bajos de $ETCO_2$, FR y SpO_2 ($p < 0.001$). **Conclusiones:** El monitor IPI es una herramienta valiosa para la monitorización de los pacientes pediátricos sometidos a sedación con propofol o ketamina. Ambos fármacos se asocian a puntuaciones de IPI comparables. Los valores de $ETCO_2$, FR y SpO_2 son más bajos en pacientes con puntuaciones de IPI bajas (1-3 puntos).

Palabras clave: Gastrointestinal. IPI. Ketamina. Sala no quirúrgica. Propofol.

Introduction

Respiratory depression and airway obstruction are the main causes of morbidity during sedation. Although pulse oximetry is a part of routine monitoring, it has disadvantages such as delayed display of arterial deoxygenation and lack of information about ventilation¹. Capnography was found to be superior to pulse oximetry in detecting hypoventilation and airway obstruction, and the use of capnography decreased hypoxemia and apnea episodes during sedation²⁻⁴. Current clinical practice guidelines suggest that capnography should be used routinely, especially in moderate and deep sedation where verbal communication with patients cannot be maintained⁵⁻⁷.

The integrated pulmonary index (IPI™) is a commercially available tool developed to detect changes in respiratory status of patients during sedation. This software works with a principle using the fuzzy logic mathematical model algorithm⁸. It analyzes patients' physiological parameters such as end-tidal carbon dioxide ($ETCO_2$), respiratory rate (RR), peripheral oxygen saturation (SpO_2), and pulse rate (PR) together and displays a number between 1 and 10 on the monitor (Supplementary data table 1). Studies have shown that there is agreement between IPI values and clinical assessment of the patient's respiratory status⁹. Effects of many sedative drugs individually on IPI have still not been studied.

Propofol and ketamine are among the agents frequently used in endoscopy procedures. While propofol provides rapid recovery with its short half-life, it can cause severe hemodynamic instability and respiratory failure. On the other hand, ketamine causes less hemodynamic instability and less respiratory failure, but it may cause increased secretion, agitation on awakening, and prolonged recovery¹⁰.

The aim of this study was to investigate the periprocedural IPI values in pediatric patients sedated with propofol and ketamine. The primary outcome measure was average periprocedural IPI values. The secondary outcome measures were recovery time, endoscopist, and anesthetist satisfactions. The hypothesis of the study was that the patients sedated with ketamine would have higher IPI values.

Methods

This study was approved by the Marmara University Faculty of Medicine Clinical Research Ethics Committee (Date: May 07, 2021/No: 09.2021.511) and recorded on the public website (<https://clinicaltrials.gov/>, No: NCT05137574). Pediatric patients undergoing elective upper and/or lower gastrointestinal endoscopy under sedation at our hospital between May 2021 and November 2021 were included in the study. Signed written informed consent was obtained from the patients' first-degree relatives or guardians. Inclusion criteria for the study were as follows: being older than 1 month and younger than 18 years of age and having American Society of Anesthesiologists (ASA) I-III physical status. Exclusion criteria were as follows: refusal to participate in the study, having an ASA IV or higher physical status, patients unsuitable for non-operating room anesthesia, and patients with a known allergy to drugs to be used for sedation. Anaphylactic reaction that may develop due to the drugs applied; respiratory failure requiring endotracheal intubation; hemodynamic failure; and gastrointestinal perforation, bleeding, or other major complications of endoscopy were determined as exclusion criteria from the study after the enrollment.

Patients were divided into two groups where propofol was used for sedation for the first group (Group P) and ketamine for the second (Group K). Patients were blind to the study design and did not know which group they would be in. The anesthetist administering the drugs and taking the measurements was the same and was not blinded to the study. The pediatric gastroenterologist performing the endoscopy procedure was blind to the study.

Routine monitoring, including pulse oximetry (SpO_2), three-lead electrocardiography, and non-invasive blood pressure, was performed after the patient entered the procedure room. A balanced crystalloid solution was started as an intravenous infusion at a rate of 10 mL/kg/h after a peripheral vascular access was placed. Midazolam 0.05 mg/kg intravenously (IV) was administered for sedation and anxiolysis. In addition,

a special nasal cannula (Smart CapnoLine Plus; Oridion Medical, Needham, MA, USA) for IPI was applied to the patient and connected to the monitor (Capnostream 20; Oridion Medical, Needham, MA, USA). Oxygen was administered at 2 L/min through the nasal cannula of the monitor. After the preparation of the surgical team was confirmed, remifentanyl 0.5 µg/kg IV was administered. The patient was placed in the left lateral decubitus position and a mouthpiece was attached to the patient's mouth if upper endoscopy was to be performed. Then, propofol 0.5 mg/kg IV for Group P and ketamine 0.5 mg/kg IV for Group K was administered. The patient with a Ramsey Sedation Scale score of 5 or 6 (poor or no response to glabellar tapping) was handed over to the endoscopy team to perform the procedure. Additional doses of remifentanyl 0.5 µg/kg IV, propofol 0.5 mg/kg IV (for Group P), and ketamine 0.5 mg/kg IV (for Group K) were added at the discretion of the anesthetist when needed.

Data were collected by the anesthetist performing the sedation procedure. Demographic and anthropometric measurements of the patient, ASA physical status, amount of drugs used, procedure, and recovery times were recorded. ETCO_2 , RR, SpO_2 , and PR values were recorded at 3-min intervals starting from the anesthetic induction of the patient. Our primary outcome measure was mean IPI values. The secondary outcome measures were recovery time, endoscopist and anesthetist satisfactions. The endoscopist and anesthesiologist satisfactions were graded according to the Likert scale between 1 and 10 points. Jaw-thrust maneuvers and number of bag-valve-mask assisted ventilations applied to the patient during the procedure were also recorded. Finally, the correlation of IPI values with ETCO_2 , RR, SpO_2 , and PR was analyzed. Individual IPI values were grouped as low IPI (IPI 1-3), medium IPI (IPI 4-6), and high IPI (IPI 7-10) and ETCO_2 , RR, SpO_2 , and PR values were compared between these groups.

Jamovi software (Version 2.3, <https://www.jamovi.org>) was used for statistical analysis. Categorical data were defined by number and percentage. Continuous data were defined as mean and standard deviation, or median and interquartile range for normal and non-normal distributions, respectively. Shapiro-Wilk test and histogram graphs were used for normality testing. χ^2 test was used for the analysis of categorical data. Student's *t* and Mann-Whitney U-tests were used for the analysis of continuous data with normal and non-normal distributions, respectively. In the analysis of the continuous data regarding ETCO_2 , RR, SpO_2 , and

PR according to the IPI score categories of the participants, Kruskal-Wallis test was used with Dwass-Steel-Critchlow-Fligner (DSCF) test for the pairwise comparisons. The independent variables of the study were age, gender, height, body weight, duration of procedure, midazolam dose, remifentanyl dose, drug group, ASA score, type of intervention, and number of bag-valve-mask maneuvers. The dependent variables were ETCO_2 , RR, SpO_2 , PR, IPI scores, and IPI score categories. G*Power program (version 3.1) was used for power analysis of our study. Regarding the IPI score, which is the primary outcome measure of the study, with the α error of 5%, the power of the study being 80%, and the effect size of 0.75 obtained from the preliminary measurements on eight patients, the sample size was calculated to be 58.

Results

A total of 66 patients were eligible for the study, of which 60 were included in the final analysis. The flow diagram of the study is presented in figure 1. Thirty-six (60%) of the patients were female and the mean age was 13 (25th-75th percentiles: 9-15) years. Demographic and clinical characteristics of the patients are demonstrated in table 1. There was no significant difference between the two groups in terms of age, gender, height, body weight, ASA, type of intervention, amount of midazolam, and remifentanyl used. There was no significant difference between the two groups in terms of the number of jaw-thrust maneuvers ($p = 0.06$, Mann-Whitney U-test). However, when categorizing the maneuver, a significant difference was observed between the groups, where the number of the patients not requiring the jaw-thrust maneuver was 4 (13.3%) and 13 (43.3%) in the Groups K and P, respectively ($p = 0.023$, χ^2 test). The IPI scores were comparable between the groups (6.3 [4.9-7] vs. 6.8 [5.3-7.6] in Group P and K, respectively, $p = 0.153$, Mann-Whitney U-test). While there was no significant difference between the two groups in terms of procedure time, recovery time was significantly longer in Group K ($p < 0.001$). There was no significant difference between the two groups in terms of endoscopist and anesthetist satisfactions as well.

Descriptive statistics of IPI values and their corresponding physiological parameters are presented in figure 2 and Supplementary data table 2. Comparison of physiological parameters such as ETCO_2 , RR, SpO_2 , and PR between low, medium, and high IPI groups is given in table 2. While there was no




Figure 1. Study flow diagram.

difference between the IPI groups in terms of PR, a statistically significant difference was demonstrated between the low, medium, and high IPI groups in terms of ETCO_2 , RR, and SpO_2 ($p < 0.001$, Kruskal-Wallis test). Pairwise comparisons between the groups revealed that ETCO_2 in the low IPI group was significantly lower compared to the other two ($p < 0.001$, DSCF test); SpO_2 in the high IPI group was significantly higher compared to the other two ($p < 0.001$, DSCF test); and RR values were significantly different between all the three groups ($p < 0.001$, DSCF test).

There were no complications in the patients related to the endoscopy procedure itself. Severe hypoxemia developed in two patients in Group K, in which assisted ventilation was provided with a bag-valve-mask, and the hypoxemia resolved in a short time. No bag-valve-mask ventilation was needed in Group P.

Discussion

We investigated the effectiveness of the IPI monitoring in pediatric patients sedated with propofol and ketamine and the main result of our study was that these drugs had comparable IPI scores. There was a

good relationship between physiological parameters and IPI scores.

Cardiopulmonary complications continue to be important causes of morbidity and mortality in these procedures¹¹. Current clinical practice guidelines recommend monitoring the hemodynamic and respiratory status of patients, as well as their state of consciousness during sedation procedures. The Association of Anesthetists recommends capnography monitoring whenever verbal communication with sedated patients is interrupted⁷. Many studies have shown that the use of capnography provides additional safety in addition to pulse oximetry^{1,12,13}. Some studies have stated that derangements in respiratory pattern occurs before hypoxia develops and this can be detected by capnography^{14,15}. IPI is a technology that has been introduced for more than a decade⁸. Although it has a complicated working principle like the fuzzy logic mathematical model, the value between 1 and 10 reflected on the monitor makes it extremely easy to use. In addition, grouping these values and using a three-point scale instead of a ten-point scale (low, medium, and high IPI corresponding to 1-3, 4-6, and 7-10 points, respectively) makes the use of the

Table 1. Demographic and clinical characteristics of the patients

Parameter	Group P	Group K	p
Age (years)	14.5 (10-15)	12.0 (7-13.75)	0.089
Sex			
Female	19 (52.8)	17 (47.2)	0.598
Male	11 (45.8)	13 (54.2)	
Height (cm)	153.5 (123.5-160.75)	142.5 (120-160)	0.424
Body weight (kg)	42.5 (25.75-50.75)	39.5 (19.25-52)	0.437
BMI (kg/m ²)	19 ± 5.29	18.5 ± 6.05	0.734
ASA			
I	14 (56.0)	11 (44.0)	0.715
II	13 (44.8)	16 (55.2)	
III	3 (50)	3 (50.0)	
Procedure			
Gastrosocopy	21 (53.8)	18 (46.2)	0.066
Colonoscopy	8 (61.5)	5 (38.5)	
Gastrosocopy + colonoscopy	1 (12.5)	7 (87.5)	
Midazolam amount (mg)	2.0 (1.5-2)	2.0 (1-2)	0.187
Remifentanil amount (µg)	40.0 (21.25-90)	30.0 (20.5-50)	0.609
Propofol/Ketamine amount (mg)	104.33 ± 50.03	72.50 ± 55.20	
IPI	6.3 (4.9-7)	6.8 (5.3-7.6)	0.153
Jaw-thrust maneuver (number)	1 (1-2) (0-6)	1 (0-2) (0-5)	0.06
0	4 (23.5)	13 (76.5)	0.023*
1	13 (68.4)	6 (31.6)	
≥ 2	13 (54.2)	11 (45.8)	
Procedure duration (min)	12.0 (10-22.5)	15.5 (10-29.5)	0.229
Recovery duration (min)	15.0 (13.5-20)	28.5 (25-30)	< 0.001*
Endoscopist satisfaction (1-10-point Likert scale)	10.0 (10-10)	10.0 (10-10)	0.190
≤ 6	1	0	
7-9	0	4	
10	29	26	
Anesthetist satisfaction (1-10-point Likert scale)	10.0 (8-10)	10.0 (10-10)	0.502
≤ 6	3	4	
7-9	6	2	
10	21	24	

*p < 0.05.

Categorical variables are given as number (percentage). Continuous variables are given as median (25th-75th percentiles) or mean ± standard deviation. For jaw-thrust maneuvers minimum-maximum values are given as well.

Group P: propofol used group; Group K: ketamine used group; BMI: body mass index; ASA: American Society of Anesthesiologist physical status; IPI: integrated pulmonary index.

Table 2. Distribution of physiological parameters according to low, medium, and high IPI groups

Parameter	Low IPI (1-3)	Medium IPI (4-6)	High IPI (7-10)	p
Number of measurements	71	85	257	
ETCO ₂ (mmHg)	12 ^a (0-34.5)	44 (35-49)	42 (36-47)	< 0.001*
RR (breaths/min)	2 ^b (0-10)	13 ^a (10-20)	19 ^b (14-23)	< 0.001*
SpO ₂ (%)	96 (86.5-100)	99 (97-100)	100 ^c (99-100)	< 0.001*
PR (beats/min)	104 (85.5-125)	103 (82-128)	105 (90-116)	0.886

*p < 0.05.

^aSignificant difference compared to the medium and high IPI groups (p < 0.001, Dwass-Steel-Critchlow-Fligner (DSCF) test).

^bSignificant difference between all the IPI groups (p < 0.001, DSCF test).

^cSignificant difference compared to the low and medium IPI groups (p < 0.001, DSCF test).

Variables are given as median (25th-75th percentiles).

ETCO₂: end-tidal carbon dioxide; RR: respiratory rate; SpO₂: peripheral oxygen saturation; PR: pulse rate.

beginning of the procedure, we provided “balanced sedation” by applying remifentanil and the appointed intravenous anesthetic agent. The IPI values we obtained were consistent with the values in the propofol + midazolam or propofol + midazolam + fentanyl groups in the study of Garah et al.¹¹ In their study, the threshold value for the low midazolam dose was 0.08 mg/kg. In our study, we used midazolam at a dose of 0.05 mg/kg as a standard, and our IPI values were consistent with the low midazolam group in that study.

The presence of an anesthetist during the endoscopy procedure is a comfort for the endoscopy team. Evaluating the patient separately and creating a personalized anesthetic protocol not only increases the safety of the patient but also ensures the endoscopy team to focus on the endoscopy procedure itself. The American Society for Gastrointestinal Endoscopy recommends that endoscopists apply propofol-based sedation if it will increase patient safety, comfort, or the effectiveness of the procedure. According to the American Gastroenterological Association and Canadian Association of Gastroenterology guidelines, there is no increased risk associated with propofol administration by non-anesthetists^{17,18}. Nevertheless, more than 20 European national anesthesia societies, including the Italian Society of Digestive Endoscopy and the Spanish Society of Anesthesia, have recommended the administration of propofol to be limited only to those trained in general anesthesia practice¹⁹⁻²¹. When deciding whether to provide the sedation by the endoscopy team, either by nurses or physicians, or by the anesthesia team, factors such as the patient’s

monitor even more practical. The use of the monitor in this way is more advantageous in cases where sedation is performed by the endoscopist or a nurse^{11,16}.

In our study, the mean IPI values were 6.3 in the propofol group and 6.8 in the ketamine group. Garah et al.¹¹ demonstrated that factors such as young age, use of propofol alone, high midazolam dose, and presence of an anesthesiologist were associated with low IPI values. In our study, we initially administered a standard dose of midazolam to the patients to benefit from its anxiolytic and amnesic properties. At the




Figure 2. Distribution of physiological variables with different IPI values. ETCO₂: end-tidal carbon dioxide; RR: respiratory rate; SpO₂: peripheral oxygen saturation; PR: pulse rate. Please note that IPI is determined by the combination of all the four parameters using the fuzzy logic principle.

ASA physical status, comorbidities, the possibility of difficult airway or intubation, the types of drugs used, and probably less important factors such as the cost of the procedure should be considered altogether²². In our clinic pediatric endoscopy, procedures are routinely performed under sedation provided by the anesthesia team. We think that the endoscopist satisfaction scores in our study also reflect this practice.

In case of hypoxemia during the procedure nasal oxygen was increased, jaw-thrust maneuver was performed, and ventilation with bag-valve-mask was used if needed. No endotracheal intubation was needed in our study. The fact that the patient has an anesthesia team that only deals with sedation and the implementation of a protocol in accordance with the concept of balanced anesthesia played an important role here. While only two patients in the ketamine group were ventilated with a bag-valve-mask due to deep hypoxia, none of the patients in the propofol group required bag-valve-mask ventilation. In our study, the jaw-thrust maneuver was less in the ketamine group. Propofol, ketamine, dexmedetomidine, etomidate, midazolam, pentobarbital, and various combinations of these drugs are used for pediatric gastrointestinal endoscopy procedures. Each drug has its own advantages and disadvantages, and this should be considered. Propofol, especially when given as a bolus, may cause respiratory depression, apnea, and hemodynamic instability^{23,24}. In a large review that examined the safety profile of propofol and included 17,066 patients, 9.3% of patients had desaturation, 1.9% had apnea, and 1.4% needed assisted ventilation²⁵. Ketamine has advantages such as having analgesic effect, less respiratory depression, and not causing hemodynamic instability. Disadvantages include nausea, unpleasant recovery reactions, and rarely apnea or laryngospasm. Hallucinations usually occur in children over the age of 15. Apnea and laryngospasm occur in < 1% of patients^{26,27}. Although ketamine generally preserves airway muscle tone and protective reflexes and does not suppress spontaneous breathing, desaturation is sometimes inevitable in the case of laryngospasm. Our findings were consistent with the literature.

Satisfaction scores of the team providing the sedation were also measured. The evaluation was made on parameters such as the patient's immobility, maintenance of spontaneous breathing, analgesia, no need for jaw-thrust, or assisted ventilation. There was no significant difference between the two groups in terms of anesthetist satisfaction. Even though ketamine, considering the above-mentioned characteristics, seems

to be more advantageous, it is not surprising that the anesthetist who knows the pros and cons of both drugs achieves similar satisfaction scores with both drugs.

In our study, a statistically significant difference was found between low, medium, and high IPI groups in terms of $ETCO_2$, RR, and SpO_2 . In the low IPI group, $ETCO_2$, RR, and SpO_2 values were below the physiological limits. This was an expected result when we consider the principle of the IPI's fuzzy logic model methodology. The expected effect of propofol is that the RR and subsequently SpO_2 decrease. Ketamine can lower SpO_2 for the reasons mentioned above. Respiratory depression has various effects on $ETCO_2$. Some studies have shown that there is an increase in $ETCO_2$ values with sedation^{28,29}. However, insufficient CO_2 reaching the sample line with airway obstruction and hypoventilation will be associated with a low $ETCO_2$ reading¹. It should be kept in mind that these measured CO_2 values do not reflect arterial CO_2 values and there may be significant differences between the two³⁰. Arterial blood gas analysis should be used for the latest. The low $ETCO_2$ values in the low IPI group in our study were consistent with the findings of Berkenstadt et al.¹ In addition, RR and SpO_2 values were lower in the low IPI group in our study. Although reducing the IPI score, which we evaluate over ten scales, to three scales as low-medium-high makes the interpretation easier, it may deprive us of some valuable physiological parameters. This must be done very carefully as it may compromise the patient's follow-up.

Our study had few limitations. First, the same anesthetist provided sedation to the patient and measured the IPI scores. Blinding of this part of the study would have resulted in more objective approach to the IPI monitorization with two different drugs. Second limitation of our study was that we did not measure the level of sedation objectively. We do not routinely measure the sedation-anesthesia depth for non-operating room procedures, but in such a study, sedation depth monitoring with processed electroencephalography (EEG) could have been important in terms of the group homogeneity; yet keeping in mind that the processed EEG monitor could be unreliable in pediatric patients. Thirdly, we included both upper endoscopy and colonoscopies in the study. Anesthetic implications of these are different, specifically as regard to the airway control for the upper endoscopy and more need for analgesia for the colonoscopy. Selecting only one of these procedures for the study would have been more solid in terms of

the group homogeneity. Another is the age of the patients. We had a broad range of the age, and the pattern of the respiratory complications could have been dependent on the age, so it would be prudent to have a narrower age range for the patients. And finally, we did not measure the patients' pain levels objectively. Considering that ketamine has analgesic activity unlike propofol, having an objective pain scale would be valuable. Future studies may also include the measures of depth of sedation such as Bispectral Index and/or level of analgesia such as Analgesia Nociception Index.

Conclusion

IPI monitor is a valuable tool in the monitorization of the pediatric patients undergoing sedation for endoscopic procedures with propofol and ketamine. In this study, we found that both propofol and ketamine were associated with comparable IPI scores. IPI scores had good correlation with the physiological parameter routinely monitored; and ETCO_2 , RR, and SpO_2 values were measured lower in the patients with low IPI scores (1-3 points).

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of humans and animals. The authors declare that the procedures followed complied with the ethical standards of the responsible human experimentation committee and adhered to the World Medical Association and the Declaration of Helsinki. The procedures were approved by the Institutional Ethics Committee.

Confidentiality, informed consent, and ethical approval. The authors have followed their institution's confidentiality protocols, obtained informed consent from patients, and received approval from the Ethics Committee. The SAGER guidelines were followed according to the nature of the study. IRB number: Marmara University Faculty of Medicine Clinical Research Ethics Committee (Date: 07.05.2021/No: 09.2021.511)

Clinical trial registration number: <https://clinicaltrials.gov/>, No: NCT05137574 (<https://register.clinicaltrials.gov/prs/app/action/SelectProtocol?sid=S000BK3J&selectaction=Edit&uid=U0004KE3&ts=2&cx=-d2tci9>)

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing of this manuscript.

Supplementary data

Supplementary data are available at DOI: 10.24875/CIRU.24000615. These data are provided by the corresponding author and published online for the benefit of the reader. The contents of supplementary data are the sole responsibility of the authors.

References

- Berkenstadt H, Ben-Menachem E, Herman A, Dach R. An evaluation of the integrated pulmonary index (IPI) for the detection of respiratory events in sedated patients undergoing colonoscopy. *J Clin Monit Comput.* 2012;26:177-81.
- Vargo JJ, Zuccaro G Jr., Dumot JA, Conwell DL, Morrow JB, Shay SS. Automated graphic assessment of respiratory activity is superior to pulse oximetry and visual assessment for the detection of early respiratory depression during therapeutic upper endoscopy. *Gastrointest Endosc.* 2002;55:826-31.
- Peveling-Oberhag J, Michael F, Tal A, Welsch C, Vermehren J, Farnik H, et al. Capnography monitoring of non-anesthesiologist provided sedation during percutaneous endoscopic gastrostomy placement: a prospective, controlled, randomized trial. *J Gastroenterol Hepatol.* 2020;35:401-7.
- Deitch K, Miner J, Chudnofsky CR, Dominici P, Latta D. Does end tidal CO_2 monitoring during emergency department procedural sedation and analgesia with propofol decrease the incidence of hypoxic events? A randomized, controlled trial. *Ann Emerg Med.* 2010;55:258-64.
- Krmpotic K, Rieder MJ, Rosen D. Recommendations for procedural sedation in infants, children, and adolescents. *Paediatr Child Health.* 2021;26:128-9.
- Practice guidelines for moderate procedural sedation and analgesia 2018: a report by the American society of anesthesiologists task force on moderate procedural sedation and analgesia, the American association of oral and maxillofacial surgeons, American college of radiology, American dental association, American society of dentist anesthesiologists, and society of interventional radiology. *Anesthesiology.* 2018; 128:437-79.
- Klein AA, Meek T, Allcock E, Cook TM, Mincher N, Morris C, et al. Recommendations for standards of monitoring during anaesthesia and recovery 2021: guideline from the association of anaesthetists. *Anaesthesia.* 2021;76:1212-23.
- Taft A, Ronen M, Epps C, Waugh J, Wales R. A Novel Integrated Pulmonary Index (IPI) Quantifies Heart Rate, etCO_2 , Respiratory Rate and SpO_2 . Berlin: ScienceOpen, Inc.; 2008.
- Lightdale JR, Fredette ME, Atmadja ML, Heard L, Jiang H. M1563: pilot study of the smart capnography integrated pulmonary Index™ in a pediatric gastroenterology procedure unit. *Gastrointest Endosc.* 2010; 71:AB255.
- Goudra B, Gouda G, Mohinder P. Recent developments in drugs for GI endoscopy sedation. *Dig Dis Sci.* 2020;65:2781-8.
- Garah J, Adiv OE, Rosen I, Shaoul R. The value of integrated pulmonary index (IPI) monitoring during endoscopies in children. *J Clin Monit Comput.* 2015;29:773-8.
- Waugh JB, Epps CA, Khodneva YA. Capnography enhances surveillance of respiratory events during procedural sedation: a meta-analysis. *J Clin Anesth.* 2011;23:189-96.
- Beitz A, Riphaus A, Meining A, Kronshage T, Geist C, Wagenpfeil S, et al. Capnographic monitoring reduces the incidence of arterial oxygen desaturation and hypoxemia during propofol sedation for colonoscopy: a randomized, controlled study (ColoCap Study). *Am J Gastroenterol.* 2012;107:1205-12.
- Dar AQ, Shah ZA. Anesthesia and sedation in pediatric gastrointestinal endoscopic procedures: a review. *World J Gastrointest Endosc.* 2010; 2:257-62.

15. Lightdale JR, Goldmann DA, Feldman HA, Newburg AR, DiNardo JA, Fox VL. Microstream capnography improves patient monitoring during moderate sedation: a randomized, controlled trial. *Pediatrics*. 2006;117:e1170-8.
16. Conway A, Chang K, Rad MG, Mafeld S, Parotto M. Integrated pulmonary index during nurse-administered procedural sedation: study protocol for a cluster-randomized trial. *J Adv Nurs*. 2022;78:2245-54.
17. Early DS, Lightdale JR, Vargo JJ, Acosta RD, Chandrasekhara V, Chathadi KV, et al. Guidelines for sedation and anesthesia in GI endoscopy. *Gastrointest Endosc*. 2018;87:327-37.
18. Birk J, Bath RK. Is the anesthesiologist necessary in the endoscopy suite? A review of patients, payers and safety. *Expert Rev Gastroenterol Hepatol*. 2015;9:883-5.
19. Conigliaro R, Fanti L, Manno M, Brosolo P, Italian Society of Digestive Endoscopy (SIED) Sedation Group. Italian society of digestive endoscopy (SIED) position paper on the non-anaesthesiologist administration of propofol for gastrointestinal endoscopy. *Dig Liver Dis*. 2017;49:1185-90.
20. Alvarez J, Cabadas R, De La Matta M. Patient safety under deep sedation for digestive endoscopic procedures. *Rev Esp Enferm Dig*. 2017;109:137-43.
21. Perel A. Non-anaesthesiologists should not be allowed to administer propofol for sedation: a consensus statement of 21 European national societies of anaesthesia. *Eur J Anaesthesiol*. 2011;28:580-4.
22. Sappenfield JW, White JD. Do we really need an anesthesiologist for routine colonoscopy in American society of anesthesiologist 1 and 2 patients? *Curr Opin Anaesthesiol*. 2018;31:463-8.
23. Vespasiano M, Finkelstein M, Kurachek S. Propofol sedation: intensivists' experience with 7304 cases in a children's hospital. *Pediatrics*. 2007;120:e1411-7.
24. Emrath ET, Stockwell JA, McCracken CE, Simon HK, Kamat PP. Provision of deep procedural sedation by a pediatric sedation team at a freestanding imaging center. *Pediatr Radiol*. 2014;44:1020-5.
25. Lamond DW. Review article: safety profile of propofol for paediatric procedural sedation in the emergency department. *Emerg Med Australas*. 2010;22:265-86.
26. Sahyoun C, Krauss B. Clinical implications of pharmacokinetics and pharmacodynamics of procedural sedation agents in children. *Curr Opin Pediatr*. 2012;24:225-32.
27. Melendez E, Bachur R. Serious adverse events during procedural sedation with ketamine. *Pediatr Emerg Care*. 2009;25:325-8.
28. McQuillen KK, Steele DW. Capnography during sedation/analgesia in the pediatric emergency department. *Pediatr Emerg Care*. 2000;16:401-4.
29. Hart LS, Berns SD, Houck CS, Boenning DA. The value of endtidal CO₂ monitoring when comparing three methods of conscious sedation for children undergoing painful procedures in the emergency department. *Pediatr Emerg Care*. 1997;13:189-93.
30. Stein N, Matz H, Schneeweiss A, Eckmann C, Roth-Isigkeit A, Hüppe M, et al. An evaluation of a transcutaneous and an end-tidal capnometer for noninvasive monitoring of spontaneously breathing patients. *Respir Care*. 2006;51:1162-6.

Inclusion for disaster risk management in hospitals of the Instituto Mexicano del Seguro Social

Inclusión para la gestión del riesgo de desastres en hospitales del Instituto Mexicano del Seguro Social

Hermes M. Cortés-Meza*, Sandra Elizondo-Argueta, Everardo Martínez-Valdés, Jorge L. Ponce-Hernández, Paul R. Bautista-Santos, and Felipe Cruz-Vega†

†In memoriam.

Coordinación de Proyectos Especiales en Salud, Instituto Mexicano del Seguro Social, Mexico City, Mexico

Abstract

Objective: Determine the level of inclusion in hospital disaster risk management of the Mexican Social Security Institute. **Methods:** The methodology Disability Inclusion in Hospital Disaster Risk Management (INGRID-H) of the Pan American Health Organization was adapted for self-assessment in 276 hospitals. Expert advice sessions were conducted using a cascade format, including verification visits, and 180 days between each evaluation. **Results:** In the first self-assessment, 220 hospitals submitted their reports, and in the second, 236. 186 met inclusion criteria and were selected for analysis. In the first self-assessment the mean obtained for the level of inclusion was 36.70%, and in the second was 52.03%, with a correlation of 0.410 and $p < 0.001$, with a confidence interval of 95%, being statistically significant. **Conclusions:** The adaptation of the INGRID-H tool using self-assessments and tutorials with experts made it possible to obtain a national diagnosis and increase the level of inclusion for disaster risk management in hospitals.

Keywords: Risk management. Disaster. Disability. Inclusion. Hospitals.

Resumen

Objetivo: Determinar el nivel de inclusión para la gestión de emergencias y desastres en hospitales del Instituto Mexicano del Seguro Social. **Métodos:** Se adaptó la metodología de Inclusión para la Gestión de Riesgo de Desastres en Hospitales (INGRID-H) de la Organización Panamericana de la Salud para su aplicación mediante autoevaluación en 276 hospitales. Para ello, se realizaron sesiones de asesoría con expertos mediante un formato en cascada, visitas de verificación y cumplir con el tiempo de 180 días entre cada evaluación. **Resultados:** En la primera autoevaluación, 220 hospitales enviaron sus informes, y en la segunda lo hicieron 236. Se seleccionaron 186 hospitales que cumplieron con los criterios de inclusión para el análisis de resultados. La media obtenida para el nivel de inclusión en la primera autoevaluación fue del 36.70% y en la segunda fue del 52.03%, con una correlación de 0.410 y $p < 0.001$, con un intervalo de confianza del 95%, siendo estadísticamente significativa. **Conclusiones:** La adaptación de la herramienta INGRID-H basada en autoevaluaciones y tutorías con expertos permitió obtener un diagnóstico y realizar acciones con base en ajustes razonables para incrementar el nivel de inclusión para la gestión del riesgo de desastres en los hospitales.

Palabras clave: Gestión de riesgos. Desastres. Discapacidad. Inclusión. Hospitales.

*Correspondence:

Hermes M. Cortés-Meza
E-mail: hermesmanuel@gmail.com

Date of reception: 09-05-2024

Date of acceptance: 06-10-2024

DOI: 10.24875/CIRUE.M24000906

Cir Cir (Eng). 2025;93(6):669-674

Contents available at PubMed

www.cirugiaycirujanos.com

2444-0507/© 2024 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

It is estimated that currently 1.3 billion people suffer from a significant disability, representing approximately 16% of the world's population. This number is increasing due to the growth of non-communicable diseases and the increase in people's life expectancy^{1,2}. This population represents one of the most excluded groups and is disproportionately affected by crises, emergencies, or disasters; even under normal circumstances, people with disabilities have more difficulty accessing health, education, employment, and social participation, and are two to four times more likely to die in disasters and emergencies than people without disabilities³.

In Mexico, according to the 2020 Population and Housing Census, of 126 million inhabitants in the country, 5.7% have a disability or some mental problem or condition⁴.

The 2023 Global Survey on Persons with Disabilities and Disasters report by the United Nations Office for Disaster Risk Reduction² reveals limited progress in disability inclusion over the past 10 years. Disasters affect everyone, but they have a disproportionate impact on people with disabilities, who continue to experience barriers to participation and social exclusion, making it essential to consolidate actions and efforts at all levels, and even more so from health services, so that in times of emergencies and disasters, in accordance with the "Safe Hospitals against Disasters" strategy⁵, the premise of "leaving no one behind" is met. This paper presents the experience of the Mexican Social Security Institute (IMSS) in adapting for the implementation of the methodology of "Inclusion for Disaster Risk Management in Hospitals" (INGRID-H) from the Pan American Health Organization (PAHO) and the World Health Organization (WHO) as part of the actions being developed within the framework of the "Institutional Policy for Health Care of Vulnerable Groups, with a Human Rights Approach and Gender Perspective"⁶.

The INGRID-H methodology promotes inclusion based on addressing the needs of people with disabilities working in the health facility, is aimed at all hospital staff, and emphasizes those who work in emergency and disaster risk management. The aspects it evaluates are the visibility of people with disabilities, the participation of people with disabilities, universal accessibility, capacities developed for disaster response, and the hospital emergency and

disaster response plan. Once the hospital is evaluated, it obtains a score that categorizes it as exclusionary, probably inclusive, or inclusive, and actions are recommended for implementation to increase the level of inclusion^{7,8}.

Methods

With the support of the WHO/PAHO Collaborating Center Mex-35 for resilient health services and with the aim of standardizing interventions and advice, a team of facilitators in the INGRID-H methodology was formed for health personnel with training and experience in health service management and accredited as evaluators of Mexico's "Safe Hospital Program." These health professionals joined the task teams both in disseminating information and providing advice regarding the INGRID-H methodology and in subsequent verification visits throughout the country.

The need to adapt the INGRID-H methodology so that it could be applied in a self-assessment format rather than an evaluation by facilitators was observed, in order to achieve national deployment and widespread dissemination of this activity, and to have information from IMSS hospitals within defined timeframes (180 days between each evaluation). The universe of hospitals was 276, including second and third-level units, and a goal of 80% submission compliance for self-assessments was established.

Advisory phase

From July to September 2022, four in-person training workshops were held for each region of the country (IMSS medical units at all three levels of care are organized for practical purposes into four geographical regions in the Mexican Republic: central, south-east, west, and north), with the participation of a total of 230 health professionals assigned to different hospitals. The workshops included conceptual aspects of disability, inclusion, methodology, and the use of the INGRID-H tool.

During October and November 2022, 404 national hospital directors were trained through regional virtual sessions on how to conduct self-assessments using the INGRID-H methodology, during which commitments were established:

- The governing body had to establish its task team for the self-assessment.
- The dates and method for submitting self-assessment forms along with graphic evidence.

Self-assessment phase

The initial self-assessment and the 180-day self-assessment (according to the cycle specified by INGRID-H) were received via email, and a shared folder was enabled in the institutional cloud to receive evidence of the actions implemented in each hospital, in accordance with the basic INGRID-H action plan.

The units submitted the basic action plan and the evaluation report with the INGRID-H tool in electronic format.

To facilitate monitoring of progress and evaluation results, a Microsoft SharePoint web portal for internal communication and a dashboard in Microsoft Power BI were developed, which allowed for a higher degree of compliance with the commitments established for national deployment.

Verification and advisory phase

When the facilitator team decided to adapt the INGRID-H methodology to a self-assessment format, it was considered that there could be a bias in the information, both due to the risk of unfamiliarity with concepts and due to underestimation or overestimation in the way of self-assessment, and for this reason, the need to implement verification visits and validate the information was established. During these visits, the self-assessment information, the hospital emergency and disaster plan, and the actions implemented in the hospital were reviewed and validated, and a tour of the facilities was also conducted, and personnel were interviewed. As an added value, the hospital session was scheduled to take place on the day of the visit so that the facilitator team could present the topic "For a Culture of Inclusion" to the staff.

Results

For the national deployment of INGRID-H, a total of 276 hospital medical units were included: 240 second-level care, 33 third-level care, and 3 outpatient care.

In the first self-assessment, 220 reports were received, representing 80% of the total units included, and for the second self-assessment, 236 reports were received, representing 86% of the total units.

The first self-assessment is considered the unit's initial diagnosis, and the second self-assessment allows for tracking and implemented actions to increase its level of inclusion. According to the

methodology established by PAHO/WHO, the cycle should be completed in 180 days, but in reality, the timeframes for collecting information, validating information, conducting verification visits, analyzing results, and monitoring action plans cause processes to take longer to conclude, sometimes requiring periods between each cycle longer than defined.

When comparing the self-assessments and for the analysis of the results, 90 units were excluded, either because they did not submit one of the self-assessments or because the requested information was incomplete; therefore, the comparative analysis of results was performed on 186 hospitals. The classification of hospitals according to their overall score is shown in figure 1.

In the first self-assessment, 125 hospitals were classified as exclusionary, 45 as probably inclusive, and 16 as inclusive. In the second self-assessment, 67 hospitals were classified as exclusionary, 77 as probably inclusive, and 42 as inclusive. We observed a 46% reduction in hospitals classified as exclusionary, as well as a 70% increase in hospitals classified as probably inclusive and a 163% increase in the number of units classified as inclusive. The minimum score obtained in the first self-assessment was 1.86% and the maximum was 93.76%; in the second self-assessment, the minimum score was 8.94% and the maximum was 98.04% (Fig. 2).

For each of the aspects evaluated by INGRID-H, a comparative analysis of the results was carried out and three compliance ranges were established: 0% to < 44.9%, $\geq 44.9\%$ to < 72.5% and $\geq 72.5\%$ to 100%, according to the INGRID-H index scale (Fig. 3). For statistical analysis, the t-test for related samples was applied, using *SPSS software* version 25.0. It was found that the mean obtained in the first self-assessment was 36.70% and in the second self-assessment it was 52.03%, with a correlation of 0.410 and a p value < 0.001, with a 95% confidence interval, so the difference between the overall results is statistically significant (Tables 1 and 2).

Discussion

The midterm review of the Sendai Framework⁹ emphasizes the importance of adopting a whole-of-society approach to encompass all of society for the implementation of inclusive disaster risk reduction, and underscores the urgency of involving all segments of society, including people with disabilities who are often overlooked, in disaster




Figura 1. Classification of hospitals according to their level of inclusion.




Figura 2. Overall results of the two self-assessments.

preparedness, response, and recovery efforts. People with disabilities are not visualized before, during, or after an emergency, or even more so, they are not integrated, consulted, or represented in emergency management, and therefore we do not know what their needs are¹⁰.

Published works on accessibility and inclusion have focused on the resources and perceived barriers by users and managers of health facilities with a focus on access to health services. In this regard, Villarreal et al.¹¹ found limited accessibility at the second and first levels of care in the IMSS, considering the barriers that users have to overcome to access health services and the available resources. There are also extensive reports of accessibility evaluations with a focus on physical barriers, the chain of accessibility, and communication, such as the one published by the Coordinating Federation of People with Physical

Tabla 1. Statistics of paired samples

Aspect to compare	Mean	n	Std	Mean error deviation
Pair 1 INGRID-H 1	36.70%	186	20.98%	1.54%
INGRID-H 2	52.03%	186	22.99%	1.69%
Pair 2 Visibility 1	32.17%	186	31.66%	2.32%
Visibility 2	52.70%	186	32.28%	2.37%
Pair 3 Representation 1	17.20%	186	34.35%	2.52%
Representation 2	39.43%	186	44.23%	3.24%
Pair 4 Accessibility 1	47.33%	186	19.49%	1.43%
Accessibility 2	55.50%	186	18.88%	1.38%
Pair 5 Capabilities	36.52%	186	35.31%	2.59%
Capabilities 2	47.99%	186	34.43%	2.52%
Pair 6 Hospital plan	48.31%	186	32.01%	2.35%
Hospital plan 2	61.83%	186	29.45%	2.16%

Tabla 2. Correlations of paired samples

Aspect to compare	n	Correlation	p
Pair 1 INGRID-H 1 and INGRID-H 2	186	0.41	0.000
Pair 2 Visibility 1 and visibility 2	186	0.293	0.000
Pair 3 Representation 1 and representation 2	186	0.235	0.001
Pair 4 Accessibility 1 and accessibility 2	186	0.351	0.000
Pair 5 Capabilities 1 and capabilities 2	186	0.306	0.000
Pair 6 Hospital plan 1 and hospital plan 2	186	0.285	0.000

Disabilities of Gipuzkoa¹², which undoubtedly contribute to guaranteeing access to services and represent a solid basis for inclusion in disaster risk management. Gutierrez et al.¹³ validated an inclusion assessment scale for centers and services for people with disabilities, in order to help organizations self-assess their status regarding inclusive values and processes, and thus facilitate the development of inclusive cultures, policies, and practices.

The adaptation of the PAHO/WHO INGRID-H methodology into a self-assessment format allowed for a large-scale initial diagnosis to understand our current position and define, according to the action plan, specific activities to strengthen inclusion. A total of 186 units submitted their information in the first self-assessment, which allowed us to observe that we do




Figure 3. Comparison of results obtained by hospitals in each aspect of INGRID-H.

not know people with disabilities, much less involve them in risk management. Considering that the visibility and participation emphasized by INGRID-H are based on the personnel working in the workplaces, it allowed the medical units to generate actions from the governance to include them in their committees and plans. The WHO's Guidance Note on Disability Risk Management and Health Emergencies states that before an emergency, we must identify pre-existing vulnerabilities, resources, and service gaps for people with disabilities, as well as opportunities to reinforce and strengthen capacities. When comparing these items in the second self-assessment, we observed that in both visibility and participation, the compliance percentage increased in the units¹⁴.

An example of the need to continue promoting the inclusion of people with disabilities is found in the publication by Qi and Hu¹⁵, which refers to the fact that COVID-19 emergency declarations in China lacked an inclusion perspective by overlooking the needs of people with disabilities, which shows that in the face of a serious public health crisis, and with knowledge of this right, they continue to be left aside. Inclusive communication from hospitals is feasible, and we could observe this from the impulse given to specific actions in accessibility to strengthen the autonomy of people with disabilities; an example of this was the adaptation of signage in evacuation routes, and during verification visits, it was observed that in some units, health personnel took Mexican sign language courses.

The integration of people with disabilities in disaster risk management allows them to provide us with the

necessary information so that their specific capabilities and needs are integrated into the plans. An example of how plans can be inclusive is the 2022 National Fire Protection Association (NFPA) document, Emergency Evacuation Plan, Guide for People with Disabilities¹⁶, which provides guidance for developing an emergency evacuation plan that includes people with disabilities.

Upon reviewing the results obtained in the self-assessments, hospital response plans, although they considered people with disabilities, did not actually comply with what is reviewed in INGRID-H, so emphasis was placed on how to integrate the inclusion variable, but above all with the vision that it is directed towards the medical unit worker, in addition to how to strengthen their capacities and the importance of conducting inclusive simulation exercises and drills.

As mentioned, the INGRID-H methodology is based on disaster risk management from the inclusion of people with disabilities. It is necessary to continue strengthening inclusion from a human rights perspective in which universal access to health services becomes a reality, and why not, visualize that through this strategy, interventions can also be promoted to eliminate any type of cultural, educational, economic, political barriers, etc.,^{17,18}.

Conclusions

The INGRID-H methodology focuses on hospital workers with disabilities and encourages them to participate in disaster risk management. The adaptation and implementation of INGRID-H through self-assessment from the strategic level, with a

mentoring-based methodology with experts directed to the tactical and operational levels in a cascading manner, allowed for greater reach in less time and a large-scale diagnosis to focus resources and efforts with the aim of increasing the level of inclusion in hospitals facing emergencies and disasters, but above all, to fulfill the premises of “leaving no one behind” and “nothing about us without us”¹⁹.

Inclusion in disaster risk management is one of the pillars of hospital and health sector resilience, and it is through the institutionalization of these strategies that a greater impact can be achieved. The experience gained with this project sets the stage for further progress in the culture of inclusion and respect for the rights of people with disabilities.

Acknowledgments

The authors thank Daniela Becerril, Alejandra Cortés, Mariela Santiago, and Dora Bernal for all the support provided for the development of this project.

Funding

The authors declare that they have not received funding for this study.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Ethical considerations

Protection of people and animals. The authors state that no experiments on humans or animals were performed for this research.

Confidentiality, informed consent, and ethical approval. The study does not involve personal patient data. The recommendations of the SAGER guidelines have been followed, according to the nature of the study.

Declaration on the use of artificial intelligence. The authors declare that they did not use any type of generative artificial intelligence for the writing of this manuscript.

References

1. Organización Mundial de la Salud. Discapacidad. OMS; 2023. (Consultado el 09-05-2024.) Disponible en: <https://www.who.int/es/news-room/fact-sheets/detail/disability-and-health>.

2. United Nations Office for Disaster Risk Reduction. 2023 Global Survey Report on Persons with Disabilities and Disasters. Geneva, Switzerland: United Nations Office for Disaster Risk Reduction; 2023. (Consultado el 09-05-2024.) Disponible en: <https://www.undrr.org/report/2023-global-survey-report-on-persons-with-disabilities-and-disasters>.
3. Organización Panamericana de la Salud. Discapacidad. (Consultado el 09-05-2024.) Disponible en: <https://www.paho.org/es/temas/discapacidad>.
4. Instituto Nacional de Estadística y Geografía. Estadísticas a propósito del día internacional de las personas con discapacidad (datos nacionales). México: INEGI; 3 de diciembre 2021. Comunicado de prensa 713/21. (Consultado el 09-05-2024.) Disponible en: https://www.inegi.org.mx/contenidos/saladeprensa/aproposito/2021/EAP_PersDiscap21.pdf.
5. Organización Panamericana de la Salud. Hospitales seguros. (Consultado el 09-05-2024.) Disponible en: <https://www.paho.org/es/emergencias-salud/hospitales-seguros>.
6. Instituto Mexicano del Seguro Social. Política Institucional para la Atención de la Salud de Grupos en Situación de Vulnerabilidad, con Enfoque de Derechos Humanos y Perspectiva de Género. México: IMSS; 2023. (Consultado el 09-05-2024.) Disponible en: <http://www.imss.gob.mx/sites/all/statics/profesionalesSalud/investigacionSalud/historico/programas/01-politica-institucional-gsv.pdf>.
7. Organización Panamericana de la Salud. INGRID-H: Inclusión para la Gestión del Riesgo de Desastres en Hospitales. Una metodología de “evaluación-acción” para mejorar la inclusión de las personas con discapacidad. Suplemento sobre la reducción del riesgo de desastres para la salud. PAHO/WHO Emergencias News; octubre 2020. (Consultado el 09-05-2024.) Disponible en: <https://www3.paho.org/disasters/newsletter/626-ingrid-h-inclusion-para-la-gestion-del-riesgo-de-desastres-en-hospitales-una-metodologia-de-evaluacion-accion-para-mejorar-la-inclusion-de-las-personas-con-discapacidad-309-101-es.html>.
8. Organización Panamericana de la Salud. Inclusión para la gestión del riesgo de desastres en hospitales. Washington, D.C.: OPS; 2018. (Consultado el 09-05-2024.) Disponible en: <https://iris.paho.org/handle/10665.2/51059>.
9. United Nations Office for Disaster Risk Reduction. The Report of the Midterm Review of the Implementation of the Sendai Framework for Disaster Risk Reduction 2015-2030. UNDRR; 2023. (Consultado el 09-05-2024.) Disponible en: <https://www.undrr.org/publication/report-midterm-review-implementation-sendai-framework-disaster-risk-reduction-2015-2030>.
10. World Health Organization. Guidance note on disability and emergency risk management for health. Geneva: WHO; 2013. (Consultado el 09-05-2024.) Disponible en: https://iris.who.int/bitstream/handle/10665/90369/9789241506243_eng.pdf?sequence=1.
11. Villarreal E, González J, Salinas AM, Garza ME, Núñez G, Uribe MG. Evaluación de la accesibilidad en los servicios de salud. Aten Primaria. 2000;25:475-8.
12. Elkartu, Federación Coordinadora de Personas con Discapacidad Física de Gipuzkoa. Diagnóstico de accesibilidad en los hospitales de la CAPV para las personas con discapacidad. Informe extraordinario de la institución del Ararteko al Parlamento Vasco. Elkartu, Federación Coordinadora de Personas con Discapacidad Física de Gipuzkoa; 2017. (Consultado el 09-05-2024.) Disponible en: https://www.ararteko.eus/sites/default/files/2021-12/0_4346_3.pdf.
13. Gutiérrez M, Hernández R, Jenaro C. Escala de Evaluación de la Inclusión (EEI) para centros y servicios para personas con discapacidad: validación preliminar. Revista Nacional e Internacional de Educación Inclusiva. 2019;12:97-118.
14. World Health Organization. Guidance note on disability and emergency risk management for health. Geneva; 2013. (Consultado el 09-05-2024.) Disponible en: https://iris.who.int/bitstream/handle/10665/90369/9789241506243_eng.pdf?sequence=1.
15. Qi F, Hu L. Including people with disability in the COVID-19 outbreak emergency preparedness and response in China. Disability & Society. 2020;35:848-53.
16. National Fire Protection Association. Emergency Evacuation Planning Guide for People with Disabilities. November 2022. (Consultado el 09-05-2024.) Disponible en: <https://epic-fsc.com/wp-content/uploads/NFPA-Evacuation-Guide-for-People-with-Disabilities.pdf>.
17. Kusumawardoyo C, Tamtomo K. Reflections on implementing the Sendai Framework in the Asia-Pacific: beyond adding disability inclusion to disaster risk reduction. Disasters. 2022;46:857-78.
18. Twigg J, Kett M, Lovell E. Disability inclusion and disaster risk reduction: overcoming barriers to progress. London: University College London; 2018. (Consultado el 09-05-2024.) Disponible en: <https://odi.cdn.ngo/media/documents/12324.pdf>.
19. Grupo de las Naciones Unidas para el Desarrollo Sostenible. Valores universales. Ginebra, Suiza: UNSDG; 2024. (Consultado el 09-05-2024.) Disponible en: <https://unsdg.un.org/es/2030-agenda/valores-universales>

Is it possible for end-tidal carbon dioxide analysis to replace non-invasive blood pressure measurement?

Es posible que el análisis de dióxido de carbono al final de la marea sustituya a la medición no invasiva de la presión arterial?

Halide H. Şahinkaya*^{ORCID} and Gözde Gürsoy-Çirkinoğlu^{ORCID}

Anesthesiology and Reanimation Department, Izmir City Hospital, Izmir, Turkey

To the Editor,

The number and the complexity of interventions performed in the pediatric cardiac catheterization laboratory (CCL) have been increasing in recent years. Anesthesia-related cardiac arrests have been reported as 1.2–8.7 times higher in the CCL than that for events occurring under general pediatric anesthesia¹. Therefore, providing continuous hemodynamic monitoring is the most important issue in CCL.

Radiofrequency ablation (RF) has proven to be an effective method for the treatment of cardiac arrhythmias in pediatric patients and has been used frequently in CCL. The procedure is preceded by an electrophysiological study that performs a systematic analysis of arrhythmias in the basal state and by evaluating the patient's response to programmed electrical stimulation. RF ablation is performed on the abnormal pathway after the arrhythmia has been induced, following an electrophysiologic study and cardiac mapping, which are used to identify and locate the area responsible for the arrhythmia. Interaction between tissue and catheter is crucial for safety and success. Contact force (CF) refers to the contact between the catheter and tissue. Rapid cardiac pacing, or rapid atrial pacing (RAP), improves the stability of and helps better CF². Deep hypotensive periods, which begin and end abruptly secondary to tachycardia, may be observed during periods of RAP and induced tachyarrhythmia. In addition, the type of arrhythmia and cardiac reserve may cause hemodynamic variability. There is a delayed time for detecting

hypotension when only non-invasive blood pressure is monitored for follow-up. This raises the question: is standard non-invasive monitoring sufficient for these patients? Do we need invasive arterial blood pressure monitoring for all patients?

Arterial cannulation for monitoring in children has minor complications, such as localized hematoma, or major complications such as thrombosis, acute interruption of arterial blood supply resulting in permanent ischemia and infection. Therefore, invasive arterial blood pressure monitoring is preferred in selected patients for special procedures. This raises the second question: What other monitoring method can we use to monitor hypotension in pediatric cardiac RF patients?

End-tidal carbon dioxide (ETCO₂) analysis and blood pressure measurement are required according to American Society of Anesthesiologists (ASA) standards for basic anesthetic monitoring³. Continual ETCO₂ analysis is a non-invasive, real-time monitoring as a cardiac-output-related parameter and also assesses the ventilation⁴. ETCO₂ has been found to be well correlated with hemodynamic changes. When minute volume of respiration stays stable, it may be assumed that low values of ETCO₂ may be attributed to changes in cardiac output. Any level of ETCO₂ measured during cardiopulmonary resuscitation correlates with return of spontaneous circulation or survival in adults experiencing cardiac arrest⁵.

Tachyarrhythmia begins and resolves rapidly in radiofrequency ablation. This is rapidly reflected in

***Correspondence:**

Halide H. Şahinkaya

E-mail: drhhande@yahoo.com

Date of reception: 08-05-2025

Date of acceptance: 11-09-2025

DOI: 10.24875/CIRUE.M25000907

Cir Cir (Eng). 2025;93(6):675-676

Contents available at PubMed

www.cirugiyacirujanos.com

2444-0507/© 2025 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

ETCO₂, as well. Hence, continual ETCO₂ monitoring allows us to detect hypotension early by comparing the baseline value.

As a result, the answer stays here; ETCO₂ can be easily used to detect hypotension immediately without waiting for the non-invasive blood pressure measurement time, and early intervention is possible during pediatric RF ablation.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of humans and animals. The authors declare that no experiments involving humans or animals were conducted for this research.

Confidentiality, informed consent, and ethical approval. The study does not involve patient personal data nor requires ethical approval. The SAGER guidelines do not apply.

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing of this manuscript.

References

1. Odegard KC, Bergersen L, Thiagarajan R, Clark L, Shukla A, Wypij D, et al. The frequency of cardiac arrests in patients with congenital heart disease undergoing cardiac catheterization. *Anesth Analg*. 2014;118:175-82.
2. Aizer A, Cheng AV, Wu PB, Qiu JK, Barbhuiya CR, Fowler SJ, et al. Pacing mediated heart rate acceleration improves catheter stability and enhances markers for lesion delivery in human atria during atrial fibrillation ablation. *JACC Clin Electrophysiol*. 2018;4:483-90.
3. Standards for Basic Anesthetic Monitoring. Committee of Origin: Standards and Practice Parameters (Approved by the ASA House of Delegates on October 21, 1986, Last Amended October 20, 2010, Last Affirmed on December 13, 2020). Available from: <https://www.asahq.org/-/media/sites/asahq/files/public/resources/standards-guidelines/standards-for-basic-anesthetic-monitoring.pdf> [Last accessed on Jun 11, 2024].
4. Ornato JP, Garnett AR, Glauser FL. Relationship between cardiac output and the end-tidal carbon dioxide tension. *Ann Emerg Med*. 1990;19:1104-6.
5. Paiva EF, Paxton JH, O'Neil BJ. The use of end-tidal carbon dioxide (ETCO₂) measurement to guide management of cardiac arrest: a systematic review. *Resuscitation*. 2018;123:1-7.

Letter to the editor about the article “Association between complete blood count parameters and histologically proven acute appendicitis”

Carta al editor sobre el artículo “Asociación entre los parámetros del hemograma y la apendicitis aguda histológicamente comprobada”

Enes Ağırman^{1*}, Rifat Peksöz², and Sabri S. Atamanalp²

¹Department of General Surgery, Erzurum City Hospital; ²Department of General Surgery, Atatürk University Research Hospital. Erzurum, Turkey

To the Editor,

We read with great interest the article by AlHarmi R., titled “Association between complete blood count parameters and histologically proven acute appendicitis.” This publication highlights the importance of laboratory values in the diagnosis of acute appendicitis¹. Acute appendicitis is the most common surgical emergency in both pediatric and adult populations. Its diagnosis relies on clinical findings, blood parameters, and imaging modalities. While the diagnostic efficacy of imaging methods currently exceeds 90%, their utilization can be restricted by various factors. In pediatric patients, challenges include patient non-compliance, concerns regarding radiation exposure from computed tomography (CT), and the inherent difficulty for pediatric patients to articulate symptoms. In adults, pregnancy and contrast-induced nephropathy are additional considerations limiting CT use. Ultrasonography offers a non-invasive and safe alternative, but its widespread applicability is hindered by the necessity for specialized operator skills, preventing its routine use by all clinicians. Particularly for physicians practicing in rural settings, blood parameters emerge as the most readily available and crucial diagnostic tool. Among laboratory assessments, the complete blood count holds a prominent position. Its rapid turnaround time, near-universal accessibility in healthcare

facilities, and satisfactory diagnostic value render it a valuable diagnostic instrument. Consequently, studies investigating blood parameters in patients with acute appendicitis are of considerable significance. The authors’ valuable contribution will undoubtedly enrich the existing literature. However, engaging in a discussion about this study will contribute to both the current study and future investigations. We’ve shared some aspects of this study that piqued our interest and offered recommendations for the article.

First, the authors indicated that the platelet-to-lymphocyte ratio (PLR) demonstrates diagnostic utility, even if at a low value. However, the diagnostic efficacy values of PLR (AUC, sensitivity, and specificity) are insufficient for its use in diagnosis or differential diagnosis. It’s possible that other parameters not assessed in the study might have similar diagnostic efficacy to PLR.

Second, bilirubin has recently emerged as a significant laboratory parameter under investigation for determining acute appendicitis and predicting complicated appendicitis². Notably, it serves as an important indicator for detecting perforation, particularly in pediatric patients³. We recommend that the authors consider incorporating bilirubin levels into their current or prospective studies, as this could yield more impactful findings.

***Correspondence:**

Enes Ağırman

E-mail: agirman_enes@hotmail.com

2444-0507/© 2025 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 29-06-2025

Date of acceptance: 07-07-2025

DOI: 10.24875/CIRUE.M25000908

Cir Cir (Eng). 2025;93(6):677-678

Contents available at PubMed

www.cirugiaycirujanos.com

Third, the inclusion and exclusion criteria should be clearly defined. Many conditions, such as systemic diseases, hematological disorders, malignancies, and pregnancy, directly influence blood parameters⁴. Therefore, patients with these conditions should be excluded from the study. The article does not provide information on whether such patients were included.

Another point to consider is the negative appendectomy rate, which is reported in the literature to be between 10% and 15%, with some studies even reaching 30%⁵. However, in your study, this rate appears to be 0%. Could you please explain how this is accounted for? Or, if such cases existed, were these patients excluded from the study?

Finally, when describing the clinical characteristics of the study groups, including information such as duration of symptoms (days), length of hospital stay, and any complications that developed, would lead to a better understanding of the patient cohorts. This information could be presented in descriptive statistics of the study groups table

We congratulate the authors on their successful study and eagerly await their response to our letter.

Sincerely,

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of humans and animals. The authors declare that no experiments involving humans or animals were conducted for this research.

Confidentiality, informed consent, and ethical approval. The study does not involve patient personal data nor requires ethical approval. The SAGER guidelines do not apply.

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing of this manuscript.

References

1. AlHarmi R. Association between complete blood count parameters and histologically proven acute appendicitis. *Asociación entre los parámetros del hemograma y la apendicitis aguda histológicamente probada.* Cir Cir (Eng). 2025;93:28-34.
2. Peksöz R, Dişçi E, Kaya A, Ağırman E, Korkut E, Aksungur N, et al. Significance of laboratory parameters in diagnosing acute appendicitis during pregnancy. *ANZ J Surg.* 2022;92:121-7.
3. Koirala DP, Gupta AK, Dahal GR, Shrestha BM, Shrestha S, Neupane S, et al. Role of hyperbilirubinaemia as a predictor of complicated appendicitis in paediatric population. *Afr J Paediatr Surg.* 2022;19:61-4.
4. Peksöz R, Bayar B. The role of complete blood count parameters in diagnosing acute appendicitis and measuring the severity of inflammation. *Ulus. Travma Acil Cerrahi Derg.* 2021;27:654-61.
5. Jukić M, Nizeteo P, Matas J, Pogorelić Z. Trends and predictors of pediatric negative appendectomy rates: A single-centre retrospective study. *Children (Basel).* 2023;10:887.