



ÓRGANO DE DIFUSIÓN
CIENTÍFICA DE LA
ACADEMIA MEXICANA
DE CIRUGÍA

FUNDADA EN 1933

CIRUGÍA Y CIRUJANOS

Content

Original articles

- 229 Impact of COVID-19 history on anxiety and complications in elective cesarean sections**
Kubra Kalkisim, Ali Akdogan, and Davut Dohman
- 235 Impact of psoas muscle area measurement in complex open abdominal aortic aneurysm repair**
Ramon García-Alva, Yuki Yoshi Kimura-Sandoval, Luis O. Bobadilla-Rosado, Javier E. Anaya-Ayala, Ximena Rosas-Flota, Natalia Sanchez-Garrido, Hugo Laparra-Escareno, Lizeth Luna, Luis H. Arzola, Adriana Torres-Machorro, and Carlos A. Hinojosa
- 240 Predictors of postoperative neurological deficits in traumatic thoracolumbar fractures**
Alp Karaaslan
- 248 Arthroscopic treatment with absorbable bone cement for intra-articular comminuted fracture of the distal radius**
Bangxian Dou, Huiling Tian, and Xuedong Li
- 255 The effect of ethyl pyruvate administration on intestinal mucosal damage in neonatal rat models of necrotizing enterocolitis**
Tugce Merve-Orbay, Aysenur Celayir, and Hulya Yavuz
- 264 Communication as the most important non-technical skill in the operating room**
Miguel Vassallo, Kimberly G. Ordaz-Cedeño, Rhayniveth Sequera, Gabriel Rondón, and José Morao
- 269 Incidence of surgical site infection in head and neck surgery: a prospective cohort study**
Diego Rodríguez-Villar, Enrique Gea-Izquierdo, Pilar Cárdenas-Soriano, M.ª Concepción Villar-del-Campo, Manuel Durán-Poveda, and Gil Rodríguez-Caravaca
- 277 Ethical considerations in dietary supplement consumption behaviors**
İlknur Arslan-Aras and Aykut Ekiyor
- 287 Is there any effect of lycopene's preventing peritoneal adhesion formation in rats: an experimental study**
Ahmet Bilgi, Mustafa C. Terek, Gürkan Yiğittürk, Dilek Taşkıran, Orkun İlgen, İsmet Hortu, Bulent Yılmaz, Cavit Kart, Fazıl Avcı, and Mehmet Kulhan
- 295 Machine learning for COVID-19 mortality prediction: enhancing cart models with node-specific odds ratios**
Argelia Pérez-Pacheco, Alejandra E. Herrera-Suárez, Jonathan Vélez-Mata, Liza M. Magdaleno-Fourlong, Amy D. Avendaño-Carrera, Catalina Casillas-Suárez, and Adolfo Pérez-García



PERMANYER MÉXICO
www.permayer.com

Volume 94, No. 3, May-June 2026

Journal Citation Reports™ from Clarivate, 20, Medline/PubMed, EBSCO Essentials™

Impact of COVID-19 history on anxiety and complications in elective cesarean sections

Impacto del antecedente de COVID-19 en la ansiedad y las complicaciones en las cesáreas electivas

Kubra Kalkisim^{id}, Ali Akdogan^{*id}, and Davut Dohman^{id}

Department of Anesthesiology and Critical Care, Faculty of Medicine, Karadeniz Technical University, Trabzon, Turkey

Abstract

Objectives: This study aims to examine the effect of COVID-19 history on pre-operative anxiety and intraoperative complications in elective cesarean section patients. **Methods:** A total of 200 patients undergoing elective cesarean section were divided into two groups based on COVID-19 history: Group C (n = 100, with COVID-19) and Group K (n = 100, without COVID-19). Anxiety was assessed using the state-trait anxiety inventory (STAI) I and II. Vital signs were monitored peroperatively. Intraoperative complications and APGAR scores were recorded. **Results:** Pregnant women with a history of COVID-19 had significantly higher state anxiety (STAI-1) scores than those without (p = 0.017). No significant difference in intraoperative or post-operative complications was observed between the groups. **Conclusions:** Anxiety levels were higher in pregnant women undergoing elective cesarean sections, especially those with a history of COVID-19. Managing increased anxiety is essential because of its potential effects on maternal and neonatal health outcomes.

Keywords: Pre-operative anxiety. COVID-19. Elective cesarean section. Intraoperative complications.

Resumen

Objetivos: Examinar el efecto de los antecedentes de COVID-19 sobre la ansiedad preoperatoria y las complicaciones intraoperatorias en pacientes sometidas a cesárea electiva. **Métodos:** Un total de 200 pacientes sometidas a cesárea electiva se dividieron en dos grupos en función de los antecedentes de COVID-19: grupo C (n=100, con COVID-19) y grupo K (n = 100, sin COVID-19). La ansiedad se evaluó mediante el Inventario de Ansiedad Estado-Rasgo I y II. Se controlaron las constantes vitales durante la operación. Se registraron las complicaciones intraoperatorias y las puntuaciones APGAR. **Resultados:** Las embarazadas con antecedente de COVID-19 presentaron puntuaciones de ansiedad estado (STAI-1) significativamente más altas que las que no lo tenían (p = 0.017). No se observaron diferencias significativas en las complicaciones intraoperatorias y posoperatorias entre los grupos. **Conclusiones:** Los niveles de ansiedad fueron mayores en las embarazadas sometidas a cesárea electiva, especialmente en aquellas con antecedente de COVID-19. Es esencial controlar el aumento de la ansiedad debido a sus posibles efectos en los resultados de salud maternos y neonatales.

Palabras clave: Ansiedad preoperatoria. COVID-19. Cesárea electiva. Complicaciones intraoperatorias.

*Correspondence:

Ali Akdogan

E-mail: draliakdogan@yahoo.com

2444-0507/© 2025 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 13-12-2024

Date of acceptance: 23-04-2025

DOI: 10.24875/CIRUE.M25000939

Cir Cir (Eng). 2026;94(3):229-234

Contents available at PubMed

www.cirugiaycirujanos.com

Introduction

Pre-operative anxiety, characterized by tension toward anesthesia and surgery, typically initiates with the anticipation of hospitalization.¹ High anxiety levels can lead to emotional and physical stress, affecting post-operative recovery, pain, and potentially delaying same-day discharge. Anesthesiologists play a key role in reducing anxiety by establishing trust and guiding the patient through the process, from preparation to recovery.²

Pregnant women commonly experience anxiety pre-operatively due to concerns about post-operative pain, fear of not waking up from anesthesia, and worries about divulging personal information under its influence.³ Pre-operative anxiety in pregnant women undergoing cesarean section is associated with perioperative complications. The mental health of the pregnant woman is crucial as it can impact the emotional bond between mother and fetus and increase the likelihood of adverse neonatal outcomes such as intrauterine growth retardation, preterm delivery, and low birth weight.⁴

The COVID-19 pandemic, declared a global pandemic by the World Health Organization on March 11, 2020, has significantly impacted global health, with cases detected across 113 countries. The pandemic's widespread effects have heightened anxiety levels and increased uncertainties regarding risks, particularly impacting the mental health of pregnant individuals.⁵

Understanding pre-operative anxiety in pregnant women is essential for developing strategies to manage anxiety effectively and mitigate perioperative complications. This study aims to contribute to the literature by exploring the relationship between pregnancy data, demographic characteristics, COVID-19 history, and anxiety levels in pregnant women scheduled for elective cesarean section. In addition, we discuss factors that may influence anxiety levels and intraoperative complications in this population.

Methods

Our study included 200 volunteer patients aged between 20 and 45 years, classified as American Society of Anesthesiologists risk II-III, scheduled for elective cesarean section at the gynecology and obstetrics clinic between July 1, 2022, and May 1, 2023, following approval from the local ethics

committee (2022/145). The study was conducted prospectively.

Exclusion criteria: patients with a body mass index > 35 kg/m², allergies to induction agents or opioids, use of medications for chronic pain, active infections, autoimmune, cardiovascular, gastrointestinal, or neurological diseases, renal or hepatic failure, chronic respiratory pathology, malignancies (colorectal, gastric, pancreatic, esophageal, lung, gynecological, etc.), chronic systemic diseases (hypertension and diabetes mellitus), hematological disorders (leukocyte and platelet diseases), and psychiatric disorders under treatment were excluded from the study. Patients who were uncooperative or unable to undergo physical and verbal performance comparisons were also excluded from the study. New participants were recruited to replace any excluded patients to achieve the planned sample size.

All patients underwent preanesthetic examinations the day before surgery. They received verbal and written information about the study and provided informed consent. Patients with a history of COVID-19, asymptomatic (no symptoms were present during the infection period), moderate (patients who had mild symptoms that did not require hospitalization), and severe (patients who required hospitalization or intensive care for their COVID-19 infection) were classified.

Pre-operative anxiety levels were assessed using the state-trait anxiety inventory (STAI), consisting of STAI-I (state anxiety) and STAI-II (trait anxiety) tests. Patients scoring 60 and above on STAI-II were excluded. Anxiety levels were categorized as follows: scores < 30 indicating little or no anxiety, 31-49 indicating moderate anxiety, and 50 and above indicating severe anxiety.

Patients were divided into two groups based on whether they had a history of COVID-19 in the past year: Group C (with COVID-19 history, n = 100) and Group K (without COVID-19 history, n = 100).

Non-invasive measurements, including mean arterial pressure (MAP), systolic arterial pressure (SAP), diastolic arterial pressure (DAB), heart rate (HR), peripheral oxygen saturation (SpO₂), and electrocardiogram were recorded. Intravenous access was established with an 18-G catheter, and 0.9% NaCl infusion was administered at 5-10 mL/kg/h. Spinal anesthesia was performed using 12 mg bupivacaine with a 25 g spinal needle. Oxygen supplementation was provided through a transparent mask as needed.

Data collection

Vital signs (SAB, DAB, MAP, HR, and blood pressure) were recorded during ward follow-up, pre-operative waiting period, intraoperative periods (0 min, 15 min, 30 min, 45 min, 60 min, 90 min after surgical incision), and post-operative period. SpO₂ values were documented accordingly. Intraoperative complications and APGAR scores of newborns were also noted.

Following surgery, patients were transferred to the post-anesthesia care unit (PACU). Vital signs (SAB, DAB, MAP, HR, and SpO₂) upon PACU admission and discharge were recorded. Stable patients were then transferred to the obstetrics and gynecology service.

Statistical analysis

Data were analyzed using the Statistical Package for the Social Sciences 23.0. Descriptive statistics were presented as numbers/percentages for categorical variables and as mean, standard deviation, minimum, and maximum for numerical variables. To control for confounding variables such as age, gestational week, previous cesarean history, and chronic illness, multivariate regression analysis was performed. Normal distribution was assessed using the Kolmogorov-Smirnov test. Parametric tests (Student's t-test, analysis of variance) were used for normally distributed data, whereas non-parametric tests (Mann-Whitney U test, Kruskal-Wallis test) were used otherwise. Chi-square analysis was employed for categorical data. *Post hoc* analysis evaluated differences between groups. Significance level was set at $p < 0.05$.

Sample size calculation

The Open Epi 3.01 program was used to calculate the sample size. Based on previous studies, 25% of pregnant women without severe acute respiratory syndrome coronavirus 2 infection were reported to have anxiety. Assuming a 45% anxiety rate in pregnant women with COVID-19, with a type 1 error of 5% and power of 80%, the study required a total of 196 participants (98/group).

Results

The study initially involved 304 pregnant women. Of these, 61 were excluded due to emergency situations, 24 due to chronic diseases, and 19 declined to

participate, resulting in a final analysis of data from 200 patients.

Demographic characteristics, including age, weight, education level, income-generating employment, smoking status, infant gender, gestational week, history of miscarriage, and history of *in vitro* fertilization (IVF), were compared between the groups with and without a history of COVID-19 infection. Statistical analysis (Table 1) showed no significant differences between the groups ($p > 0.05$).

The intraoperative MAP follow-up values were analyzed based on the history of COVID-19 infection. Graph 1 illustrates that there was no statistically significant difference in MAP values between patients with and without a COVID-19 history ($p > 0.05$) (Fig. 1).

STAI-1 (state anxiety) and STAI-2 (trait anxiety) scores were evaluated in relation to COVID-19 history among pregnant participants. Table 2 reveals that STAI-1 scores were significantly higher in Group C (with COVID-19 history) compared to Group K (without COVID-19 history) ($p = 0.017$). However, there was no significant difference in STAI-2 scores between the two groups.

Complications during the perioperative period were compared between patients with and without COVID-19 history. Table 3 shows that there was no statistically significant difference in the development of specific complications between the groups, except for bradycardia ($p > 0.05$). Complications occurred in 67 (33.5%) of the pregnant women, with the most common being nausea and vomiting in 47 (23.5%) patients.

Apgar scores at 1st and 5th min were analyzed according to COVID-19 history. Table 4 indicates that there were no statistically significant differences in APGAR scores between the groups at the 1st min and 5th min ($p > 0.05$).

Discussion

In this study, we investigated the pre-operative anxiety levels in pregnant women undergoing elective cesarean section with and without a history of COVID-19 infection. We found a significant difference in state anxiety (STAI-1) between the two groups, suggesting that a history of COVID-19 may contribute to higher levels of acute anxiety in pregnant women facing surgery. We also identified factors such as a history of IVF and non-active employment status that influenced anxiety levels in pregnant women with COVID-19 history.

Table 1. Comparison of demographic characteristics between groups with and without a COVID-19 history

Variables	Total (n = 200)	Group C (n = 100)	Group K (n = 100)
Age (years), mean $\pm$ SD	32.0 $\pm$ 5.5	32.6 $\pm$ 5.7	31.4 $\pm$ 5.4
Weight (kg), mean $\pm$ SD	81.7 $\pm$ 12.9	81.0 $\pm$ 12.1	82.4 $\pm$ 13.8
COVID-19 symptoms, n (%)			
Asymptomatic	82	82 (72)	NA
Moderate	13	13 (20.0)	NA
Severe	5	5 (51.0)	NA
Working status, n (%)			
Yes	47	29 (29.0)	18 (18.0)
No	153	71 (71.0)	82 (82.0)
Smoking status, n (%)			
None	175 87.5	89 89.0	86 86.0
Before pregnancy	14 7.0	6 6.0	8 8.0
Before and during pregnancy	11 5.5	5 5.0	6 6.0
Gestational week, mean $\pm$ SD	37.4 $\pm$ 2.1	37.2 $\pm$ 1.8	37.6 $\pm$ 2.3
Baby gender, n (%)			
Female	105	52 (52.0)	53 (53.0)
Male	95	48 (48.0)	47 (47.0)
History of miscarriage, n (%)			
Yes	43	19 (19.0)	24 (24.0)
No	157	81 (81.0)	76 (76.0)
History of <i>in vitro</i> fertilization, n (%)			
Yes	7	4 (4.0)	3 (3.0)
No	193	96 (96.0)	97 (97.0)

SD: standard deviation; NA: not applicable.

Pregnancy inherently involves physiological and psychological changes, often accompanied by increased levels of fear and anxiety, particularly in the context of impending surgery.⁶ Studies by Rubertsson et al. and Morris et al. have consistently shown high anxiety levels among pregnant women scheduled for cesarean sections, with a significant portion experiencing severe anxiety.^{7,8} Our findings align with these studies, indicating that pre-operative anxiety remains a prevalent concern among pregnant women.

The COVID-19 pandemic has significantly influenced global mental health, exacerbating anxiety and depression in the general population.⁴ Our study aimed to evaluate this impact specifically among pregnant women who had experienced COVID-19. Studies by Khubchandani et al. and Ayaz et al. highlighted increased psychological distress and anxiety symptoms in COVID-19 survivors and pregnant women during the pandemic.^{9,10} Similarly, our findings indicate

Table 2. Comparison of STAI-1 and STAI-2 scores according to COVID-19 history

Variables	Group C	Group K	p
STAI-1 score	50.9 $\pm$ 13.0	47.0 $\pm$ 10.2	0.017*
STAI-2 score	46.4 $\pm$ 9.1	44.9 $\pm$ 7.7	0.142

*p < 0.05 indicates statistical significance.

STAI: state-trait anxiety inventory.


Table 3. Comparison of complication development according to COVID-19 history

Complications	Group C n (%)	Group K n (%)	p
Any complication present			
Yes	34 (34.0)	33 (33.0)	1.000
No	66 (66.0)	67 (67.0)	1.000
Nausea-vomiting			
Yes	23 (23.0)	24 (24.0)	1.000
No	77 (77.0)	76 (76.0)	1.000
Mask oxygen requirement			
Yes	8 (8.0)	5 (5.0)	0.568
No	92 (92.0)	95 (95.0)	1.000
Spinal block level increase			
Yes	5 (5.0)	5 (5.0)	1.000
No	95 (95.0)	95 (95.0)	1.000
Bradycardia			
Yes	1 (1.0)	5 (5.0)	0.097
No	99 (99.0)	95 (95.0)	1.000
Bleeding			
Yes	2 (2.0)	2 (2.0)	1.000
No	98 (98.0)	98 (98.0)	1.000

Table 4. Comparison of APGAR scores (1st and 5th min) according to COVID-19 history

Variables	Group C	Group K	p
APGAR 1 st min (mean $\pm$ SD)	6.3 $\pm$ 1.4	6.6 $\pm$ 1.4	0.098
APGAR 5 th min (mean $\pm$ SD)	7.8 $\pm$ 1.1	7.8 $\pm$ 1.3	0.743

SD: standard deviation.

**Figure 1. Intraoperative mean arterial pressure follow-up values according to COVID-19 history.**

that pregnant women with a history of COVID-19 exhibited significantly higher state anxiety levels compared to those without such a history.

In a survey study conducted by Zainiyah and Susanti among 70 pregnant women during the coronavirus pandemic, it was shown that 31.4% of pregnant women experienced very severe anxiety, and 12.9% experienced severe anxiety. It was emphasized that anxiety during the coronavirus pandemic is an important issue that should be addressed in pregnant women to prevent negative effects on the mother and unborn child.¹¹ In our study, the STAI-1 (state) score of pregnant women with a history of COVID-19 was statistically significantly higher than that of those without a history of COVID-19. In terms of STAI-1 levels, severe anxiety level was found to be statistically significantly higher than moderate and mild anxiety levels in pregnant women with a history of COVID-19 compared to those without a history of COVID-19.

Literature reports varying impacts of COVID-19 on perioperative complications among pregnant women.¹² Studies like those by Zhang et al. have observed higher incidences of complications such as hypotension in COVID-19-positive pregnant women undergoing cesarean sections.¹³ However, our study, consistent with findings from Kho et al., did not find significant differences in complication rates between pregnant women with and without a history of COVID-19.¹⁴

The relationship between COVID-19 during pregnancy and neonatal outcomes remains an area of active investigation. There are studies that found a significant association between low Apgar scores and neonatal intensive care unit admission in pregnant women with active COVID-19 infection, as well as studies showing that pregnancy-related anxiety increased during the COVID-19 pandemic, but this did not have a direct effect on neonatal outcomes after cesarean section.¹⁵⁻¹⁸ In our study, we did not find significant differences in Apgar scores at 1st and 5th min between groups with and without COVID-19 history, likely due to the absence of active infection and emergency cases in our study cohort.

Previous research has established a link between maternal anxiety levels and neonatal outcomes, including lower Apgar scores.^{19,20} Our study also found that higher state anxiety levels (STAI-1) were associated with lower Apgar scores at both 1st and 5th min. This underscores the importance of managing maternal anxiety to potentially improve neonatal outcomes.

Limitations

This study has limitations, including its single-center design and exclusion criteria that may limit generalizability. In addition, the study did not assess long-term maternal mental health outcomes beyond the immediate perioperative period. Future studies could benefit from larger, multicenter designs and longitudinal follow-up to comprehensively evaluate these relationships.

Conclusions

Our study highlights the heightened anxiety levels among pregnant women with a history of COVID-19 undergoing elective cesarean section. The pandemic context has exacerbated anxiety, which may impact perioperative and neonatal outcomes. Effective management of pre-operative anxiety is crucial to mitigate potential complications and improve maternal and neonatal health outcomes. Future research should continue to explore the long-term impacts of the COVID-19 pandemic on maternal mental health and pregnancy outcomes.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of human subjects and animals. The authors declare that no experiments on humans or animals were performed for this research.

Confidentiality, informed consent, and ethical approval. The authors have followed their institution's confidentiality protocols, obtained informed consent from all patients, and secured approval from the Ethics Committee. SAGER guidelines have been followed as applicable to the nature of the study. Ethical approval for this study (Ethical Committee 2022/145) was provided by the Local Ethical Committee.

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing or creation of the content of this manuscript.

References

1. Maranets I, Kain ZN. Preoperative anxiety and intraoperative anesthetic requirements. *Anesth Analg*. 1999;89:1346-51.
2. Friedrich S, Reis S, Meybohm P, Kranke P. Preoperative anxiety. *Curr Opin Anaesthesiol*. 2022;35:674-8.
3. Alipour Z, Lamyian M, Hajizadeh E, Vafaei MA. The association between antenatal anxiety and fear of childbirth in nulliparous women: a prospective study. *Iran J Nurs Midwifery Res*. 2011;16:169-73.
4. Mehdizadehkashi A, Chaichian S, Haghighi L, Eshraghi N, Bordbar A, Hashemi N, et al. The impact of COVID-19 pandemic on stress and anxiety of non-infected pregnant mothers. *J Reprod Infertil*. 2021;22:125-32.
5. Dashraath P, Wong JL, Lim MX, Lim LM, Li S, Biswas A, et al. Coronavirus disease 2019 (COVID-19) pandemic and pregnancy. *Am J Obstet Gynecol*. 2020;222:521-31.
6. Domínguez-Solís E, Lima-Serrano M, Lima-Rodríguez JS. Non-pharmacological interventions to reduce anxiety in pregnancy, labour and postpartum: a systematic review. *Midwifery*. 2021;102:103126.
7. Rubertsson C, Hellström J, Cross M, Sydsjö G. Anxiety in early pregnancy: prevalence and contributing factors. *Arch Womens Ment Health*. 2014;17:221-8.
8. Morris JR, Jaswa E, Kaing A, Hariton E, Andrusier M, Aliaga K, et al. Early pregnancy anxiety during the COVID-19 pandemic: preliminary findings from the UCSF ASPIRE study. *BMC Pregnancy Childbirth*. 2022;22:272.
9. Khubchandani J, Price JH, Sharma S, Wiblishauser MJ, Webb FJ. COVID-19 infection survivors and the risk of depression and anxiety symptoms: a nationwide study of adults in the United States. *Eur J Intern Med*. 2022;97:119-21.
10. Ayaz R, Hocaoglu M, Günay T, Yardımcı OD, Turgut A, Karateke A. Anxiety and depression symptoms in the same pregnant women before and during the COVID-19 pandemic. *J Perinat Med*. 2020;48:965-70.
11. Zainiyah Z, Susanti E. Anxiety in pregnant women during Coronavirus (COVID-19) pandemic in East Java, Indonesia. *MKB*. 2020;52:149-53.
12. COVIDSurg Collaborative. Mortality and pulmonary complications in patients undergoing surgery with perioperative SARS-CoV-2 infection: an international cohort study. *Lancet*. 2020;396:27-38.
13. Zhang Y, Chen R, Cao C, Gong Y, Zhou Q, Wei M, et al. The risk of neuraxial anesthesia-related hypotension in COVID-19 parturients undergoing cesarean delivery: a multicenter, retrospective, propensity score matched cohort study. *Front Med (Lausanne)*. 2021;8:713733.
14. Kho RM, Chang OH, Hare A, Schaffer J, Hamner J, Northington GM, et al. Surgical outcomes in benign gynecologic surgery patients during the COVID-19 pandemic (SOCOVID study). *J Minim Invasive Gynecol*. 2022;29:274-83.e1.
15. Ababneh O, Alrabayah M, El-Share' Al, Bsisu I, Bahar Y, Dabousi B, et al. Perioperative outcomes in COVID-19 obstetric patients undergoing spinal anesthesia for cesarean section: a prospective observational study. *Healthcare (Basel)*. 2021;10:23.
16. Abedzadeh-Kalahroudi M, Sehat M, Vahedpour Z, Talebian P. Maternal and neonatal outcomes of pregnant patients with COVID-19: a prospective cohort study. *Int J Gynaecol Obstet*. 2021;153:449-56.
17. Smithgall MC, Murphy EA, Rand S, Sukhu A, Singh S, Schatz-Siemers N, et al. Placental pathology, neonatal birth weight, and Apgar score in acute and distant SARS-CoV-2 infection. *J Clin Transl Res*. 2022;8:351-9.
18. Yang R, Mei H, Zheng T, Fu Q, Zhang Y, Buka S, et al. Pregnant women with COVID-19 and risk of adverse birth outcomes and maternal-fetal vertical transmission: a population-based cohort study in Wuhan, China. *BMC Med*. 2020;18:330.
19. Ossola P, Ampollini P, Gerra ML, Tonna M, Viviani D, Marchesi C. Anxiety, depression, and birth outcomes in a cohort of unmedicated women. *J Matern Fetal Neonatal Med*. 2021;34:1606-12.
20. Smorti M, Ponti L, Tani F. The effect of maternal depression and anxiety on labour and the well-being of the newborn. *J Obstet Gynaecol*. 2019;39:492-7.

Impact of psoas muscle area measurement in complex open abdominal aortic aneurysm repair

Impacto de la medición del área del músculo psoas en la reparación de aneurismas de aorta abdominal complejos

Ramon García-Alva¹, Yuki Yoshi Kimura-Sandova², Luis O. Bobadilla-Rosado¹, Javier E. Anaya-Ayala¹, Ximena Rosas-Flota³, Natalia Sanchez-Garrido⁴, Hugo Laparra-Escareno¹, Lizeth Luna¹, Luis H. Arzola¹, Adriana Torres-Machorro⁵, and Carlos A. Hinojosa^{1*}

¹Department of Surgery, Section of Vascular Surgery and Endovascular Therapy; ²Department of Radiology and Interventional Radiology;

³Department of Clinical Nutrition; ⁴Department of Geriatric. Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City, Mexico;

⁵Division of Vascular and Endovascular Surgery, Instituto Nacional de Cardiología Ignacio Chavez, Mexico City, Mexico

Abstract

Objective: Cross-sectional psoas muscle area (PMA), measured by computed tomography images, is a surrogate of the muscle mass. We aimed to evaluate the impact of PMA and length of stay (LOS) after complex open abdominal aortic aneurysm repair (OAAAR). **Methods:** A retrospective review between January 2013 and January 2017 was conducted. The main outcome was to assess the PMA as a predictor of LOS. Demographics and clinical variables were collected, and Spearman's rank coefficient was used to define the correlation of PMA and LOS. **Results:** A total of 72 patients were studied with a mean age of 72 (standard deviation $\pm$ 10.3) years, and 54 (75%) were males. The median post-operative LOS was 10 (7-14) days; lower PMA was correlated with longer LOS ($R = -0.47$, $p \leq 0.001$). Thirty-six patients (50%) had a PMA $< 17 \text{ cm}^2$, which was associated with greater post-operative 30-day complications ($p = 0.004$). There was no difference in 30-day mortality between patients with PMA $< 17 \text{ cm}^2$ and those with PMA $> 17 \text{ cm}^2$ ($p = 0.147$). **Conclusions:** Lower PMA was associated with longer LOS and a higher rate of 30-day complications; however, no statistically significant difference was found in mortality.

Keywords: Psoas muscle area. Length of stay. Abdominal aortic aneurysm. Open repair. Clinical outcomes.

Resumen

Objetivo: Evaluar el impacto de la medición del área del músculo psoas (AMP) en la estancia hospitalaria (EH) posterior a la reparación abierta de los aneurismas de aorta abdominal (RAAAA) complejos. **Métodos:** Revisión de pacientes que fueron sometidos a RAAAA entre enero de 2013 y enero de 2017. El objetivo principal fue la evaluación de la medición del AMP como predictor de EH. Se documentaron variables demográficas y clínicas, y se utilizó el coeficiente de Spearman para correlacionar el AMP y la EH. **Resultados:** Se incluyeron 72 pacientes con una media de edad de 72 años (DE: 10.3), de los que 54 (75%) eran hombres. La mediana de EH fue de 10 (7-14) días. Un AMP baja se correlacionó con una EH prolongada ($R = -0.47$; $p < 0.001$). Treinta y seis pacientes (50%) tuvieron AMP $< 17 \text{ cm}^2$, y se asoció con mayor tasa de complicaciones a 30 días ($p = 0.004$). No se observaron diferencias en cuanto a mortalidad ($p = 0.147$). **Conclusiones:** Un AMP baja se asoció a una EH más larga y a mayores tasas de complicaciones a 30 días; sin embargo, no hubo diferencia significativa en cuanto a mortalidad.

Palabras clave: Área del músculo psoas. Estancia hospitalaria. Aneurisma de aorta abdominal. Reparación abierta. Resultados clínicos.

*Correspondence:

Carlos A. Hinojosa

E-mail: carlos.a.hinojosa@gmail.com

Date of reception: 07-09-2023

Date of acceptance: 01-10-2023

DOI: 10.24875/CIRUE.M23000940

Cir Cir (Eng). 2026;94(3):235-239

Contents available at PubMed

www.cirugiaycirujanos.com

2444-0507/© 2023 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The proportion of older adults is expected to increase within the next few years, particularly in countries where the demographic transition is accelerated.¹ By 2050, almost 23% of the total population will be 60 years or older, 3 times greater than what it was in 2010. Regrettably, this has seemed to be accompanied by an increase in chronic diseases.¹ Due to the increased prevalence of arterial hypertension and diabetes, there will also be an increase in the incidence of arterial diseases; therefore, more vascular interventions in the elderly population will be needed. Age-related changes have demonstrated a negative impact on outcomes in the older population, such as increased mortality and complications after abdominal surgery, and some others related just to hospitalization. Some age-related complications, such as delirium, urinary incontinence, pressure ulcers, depression, infection, and decline in functionality, can influence post-operative outcomes.^{2,3} In terms of surgery decision making, the eligibility of older adults has become an important issue, and the need for objective parameters that could help to identify patients at higher risk of post-operative complications and hospitalization length of stay (LOS) is still necessary.⁴

Sarcopenia is a geriatric syndrome characterized by a decline in skeletal muscle mass and decreased strength,⁵ it has been studied in multiple scenarios. It is known to be related to higher mortality and post-operative complications in older adults after abdominal and cardiovascular surgeries.^{2,6,7} There are many ways to identify sarcopenia, and the choice of measurement depends on the patient and the resources that each hospital can access. Psoas muscle area (PMA) measured by computed tomography (CT) has been used to identify sarcopenia in pre-operative patients.⁵ Some authors have found PMA association with adverse outcomes in cardiovascular surgeries.⁸ This study aims to determine the association of PMA and LOS after open aortic aneurysm repair, and whether PMA has an impact on post-operative complications in these patients.

Methods

We performed a retrospective perioperative CT scan analysis of patients who underwent open abdominal aortic aneurysm repair (OAAAR) at our third-level referral center in Mexico City. The PMA was measured

at the level of the L3 vertebrae; this was defined as the CT slice including the most inferior portion of the L3 vertebral body. Data of patients with an OAAAR treated between January 2013 and January 2017 were collected. Patients with an acute rupture were excluded because of higher morbidity and mortality. Demographic characteristics, complications, and outcomes were obtained from medical records.

Patients were divided into two groups to compare LOS, we defined the cut-off point, as the median of days after surgery in all patients. As a secondary outcome, we divided the patients by their mean PMA into two groups for comparison. This research project follows the regulations of an institution and the generally accepted guidelines governing this work. For this study, informed consent was not required.

PMA determination

CT scans were performed before all procedures for clinical evaluation, and Osirix MC (Pixmeo SARL, Geneva, Switzerland) multiplanar reformatting software was used to measure the right and left PMA in an axial plane. (Fig. 1) The L3 level was selected in accordance with the European Working Group on Sarcopenia in Older Adults recommendations for the diagnosis of sarcopenia.⁵ Interobserver mean differences were assessed in two observers researches by measuring the PMA in 20 randomized patients. PMA values were represented as continuous variables with a mean PMA in the study group.

Statistical analysis

Categorical variables are shown as frequencies and continuous variables as mean $\pm$ standard deviation or as median and interquartile range. Spearman's rank correlation coefficient was used to assess the linear correlation between PMA and LOS. Distribution of categorical variables was compared using the χ^2 test. Continuous variables were compared with a univariate test, and statistical significance was considered at a two-sided $p < 0.05$. Data were collected and analyzed using the Statistical Package for the Social Sciences (IBM SPSS, version 22, IBM Corp., NY, USA).

Results

A total of 72 patients were included in this study, with a mean age of 72 ± 10 years, and 75% (54 patients)




Figure 1. Cross-sectional view of the computed tomography with the Psoas muscle area measurement in a patient with an abdominal aortic aneurysm.

were male. Regarding comorbidity, 66% had hypertension, 20% diabetes, and 86% had a history of smoking. The median abdominal aortic aneurysm (AAA) diameter in centimeters was 64 (55-74) (Table 1).

The mean interobserver PMA difference found (one radiology fellow and one vascular surgeon fellow) in 20 patients was 0.4 cm² (95% confidence interval [CI] -0.9-1.2 cm²), and the mean interobserver difference was 0.5 cm² (95% CI -1.3-2.1 cm²). The correlation obtained by the Spearman test between PMA and post-operative LOS was negative ($R = -0.47$, $p < 0.001$), supporting the fact that patients with lower PMA were more likely to have longer LOS after surgery (Fig. 2).

The median LOS for the 72 patients included in the study was 10 (7-14) days, after dividing groups by their median, the first group included 42 patients (58.3%) who stayed < 10 days after OAAAR, with a median age at surgery of 73 (68-78) years, 33 patients (79%) were males, with median AAA diameter of 65 (56-75) cm and a median PMA of 19 (14-23) cm². After comparing the PMA between groups, those patients who stayed more than 10 days after surgery had a smaller PMA compared to those who stayed < 10 days - 14 (11-19) versus 19 (14-23) cm², $p = 0.004$ (Table 2).

As a secondary outcome, 30-day complications were assessed using PMA for the comparison between groups. The cut point used to separate groups was the mean PMA obtained in the 72 patients (17 cm²) (Table 3). We observed that complications after 30 days were present in 71% (26 patients) in the < 17 cm² PMA group compared to 39% in patients with PMA > 17 cm² ($p = 0.004$) (Table 3). None of the complications required surgery intervention. The most




Figure 2. Spearman correlation of length of stay and psoas muscle area. $R = -0.47$ ($p < 0.001$).

Table 1. Demographic and clinical characteristics of the study population

Characteristics	n = 72
Age (years)	72 ± 10
Sex male, n (%)	54 (75)
Diabetes, n (%)	15 (20)
Hypertension, n (%)	48 (66)
CAD, n (%)	12 (16)
Chronic kidney insufficiency, n (%)	15 (20)
Smoke, n (%)	62 (86)
Height (cm)	163 (153-171)
Weight (kg)	67 ± 13
Aortic aneurysm diameter (cm)	64 (55-74)
Creatinine (mg/dL)	1.08 (0.8-1.3)
Albumin level g/dL	3.7 ± 0.66
Psoas muscle area (cm ²)	17 ± 5.8
LOS (days)	10 (7-14)
30-day complication	40 (55)

LOS: length of stay; CAD: coronary artery disease.

common complications that we found were: respiratory complications (30%) as pulmonary embolism and pneumonia; cardiac complications (25%) as arrhythmias, myocardial infarction, and cardiac failure; renal complications (15%) as acute renal failure, urinary retention, and urinary tract infections; and other complications (30%) such as incisional hernia, small bowel obstruction or wound infection. There was no significant difference in death between groups ($p = 0.147$); however, those patients with PMA < 17 cm² had longer LOS 12 (9-17) versus 8 (5-11) days ($p = 0.001$).

Table 2. Comparison of patients divided by length of stay in the post-operative

Characteristics	LOS < 10d (n = 42)	LOS > 10d (n = 30)	p
Age (years)	73 (68-78)	74 (67-80)	0.468
Sex male, n (%)	33 (79)	21 (70)	0.408
Diabetes, n (%)	9 (21)	6 (20)	0.888
Hypertension, n (%)	29 (69)	19 (63)	0.612
CAD (%)	7 (17)	5 (17)	1.0
Chronic renal failure, n (%)	8 (19)	7 (23)	0.659
Smoke, n (%)	38 (90)	24 (80)	0.205
Height (cm)	161 (150-172)	165 (156-170)	0.441
Weight (kg)	68 ± 13	65 ± 13	0.392
Aortic aneurysm diameter (cm)	65 (56-75)	61 (55-72)	0.355
Creatinine (mg/dL)	1.08 (0.8-1.3)	1.08 (0.8-1.5)	0.918
Albumin g/dL	3.8 ± 0.63	3.5 ± 0.68	0.065
Psoas muscle area (cm ²)	19 (14-23)	14 (11-19)	0.004*

*p < 0.05.

LOS: length of stay; CAD: coronary artery disease.

Table 3. Comparison of patients divided by Psoas muscle area in the post-operative

Characteristics	PMA < 17cm ² (n = 36)	PMA > 17cm ² (n = 36)	p
Age (years)	75 (69-81)	71 (66-77)	0.32
Sex male, n (%)	20 (56)	34 (94)	0.000*
Diabetes, n (%)	8 (22)	7 (19)	0.772
Hypertension, n (%)	22 (61)	26 (72)	0.317
CAD, n (%)	6 (17)	6 (17)	1.0
Chronic renal failure, n (%)	6 (17)	9 (25)	0.384
Smoke, n (%)	28 (78)	34 (94)	0.041*
Height (cm)	161 (152-165)	168 (158-175)	0.062
Weight (kg)	61 ± 11.5	73 ± 12	0.000*
Aortic aneurysm diameter (cm)	67 (55-80)	62 (57-71)	0.460
Creatinine (mg/dl)	1.0 (0.8-1.3)	1.1 (0.8-1.4)	0.288
Albumin g/dL	3.6 ± 0.7	3.8 ± 0.6	0.214
Mortality	10 (28)	5 (14)	0.147
30-day complications	26 (71)	14 (39)	0.004*
LOS	12 (9-17)	8 (5-11)	0.001*

*p < 0.05.

PMA: psoas muscle area; CAD: coronary artery disease; LOS: length of stay.

Discussion

Once an aneurysm in the abdominal aorta has been diagnosed and after it reaches the threshold for elective repair, the risk for post-operative complications and death must be assessed against the risk of rupture.⁹ Consequently, many prediction models have been described to estimate morbidity and mortality; however, many of these include poor suitability variables for OAAAR.^{10,11} Frailty is a geriatric syndrome where low muscle mass and strength play an important role, contributing to a diminished homeostatic capability and leading patients to a higher vulnerability to stressors such as surgical interventions and post-operative time.¹¹⁻¹⁶ This syndrome becomes more prevalent with increasing age, and sarcopenia, which is defined as a loss of muscle mass and strength, is an important characteristic of the latter.^{5,17} In addition, as it has been reported as an easy component to measure by imaging techniques such as CT⁹ and due to the relevant association that sarcopenia has with post-operative outcomes such as LOS and major adverse complications, it has gained interest by many authors in the surgical field.^{12,16,18-22}

Nowadays, there is growing evidence that PMA can predict negative post-operative outcomes such as an increasing patient morbidity and mortality.^{12,16,19-22} In regard to the proposed threshold to define low PMA in these patients, there is no universally accepted cutoff point, since some authors have chosen to stratify their patients in percentiles^{7,14,16,18} and others have chosen a random cut point.^{12,20,23} As no threshold value for sarcopenia exists currently in the literature regarding PMA,^{21,22} we decided to stratify our post-operative patients into those above the mean PMA obtained from patients' CT scans reviewed (mean = 17 cm²). Using our proposed threshold, 36 patients (50%) were defined as having a low PMA, showing a statistically significant association (p = 0.004) with greater post-operative 30-days complications as well as an increased LOS (> 10 days) (p = 0.001) identifying PMA as a radiologic biomarker of frailty, helping in the prediction of post-operative morbidity in patients considered for complex abdominal aortic aneurysm open repair.

Clinical research has been growing, focusing on the association of a low PMA with an increased post-operative morbidity and mortality, as stated in a systematic review by Antoniou and colleagues²⁴ confirming the relevance of low skeletal muscle mass as a prognostic factor for survival. LOS then becomes a

crucial post-operative outcome to consider for a physician when deciding on the best course of treatment for patients, as an increased LOS leads to higher costs for patients and the healthcare system. We found in the present study that a lower PMA was significantly associated with longer LOS, which supports the evidence from previous studies. Our results are in concordance with those reported by Zuckerman et al.,¹² who reported an increased LOS in patients with a lower PMA.

Limitations of the present study are those inherent in the retrospective nature of our work and possible selection bias.

Conclusion

Patients with the lower PMA (below 17 cm²) were associated with longer LOS and a higher rate of 30-day complications. However, no statistically significant difference was found in mortality. PMA measurement might assist as a postoperative predictor of LOS in major vascular surgery.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of human subjects and animals. The authors declare that the procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation and with the World Medical Association and the Declaration of Helsinki. The procedures were authorized by the Institutional Ethics Committee.

Confidentiality, informed consent, and ethical approval. The authors have followed their institution's confidentiality protocols, obtained informed consent from all patients, and secured approval from the Ethics Committee. SAGER guidelines have been followed as applicable to the nature of the study.

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing or creation of the content of this manuscript.

References

- Angel JL, Vega W, López-Ortega M. Aging in Mexico: population trends and emerging issues. *Gerontologist*. 2017;57:153-62.
- Massarweh NN, Legner VJ, Symons RG, McCormick WC, Flum DR. Impact of advancing age on abdominal surgical outcomes. *Arch Surg*. 2009;144:1108-14.
- Creditor MC. Hazards of hospitalization of the elderly. *Ann Intern Med*. 1993;118:219-23.
- Houghton JS, Nickinson AT, Morton AJ, Nduwayo S, Pepper CJ, Rayt HS, et al. Frailty factors and outcomes in vascular surgery patients: a systematic review and meta-analysis. *Ann Surg*. 2020;272:266-76.
- Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyere O, Cederholm T, et al. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing*. 2019;48:16-31.
- Cao Q, Xiong Y, Zhong Z, Ye Q. Computed tomography-assessed sarcopenia indexes predict major complications following surgery for hepatopancreatobiliary malignancy: a meta-analysis. *Ann Nutr Metab*. 2019;74:24-34.
- Okamura H, Kimura N, Tanno K, Mieno M, Matsumoto H, Yamaguchi A, et al. The impact of preoperative sarcopenia, defined based on psoas muscle area, on long-term outcomes of heart valve surgery. *J Thorac Cardiovasc Surg*. 2018;157:1071-9.e3.
- Ghaffarian AA, Foss WT, Donald G, Kraiss LW, Sarfati M, Griffin C, et al. Prognostic implications of diagnosing frailty and sarcopenia in vascular surgery practice. *J Vasc Surg*. 2019;70:892-900.
- Hinojosa CA, Anaya-Ayala JE, Bermúdez-Serrato K, Leal-Anaya P, Laparra-Escareno H, Torres-Machorro A. Valor predictivo del tamizaje con tomografía computarizada en la detección de aneurismas aórticos en la población mexicana mayor de 55 años [Predictive value of computed tomographic screening of aortic aneurysms in the Mexican population older than 55 years]. *Gac Med Mex*. 2017;153 Suppl 2:S27-33.
- Hinojosa CA, Anaya-Ayala JE, Carrera-Ramirez M, Leal-Anaya P, García-Alva R, Lozano-Corona R. Factors that impact kidney function during open and endovascular reconstruction of aortic aneurysms. *Cir Cir*. 2019;87:170-5.
- Indrakusuma R, Zijlman JL, Jalalzadeh H, Planken RN, Balm R, Koelmay MJ. Psoas muscle area as a prognostic factor for survival in patients with an asymptomatic infrarenal abdominal aortic aneurysm: a retrospective cohort study. *Eur J Vasc Endovasc Surg*. 2017;55:83-91.
- Zuckerman J, Ades M, Mullie L, Trnkus A, Morin JF, Langlois Y, et al. Psoas muscle area and length of stay in older adults undergoing cardiac operations. *Ann Thorac Surg*. 2017;103:1498-504.
- Michel J, Pellegrini C, Rheude T, Von Scheidt M, Trenkwalder T, Elhmidi Y, et al. The clinical impact of psoas muscle cross-sectional area on medium-term mortality after transcatheter aortic valve implantation. *Heart Lung Circ*. 2020;29:904-13.
- Jones K, Gordon-Weeks A, Coleman C, Silva M. Radiologically determined sarcopenia predicts morbidity and mortality following abdominal surgery: a systematic review and meta-analysis. *World J Surg*. 2017;41:2266-79.
- Shah N, Abeyundara L, Dutta P, Christodoulidou M, Wylie S, Richards T, et al. The association of abdominal muscle with outcomes after scheduled abdominal aortic aneurysm repair. *Anaesthesia*. 2017;72:1107-11.
- Lee JS, He K, Harbaugh CM, Schaubel DE, Sonnenday CJ, Wang SC, et al. Frailty, core muscle size, and mortality in patients undergoing open abdominal aortic aneurysm repair. *J Vasc Surg*. 2011;53:912-7.
- Walston J, Hadley EC, Ferrucci L, Guralnik JM, Newman AB, Studenski SA, et al. Research agenda for frailty in older adults: toward a better understanding of physiology and etiology: summary from the American geriatrics society/national institute on aging research conference on frailty in older adults. *J Am Geriatr Soc*. 2006;54:991-1001.
- Shen W, Punyanitya M, Wang Z, Gallagher D, St-Onge MP, Albu J, et al. Total body skeletal muscle and adipose tissue volumes: estimation from a single abdominal cross-sectional image. *J Appl Physiol* (1985). 2004;97:2333-8.
- Drudi LM, Phung K, Ades M, Zuckerman J, Mullie L, Steinmetz OK, et al. Psoas muscle area predicts all-cause mortality after endovascular and open aortic aneurysm repair. *Eur J Vasc Endovasc Surg*. 2016;52:764-9.
- Thurston B, Pena GN, Howell S, Cowled P, Fitridge R. Low total psoas area as scored in the clinic setting independently predicts midterm mortality after endovascular aneurysm repair in male patients. *J Vasc Surg*. 2018;67:460-7.
- Newton DH, Kim C, Lee N, Wolfe L, Pfeifer J, Amendola M. Sarcopenia predicts poor long-term survival in patients undergoing endovascular aortic aneurysm repair. *J Vasc Surg*. 2018;67:453-9.
- Cheng BT, Soult MC, Helenowski IB, Rodriguez HE, Eskandari MK, Hoel AW. Sarcopenia predicts mortality and adverse outcomes after endovascular aneurysm repair and can be used to risk stratify patients. *J Vasc Surg*. 2017;70:1576-84.
- Huber TC, Keefe N, Patrie J, Tracci MC, Sheeran D, Angle JF, et al. Predictors of all-cause mortality after endovascular aneurysm repair: assessing the role of psoas muscle cross-sectional area. *J Vasc Interv Radiol*. 2019;30:1972-9.
- Antoniou GA, Rojoa D, Antoniou SA, Alfahad A, Torella F, Juszczak MT. Effect of low skeletal muscle mass on post-operative survival of patients with abdominal aortic aneurysm: a prognostic factor review and meta-analysis of time-to-event data. *Eur J Vasc Endovasc Surg*. 2019;58:190-8.

Predictors of postoperative neurological deficits in traumatic thoracolumbar fractures

Predictores de déficits neurológicos posoperatorios en fracturas toracolumbares traumáticas

Alp Karaaslan 

Department of Neurosurgery, Sancaktepe Training and Research Hospital, Istanbul, Turkey

Abstract

Objective: The goal of this retrospective study was to assess the outcomes of vertebral stabilization following acute traumatic thoracolumbar (TL) fractures. **Methods:** The study examined the mechanism and location of trauma, pre-operative and post-operative neurological deficits, stabilization level, screw malposition, cerebrospinal fluid fistula, development of post-operative infections, additional pathological findings, mobilization times, and follow-up durations. **Results:** The study included 55 patients. Falls were the leading cause of trauma (75%), and L1 was the most common fracture site (30.9%). A4 was the most common fracture type (61.8%), with T11-L3 being the most frequently stabilized level (32.7%). Screw malposition occurred in 3.6% of cases, and CSF fistula was more common in types B and C. The presence of pre-operative neurodeficits and TL American Orthopedic Spine Injury Score (AOSIS) type B and C significantly increased the risk of post-operative neurological complications. Univariate analysis showed that pre-operative neurodeficits (odds ratio [OR]: 396, 95% confidence interval [CI]: 22-6935, $p < 0.001$) and TL AOSIS types B (OR: 11.78, 95% CI: 1.88-73.58, $p = 0.008$) and C (OR: 9.9, 95% CI: 1.31-74.73, $p = 0.026$) significantly increased the risk of post-operative neurodeficits. **Conclusions:** This study, therefore, takes into consideration the strong impact of pre-operative neurological deficit and injury severity with respect to TL classifications on post-operative outcomes for patients undergoing surgical stabilization for traumatic fractures.

Keywords: Traumatic spine fracture. Thoracolumbar fracture. Trauma.

Resumen

Objetivo: Evaluar los resultados de la estabilización vertebral tras fracturas traumáticas agudas toracolumbares. **Métodos:** Se evaluaron el mecanismo y la localización del trauma, los déficits neurológicos preoperatorios y posoperatorios, el nivel de estabilización, la mala posición de los tornillos, la fístula de líquido cefalorraquídeo, el desarrollo de infecciones posoperatorias, los hallazgos patológicos adicionales, los tiempos de movilización y las duraciones del seguimiento. **Resultados:** El estudio incluyó 55 pacientes. Las caídas fueron la principal causa de trauma (75%), L1 fue el sitio de fractura más común (30.9%), el tipo de fractura más frecuente fue A4 (61.8%) y T11-L3 fue el nivel estabilizado con mayor frecuencia (32.7%). La mala posición de los tornillos se observó en el 3.6% de los casos y la fístula de líquido cefalorraquídeo fue más común en los tipos B y C. La presencia de déficits neurológicos preoperatorios y tipo B, C de la clasificación TL AOSIS aumentó significativamente el riesgo de complicaciones neurológicas posoperatorias. El análisis univariado mostró que los déficits neurológicos preoperatorios (OR: 396; IC 95%: 22-6935; $p < 0.001$) tipo B (OR: 11.78; IC 95%: 1.88-73.58; $p = 0.008$) y tipo C (OR: 9.9; IC 95%: 1.31-74.73; $p = 0.026$) de TL AOSIS aumentaron significativamente el riesgo de déficits neurológicos posoperatorios.

Correspondence:

Alp Karaaslan

E-mail: alpkaaslan44@gmail.com

2444-0507/© 2025 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 04-04-2025

Date of acceptance: 07-08-2025

DOI: 10.24875/CIRUE.M25000941

Cir Cir (Eng). 2026;94(3):240-247

Contents available at PubMed

www.cirugiaycirujanos.com

Conclusiones: *Este estudio toma en consideración el fuerte impacto del déficit neurológico preoperatorio y la gravedad de la lesión con respecto a clasificaciones toracolumbares en resultados posoperatorios de pacientes sometidos a estabilización quirúrgica por fracturas traumáticas.*

Palabras clave: *Fractura traumática de columna. Fractura toracolumbar. Trauma.*

Introduction

Traumatic thoracolumbar (TL) fractures are one of the most significant causes of disability, deformity, and neurological deficits among working-age individuals in both industrialized and developing countries.¹⁻⁴ These fractures most commonly affect men between the ages of 20 and 40, and in approximately 50% of cases, they can lead to significant medical, social, and economic consequences.^{2,3,5} It is estimated that 50-60% of TL fractures occur in the transition zone between T11 and L2, 25-50% in the thoracic region, and only 10-14% in the lumbar and sacral regions.⁶⁻⁸ According to various studies, compression fractures (type A) account for 63-82% of all traumas, distraction injuries (type B) for 14-21%, and rotational injuries (type C) for 4-16%.^{9,10} The likelihood and extent of neurological deficits vary depending on the type of fracture, with an incidence of up to 55% in the latest AO classification's type C fractures.^{7,11} Spinal cord injuries are often the result of high-energy blunt trauma, with 56-67% associated with motor vehicle accidents and falls from great heights. These injuries are commonly associated with other injuries, such as rib fractures and pneumothorax, due to the high-speed nature of the trauma.^{11,12} In addition, the incidence of concurrent intra-abdominal injuries can be as high as 50%. Over time, many classifications have been developed to assist clinicians in treatment selection; however, while there is consensus on surgery for flexion-distraction and rotational injuries, a definitive recommendation on how to treat compression injuries is still lacking.^{11,13} Although various approaches and surgical techniques have been described for the treatment of TL fractures, scientific evidence is still insufficient to support the selection of one surgical technique as being more effective than others.^{7,14,15}

The goal of this retrospective study was to assess the outcomes of vertebral stabilization following acute traumatic TL fractures. We aimed to examine these outcomes with patient clinical data, the type and location of the fracture, the presence of neurological damage, and neurodeficit risk factor analysis.

Methods

A retrospective analysis was conducted on 55 patients who underwent surgery in our clinic between 2018 and 2024 (Fig. 1). The study examined the mechanism and location of trauma, pre-operative and post-operative neurological deficits, stabilization level, screw malposition, cerebrospinal fluid (CSF) fistula, development of post-operative infections, additional pathological findings, mobilization times, and follow-up durations. All patients underwent an imaging study of the vertebral column, which included anteroposterior and lateral X-rays, computed tomography, and magnetic resonance imaging. The purpose was to outline the exact morphology of the lesion and the possible spine involvement. Fractures were radiologically classified using the AO spine injury scores, and the findings were evaluated in the context of the existing literature (Fig. 2).

This study was approved by the non-interventional ethics committee of Sancaktepe Training and Research Hospital. This study was conducted in accordance with the principles of the Declaration of Helsinki, as revised in 2013. Before surgery, written informed consent was obtained from all patients or their legal guardians.

Patients who presented to the emergency department due to trauma and were determined to require surgery were included in the study. Patients diagnosed with osteoporosis who developed fractures, as well as those who sustained fractures due to trauma but did not require surgical intervention, were excluded from this study.

In this study, spine injuries were classified according to the AOSpine TL spine injury classification system. In response to the limitations of previous classification systems in gaining global acceptance, Vaccaro et al. developed the AOSpine TL spine injury classification system in 2013.¹⁶ This new system incorporates elements from both the Magerl classification and the TL injury classification and severity score. The morphologic classification is a simplified version of the Magerl system, featuring a total of nine injury patterns. These injuries are categorized into three




Figure 1. Patient flow chart.




Figure 2. Radiological image. A and B: pre-operative magnetic resonance imaging image. C: post-operative MRI image, D: post-operative 3D computed tomography image.

primary types: Type A – compression injuries, Type B – tension band injuries, and Type C – translation injuries. Type A and Type B injuries are further divided into five and three subtypes, respectively (Table 1).




Figure 3. Intraoperative view.

All patients underwent surgical stabilization of the fracture site, utilizing a posterior approach under general anesthesia.^{7,11,14,17} The number of levels fused varied depending on the type and location of the fracture, as well as the difficulty in realigning the spinal column, with the goals of restoring sagittal alignment and ensuring implant stability. In cases involving spinal canal compromise with neurological deficits, spinal cord decompression was also performed. For patients where pedicle screws alone could not provide sufficient stability due to osteoporosis, bone cement and fenestrated screws were used (Fig. 3).

The data for our study were analyzed using Jamovi software version 2.4.1. Descriptive statistics were presented as counts and percentages for categorical variables and as means with standard deviations for continuous variables. The Pearson Chi-square test was applied to determine differences between groups for categorical variables. To assess the normality of the distribution for continuous variables, the Shapiro–Wilk test was utilized. For comparisons between two groups, the independent t-test was employed. A binary logistic regression test was used to test the predictors. A $p < 0.05$ was considered statistically significant.

Results

The study included 55 patients with a mean age of 42.1 years (min-max: 12-83). The majority of the patients were male (69%). The primary cause of trauma was falls, accounting for 75% of the cases, followed by motor vehicle accidents at 18%, and other injuries at 7%. Vertebral fractures were most commonly observed at the L1 level, representing 30.9% of the cases. Other frequently affected levels included L2 (16.3%), T12 (14.6%), L3 (14.4%), and L2-L4 (7.2%). Preoperatively, 18.2% of the patients presented with neurodeficits, with 5.5% having paraparesis and 12.7% experiencing plegia. Postoperatively, the incidence of neurodeficits remained consistent, with 18.2% of the patients displaying neurological

Table 1. TL AOSIS types

Subgroup	Description	TL AOSIS
Type A: compression fractures		
A0	An injury that has no possibility of affecting the structural integrity of the spine (ie, fracture of the spinous or transverse process)	0
A1	A fracture through a single endplate that does not extend into the posterior wall	1
A2	A fracture through both endplates that does not extend into the posterior wall	2
A3	A fracture through a single endplate that does extend into the posterior wall (incomplete burst)	3
A4	A fracture through both endplates that does extend into the posterior wall (complete burst)	5
Type B: Tension band injury		
B1	A completely osseous tension band injury (ie, a bony chance fracture)	5
B2	An injury that disrupts the posterior tension band	6
B3	An injury that disrupts the anterior tension band	7
Type C: Translational injuries		
C	Any injury that results in translation of the vertebral body	8

TL: thoracolumbar.

impairment, distributed equally between paraparesis (5.5%) and plegia (12.7%). Screw malposition was noted in 3.6% of the cases, while a CSF fistula and infections occurred in 1.8% of the patients. The mean time to mobilization post-surgery was 1.2 days (min-max: 1-5). The mean follow-up period was 3.3 years (min-max: 1-6), during which the outcomes of the surgical interventions were closely monitored. During the follow-up period, no patients required reintervention (Table 2).

The TL injury classification for the 55 patients included in the study revealed that the majority of cases were categorized as A4, accounting for 61.8% of the total. This was followed by B2 injuries, which comprised 21.8% of the cases, and C injuries, representing 14.5%. A2 injuries were the least common, with only 1.8% of the cases falling into this category (Table 3).

The analysis of stabilization levels among the 55 patients revealed that the most common stabilization level was T11-L3, accounting for 32.7% of the cases. This was followed by the L1-L5 stabilization level, which was observed in 9.1% of the patients. Other frequently stabilized segments included T10-L2 (7.3%), T12-L4 (5.5%), and L2-L5 (5.5%) (Table 4).

The average age differed among the groups, with patients classified as type A having a mean age of 45.8 years, while those in groups B and C were younger, with mean ages of 38.3 years and 31.5 years, respectively, though this difference was not statistically significant ($p = 0.059$). Gender distribution did not differ significantly across the groups ($p = 0.546$). A significant

difference was observed in the presence of pre-operative neurodeficits, which were rare in type A injuries (2.9%) but more common in types B (41.7%) and C (50%) ($p < 0.001$). Post-operative neurodeficits followed a similar pattern, with higher rates in type B (41.7%) and C (37.5%) compared to type A (5.7%) ($p = 0.006$). The cause of trauma did not show a significant difference across the injury types ($p = 0.617$). Screw malposition was only observed in the type A group (5.7%) ($p = 0.553$). However, CSF fistula occurrence showed a significant difference, being more common in types B (25%) and C (25%) compared to type A (2.9%) ($p = 0.040$). Infection rates did not differ significantly among the groups ($p = 0.748$) (Table 5).

The univariate analysis revealed that the presence of pre-operative neurodeficit and the TL AOSIS type were significantly associated with post-operative neurodeficit occurrence. Specifically, having a pre-operative neurodeficit significantly increased the risk of developing a post-operative neurodeficit (OR: 396, 95% CI: 22-6935, $p < 0.001$). In addition, patients with TL AOSIS type B had a significantly higher risk of post-operative neurodeficit compared to those with type A (OR: 11.78, 95% CI: 1.88-73.58, $p = 0.008$). Similarly, those with type C also demonstrated an increased risk (OR: 9.9, 95% CI: 1.31-74.73, $p = 0.026$) (Table 6).

Discussion

In this study, we found that pre-operative neurodeficits and the type of TL injury, as classified by the AOSpine

Table 2. Basic characteristics of the patients

Basic characteristics	n = 55	%
Age (year; mean $\pm$ SD, min-max)	42.1 $\pm$ 18.3	12-83
Gender (M)	34	69
Cause of trauma		
Falls	41	75
Motor vehicle accidents	10	18
Other injuries*	4	7
Vertebral fracture levels		
T4-T5	1	1.8
T7	1	1.8
T10-T11	1	1.8
T12	8	14.6
T12-L1	1	1.8
L1	17	30.9
L2	9	16.3
L2-L4	4	7.2
L3	8	14.4
L3-L4	1	1.8
L3-L5	1	1.8
L4	3	5.5
Pre-operative neurodeficit (yes)	10	18.2
Type of neurodeficit (pre-operative)		
Paraparesis	3	5.5
Plegia	7	12.7
Post-operative neurodeficit	10	18.2
Type of neurodeficit (post-operative)		
Paraparesis	3	5.5
Plegia	7	12.7
Screw malposition (yes)	2	3.6
Infections	1	1.8
CSF fistula (yes)	1	1.8
Time for mobilization (day; mean $\pm$ SD, min-max)	1.2 $\pm$ 0.6	1-5
Follow-up (year; mean $\pm$ SD, min-max)	3.3 $\pm$ 1.5	1-6

*Other injuries: assault, epileptic seizure, gunshot.

SD: standard deviation; CSF: cerebrospinal fluid; M: male; T: thoracic; L: lumbar.

Table 3. TL AOSIS type of fracture

TL AOSIS	n	%	Cumulative %
A2	1	1.8	1.8
A4	34	61.8	63.6
B2	12	21.8	85.5
C	8	14.5	100
Total	55	100	100

TL: thoracolumbar.

system, were significant predictors of post-operative outcomes. Patients presenting with pre-operative neurodeficits had a markedly higher risk of developing post-operative neurodeficits, emphasizing the critical

Table 4. Frequencies of stabilization level

Stabilization level	n	%
L1-L3	1	1.8
L1-L4	2	3.6
L1-L5	5	9.1
L2-L5	3	5.5
L2-S1	1	1.8
L3	1	1.8
L3-L5	1	1.8
T10-L1	1	1.8
T10-L2	4	7.3
T10-L3	2	3.6
T10-T12	1	1.8
T10-L2	1	1.8
T11-L5	2	3.6
T11-L6	1	1.8
T11-L3	18	32.7
T11-L3	1	1.8
T11-L4	1	1.8
T12-L3	1	1.8
T12-L4	3	5.5
T12-L5	2	3.6
T2-T7	1	1.8
T5-T9	1	1.8
T8-T12	1	1.8

impact of early neurological impairment on surgical outcomes. Furthermore, injury types B and C were associated with significantly higher risks of post-operative neurodeficits compared to type A injuries, underscoring the severity of these injury patterns.

Despite significant advancements in diagnostic techniques and spinal instrumentation, there remains no clear consensus on the optimal treatment approach for TL fractures.^{11,18-20} Treatment options vary widely, ranging from conservative management with orthotic external bracing to more invasive surgical procedures, such as instrumented fusions.^{11,13,15} Recently, percutaneous fixation has emerged as a minimally invasive alternative, providing a middle ground between these two extremes. Neeley et al. compared deformity correction and surgical outcomes between

Table 5. Comparative analysis of items regarding TL AOSIS

Characteristics	A (n = 35)	B (n = 12)	C (n = 8)	p
Age (year)	45.8 ± 18.6	38.3 ± 17.8	31.5 ± 13	0.059
Gender (M), n (%)	20 (57)	9 (75)	5 (62.5)	0.546
Any pre-operative neurodeficit, n (%)	1 (2.9)	5 (41.7)	4 (50)	< 0.001
Any post-operative neurodeficit, n (%)	2 (5.7)	5 (41.7)	3 (37.5)	0.006
Cause of trauma, n (%)				0.617
Falls	26 (74.3)	8 (66.7)	7 (87.5)	
Motor vehicle accidents	7 (20)	2 (16.7)	1 (12.5)	
Other injuries*	2 (5.7)	2 (16.7)	0 (0)	
Screw malposition	2 (5.7)	0 (0)	0 (0)	0.553
CSF fistula	1 (2.9)	3 (25)	2 (25)	0.040
Infections	1 (2.9)	0 (0)	0 (0)	0.748

*Other injuries: assault, epileptic seizure, gunshot.

CSF: cerebrospinal fluid; M: male.

Table 6. Binary logistic regression analysis

Characteristics	Univariate analysis		p	OR	95% CI	Multivariate analysis		p
	Estimate	SE				Estimate	SE	
Age (year)	-0.0163	0.0202	0.419	0.984	0.945-1.02			
Gender (ref: Female)	1-0.442	0.754	0.558	0.643	0.147-2.189			
Any pre-operative neurodeficit	5.98	1.46	< 0.001	396	22-6935	39.44	6763	0.995
Cause of trauma								
Motor vehicle accidents - falls	-0.617	1.133	0.586	0.540	0.058-4.970			
Other injuries* - falls	1.580	1.083	0.144	4.857	0.581-40.55			
Screw malposition	-15.11	1696	0.993		0.39-16.44			
CSF fistula	0.942	0.948	0.321	2.563				
Infections	-15.08	2399	0.995					
TL AOSIS type (ref: A)								
B-A	2.47	0.934	0.008	11.78	1.88-73.57	-16.33	4619	0.997
C-A	2.29	1.031	0.026	9.9	1.31-74.73	-34.85	6763	0.996

*Other injuries: assault, epileptic seizure, gunshot.

CSF: cerebrospinal fluid. OR: odds ratio; SE: standard error; CI: confidence interval.

percutaneous instrumentation and open fusion in patients with traumatic TL fractures, highlighting the evolving landscape of treatment options.¹⁵ Reinhold et al. reported using a posterior approach in approximately half of their cases, with the remainder divided between combined anteroposterior and exclusively anterior approaches.¹⁴ Similarly, Muratore et al. emphasized a tailored approach, choosing between posterior and combined anteroposterior methods based on the anatomical and overall clinical status of each patient.¹¹ In our study, we employed a uniform strategy, performing surgical stabilization using the posterior approach

in all patients. This choice was guided by our clinical experience and the specific characteristics of the fractures observed, reinforcing the utility of a consistent approach in achieving reliable outcomes.

According to a meta-analysis, the most commonly injured vertebra was L1, with an incidence rate of 34.4% (95% CI 18.2%, 50.3%).²¹ The most frequently injured non-junctional vertebra was T7, at 3.9% (95% CI 2.81%, 4.99%), while T2 was the least affected, with an incidence of 0.26% (95% CI 0, 0.56%). The same meta-analysis found that the leading cause of TL fractures was motor vehicle collisions, accounting for

36.7% (95% CI 31.35%, 42.0%) of cases, followed by high-energy falls at 31.7% (95% CI 25%, 38.4%). Motorcycle collisions accounted for 10.05%, other causes for 9.06%, and pedestrian accidents for 4.83%. Muratore et al.¹¹ reported that fractures predominantly involved the TL region, between T11 and L2, while Neeley et al.¹⁵ found the highest incidence at the T12-L1 level. In contrast, our study identified falls as the most common cause of fractures, with the most frequent fracture levels occurring between T11 and L3.

Katsuura et al. identified type A3 or A4 fractures as the most prevalent trauma patterns.²¹ Similarly, Reinhold et al. demonstrated that type A fractures were the most common, with over half of their patients presenting with polytrauma.¹⁴ Their findings also revealed an increasing incidence of neurological deficits as the fracture severity escalated from type A to type C. Neeley et al. also observed a predominance of type A fractures.¹⁵ In our study, type A fractures were the most frequently encountered. In addition, post-operative neurodeficits in our cohort exhibited a comparable trend, with significantly higher rates observed in type B (41.7%) and type C (37.5%) fractures compared to type A fractures (5.7%).

In their meta-analysis, Azizi et al. reported a pooled spinal cord injury (SCI) event rate of 15.81% among patients with TL fractures.²² Similarly, Muratore et al. found that the probability of medullary involvement increased with the severity of the primary fracture type, showing a significant difference between type A and type C fractures, but not between type A and type B.¹¹ Katsuura et al. conducted a meta-analysis involving seven studies, which included 3,146 patients, to assess the rate of neurologic SCI in TL fracture patients.²¹ One study, which differentiated between a polytrauma group and a specific TL trauma group, was analyzed as two separate cohorts.²³ The overall rate of SCI in these patients was found to be 26.5% (95% CI 15.8%, 37.2%).²¹ In our study, neurological deficits were observed in 18.2% of the patients. Furthermore, univariate analysis revealed that the presence of pre-operative neurodeficits and the TL AOSIS type were significantly associated with the occurrence of post-operative neurodeficits.

While this study provides valuable insights into the surgical outcomes of traumatic TL fractures, it is not without limitations. First, the retrospective nature of the study introduces potential biases, including selection and recall bias, which may affect the accuracy of the data. Second, the relatively small sample size limits the generalizability of the findings, particularly

when stratifying patients by injury type and analyzing subgroup outcomes. The study did not account for potential confounding factors such as the patients' comorbidities, the timing of surgery, and variations in surgical techniques, all of which could influence the outcomes. In addition, the follow-up period, though adequate, varied among patients, which might have affected the assessment of long-term outcomes such as late-onset complications and functional recovery. Finally, the lack of randomization and a control group limits the ability to establish causality between the interventions and observed outcomes.

Conclusions

This study, therefore, takes into consideration the strong impact of pre-operative neurological deficit and injury severity with respect to TL classifications on post-operative outcomes for patients undergoing surgical stabilization for traumatic fractures. Patients who have pre-operative neurological deficits and those whose injury types are more serious, the B and C types, have a higher likelihood of having post-operative neurological deficits. This study's results bring out the importance of early and accurate assessment of injuries, further pinpointing the need for individualized surgical approaches to improve patient outcomes. However, more prospective studies on larger populations are needed to confirm these observations and sharpen clinical decision-making.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of human subjects and animals. The authors declare that the procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation and with the World Medical Association and the Declaration of Helsinki. The procedures were authorized by the Institutional Ethics Committee.

Confidentiality, informed consent, and ethical approval. The authors have followed their institution's

confidentiality protocols, obtained informed consent from all patients, and secured approval from the Ethics Committee. SAGER guidelines have been followed as applicable to the nature of the study. Ethical approval was obtained from the Sancaktepe local ethical committee.

Declaration on the use of artificial intelligence.

The authors declare that no generative artificial intelligence was used in the writing or creation of the content of this manuscript.

References

1. Azam MQ, Sadat-Ali M. The concept of evolution of thoracolumbar fracture classifications helps in surgical decisions. *Asian Spine J.* 2015;9:984-94.
2. Haq MI, Khan SA, Aurangzeb A, Ahmed E, Bhatti SN, Noman A. Radiological outcome of transpedicular screws fixation in the management of thoracolumbar spine injury. *J Ayub Med Coll Abbottabad.* 2015;27:171-3.
3. Cengiz SL, Kalkan E, Bayir A, Ilik K, Basefer A. Timing of thoracolumbar spine stabilization in trauma patients; Impact on neurological outcome and clinical course. A real prospective (rct) randomized controlled study. *Arch Orthop Trauma Surg.* 2008;128:959-66.
4. Hatakeyama Y, Hongo M, Kido T, Urayama M, Kasukawa Y, Sasaki H, et al. Neurological analysis based on the terminal end of the spinal cord and the narrowest level of injured spine in thoracolumbar spinal injuries. *Acta Med Okayama.* 2023;77:499-509.
5. Rodríguez-Morales J, Méndez-Viveros A, Pineda-Hernández C, Parra-Romero G, Ariñez-Barahona E, Guartazaca-Guerrero S, et al. Incidence of neurosurgical pathology of the spine in population of Mexico City. *Cir Cir.* 2021;89:806-10.
6. Ruiz Santiago F, Tomás Muñoz P, Moya Sánchez E, Revelles Paniza M, Martínez Martínez A, Pérez Abela AL. Classifying thoracolumbar fractures: role of quantitative imaging. *Quant Imaging Med Surg.* 2016;6:772-84.
7. Reinhold M, Knop C, Beisse R, Audigé L, Kandziora F, Pizanis A, et al. [Operative treatment of traumatic fractures of the thoracic and lumbar spinal column. Part I: epidemiology]. *Unfallchirurg.* 2009;112:33-42, 44-5.
8. Durel R, Rudman E, Milburn J. Clinical images - a quarterly column: chance fracture of the lumbar spine. *Ochsner J.* 2014;14:9-11.
9. Smith PP, Hurtado EA, Appell RA. *Post hoc* interpretation of urodynamic evaluation is qualitatively different than interpretation at the time of urodynamic study. *NeuroUrol Urodyn.* 2009;28:998-1002.
10. Schömig F, Taheri N, Kalaf H, Muellner M, Becker L, Pumberger M. Validation of the TLICS and AOSpine injury score for surgical management of paediatric traumatic spinal injuries. *Arch Orthop Trauma Surg.* 2023;143:2011-7.
11. Muratore M, Allasia S, Viglierchio P, Abbate M, Aleotti S, Masse A, et al. Surgical treatment of traumatic thoracolumbar fractures: a retrospective review of 101 cases. *Musculoskelet Surg.* 2021;105:49-59.
12. Villarreal-García FI, Martínez-Gutiérrez OA, Reyes-Fernández PM, Saavedra-Badillo LA, Avalos RM, Acosta-Olivo CA, et al. Two-year prevalence of spinal gunshot injuries in Mexico: a single center experience. *Cir Cir.* 2022;90:467-72.
13. Zhang C, Liu Y. Combined pedicle screw fixation at the fracture vertebrae versus conventional method for thoracolumbar fractures: a meta-analysis. *Int J Surg.* 2018;53:38-47.
14. Reinhold M, Knop C, Beisse R, Audigé L, Kandziora F, Pizanis A, et al. Operative treatment of 733 patients with acute thoracolumbar spinal injuries: comprehensive results from the second, prospective, internet-based multicenter study of the Spine study group of the German association of trauma surgery. *Eur Spine J.* 2010;19:1657-76.
15. Neeley OJ, Kafka B, Tecle NE, Shi C, El Ahmadi TY, Sagoo NS, et al. Percutaneous screw fixation versus open fusion for the treatment of traumatic thoracolumbar fractures: a retrospective case series of 185 patients with a single-level spinal column injury. *J Clin Neurosci.* 2022;101:47-51.
16. Vaccaro AR, Oner C, Kepler CK, Dvorak M, Schnake K, Bellabarba C, et al. AOSpine thoracolumbar spine injury classification system: fracture description, neurological status, and key modifiers. *Spine (Phila Pa 1976).* 2013;38:2028-37.
17. Jaafarpour M, Hatefi M, Vasigh A, Khani A, Njafi F, Shafiei E. Comparison between general and spinal anesthesia for lumbar disc surgery: a randomized clinical trial. *Acad J Health Sci Med Balear.* 2023;38:41-6.
18. Qureshi S, Dhall SS, Anderson PA, Arnold PM, Chi JH, Dailey AT, et al. Congress of neurological surgeons systematic review and evidence-based guidelines on the evaluation and treatment of patients with thoracolumbar spine trauma: radiological evaluation. *Neurosurgery.* 2019;84:E28-31.
19. Ghobrial GM, Jallo J. Thoracolumbar spine trauma: review of the evidence. *J Neurosurg Sci.* 2013;57:115-22.
20. Gertzbein SD. Scoliosis research society. Multicenter spine fracture study. *Spine (Phila Pa 1976).* 1992;17:528-40.
21. Katsuura Y, Osborn JM, Cason GW. The epidemiology of thoracolumbar trauma: a meta-analysis. *J Orthop.* 2016;13:383-8.
22. Azizi A, Azizzadeh A, Tavakoli Y, Vahed N, Mousavi T. Thoracolumbar fracture and spinal cord injury in blunt trauma: a systematic review, meta-analysis, and meta-regression. *Neurosurg Rev.* 2024;47:333.
23. Hsu JM, Joseph T, Ellis AM. Thoracolumbar fracture in blunt trauma patients: guidelines for diagnosis and imaging. *Injury.* 2003;34:426-33.

Arthroscopic treatment with absorbable bone cement for intra-articular comminuted fracture of the distal radius

Tratamiento artroscópico con cemento óseo reabsorbible para la fractura intraarticular conminuta del radio distal

Bangxian Dou^{1#}, Huiling Tian^{2#}, and Xuedong Li^{1*}

¹Department of Hand Surgery, Zhengzhou Orthopedic Hospital; ²School of Basic Medicine, Zhengzhou Railway Vocational and Technical College, Zhengzhou, China.

[#]These authors contributed equally and share first authorship.

Abstract

Objective: The objective of this study was to explore the clinical application value of arthroscopic injection of absorbable calcium sulfate bone cement in the treatment of distal radius intra-articular comminuted fractures. **Methods:** A total of 15 patients with distal radius intra-articular comminuted fractures treated in the Department of Hand Surgery I of our hospital from January 2020 to October 2021 were selected as the research subjects, including nine males and six females, aged 25-55. Closed reduction was performed first, followed by arthroscopic-assisted fracture reduction to eliminate step-off caused by the fracture and to inspect and repair damaged ligaments. Absorbable bone cement was injected into the fracture gap under fluoroscopy or arthroscopic guidance. The Mayo wrist score and Gartland-Werley score were used to assess wrist pain and joint function of the patients. **Results:** In the latest follow-up, 12 cases achieved excellent results in the Mayo score, and three cases achieved good results; Gartland-Werley score: 11 cases were excellent, and 4 cases were good. **Conclusions:** Arthroscopic injection of calcium sulfate bone cement for distal radius intra-articular comminuted fractures is effective, with the advantages of minimal trauma, satisfactory reduction, fewer post-operative complications, and the ability for early functional exercise in patients.

Keywords: Wrist arthroscopy. Calcium sulfate bone cement. Distal radius intra-articular comminuted fracture. Traumatic arthritis.

Resumen

Objetivo: Explorar el valor clínico de la inyección artroscópica de cemento de sulfato de calcio reabsorbible en el tratamiento de la fractura conminuta intraarticular del radio distal. **Métodos:** Se seleccionaron 15 pacientes (9 hombres y 6 mujeres), de entre 25 y 55 años, con fractura conminuta intraarticular del radio distal tratados en nuestro hospital entre enero de 2020 y octubre de 2021. Primero se realiza una reducción cerrada y luego una reducción de fractura asistida por artroscopia para eliminar la dislocación causada por la fractura, examinar y reparar los ligamentos dañados. Bajo guía de fluoroscopia o artroscopia, el cemento óseo reabsorbible se inyecta en el espacio de fractura. Para evaluar el dolor de muñeca y la función articular del paciente se utilizaron la puntuación Mayo de muñeca y la puntuación de Gartland y Werley. **Resultados:** En el seguimiento reciente, 12 pacientes alcanzaron una excelente puntuación Mayo y 3 alcanzaron una buena puntuación de Gartland y Werley: 11 casos excelentes y 4 casos buenos. **Conclusiones:** La inyección de cemento óseo de sulfato de calcio bajo artroscopia es efectiva en el tratamiento de fracturas conminutas en la articulación del radio distal, con pocas complicaciones posoperatorias y ejercicio funcional temprano en los pacientes.

Palabras clave: Artroscopia de muñeca. Cemento de sulfato de calcio. Fractura conminuta intraarticular del radio distal. Artritis traumática.

*Correspondence:

Xuedong Li

E-mail: 912001079@qq.com

2444-0507/© 2025 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 21-09-2024

Date of acceptance: 28-03-2025

DOI: 10.24875/CIRUE.M25000942

Cir Cir (Eng). 2026;94(3):248-254

Contents available at PubMed

www.cirugiaycirujanos.com

Introduction

Distal radius fractures are the most common fractures in adults, with an incidence rate of 0.35%, and this rate is showing an upward trend.¹ These fractures are typically caused by high-energy trauma and can manifest as sheare, compression, burst, or avulsion fractures. In addition to osseous structural damage, they are often accompanied by carpal bone fractures.² Non-surgical treatment is suitable for stable fractures, whereas displaced or unstable fractures usually require surgical intervention to restore upper limb alignment, functionality, and prevent post-traumatic arthritis.^{3,4} Currently, the use of the volar approach and volar locking plates for treating distal radius fractures has been widely accepted. Although it can address most extra-articular and simple intra-articular fractures, treating comminuted intra-articular fractures, such as AO type C fractures, remains a challenge in surgery. Harness pointed out that the unique anatomical structure of the distal radius fracture area with volar bone fragments makes it challenging to support the entire volar surface using standard plates for fixation, due to the proximity of the fracture line to the joint surface, presenting treatment difficulties.⁵ For elderly individuals with accompanying osteoporosis, bone deficiency is also a factor that must be considered. Moreover, distal radius intra-articular fractures often involve damage to ligaments and other structures. Without timely intervention, post-operative chronic wrist pain or wrist instability might persist.⁶ How to reduce intra-articular comminuted fractures and repair ligament damage are critical aspects of surgically treating intra-articular fractures, and the emphasis on anatomical reduction and early functional exercises in the context of accelerated recovery concepts places higher demands on surgical techniques.^{7,8} From January 2020 to October 2021, our department utilized wrist arthroscopy in combination with calcium sulfate bone cement for the treatment of 15 patients with distal radius fractures and intra-articular comminuted fractures. The results are reported as follows.

Methods

General information

Between January 2020 and October 2021, 45 patients underwent surgery for distal radius

intra-articular comminuted fractures in our trauma center. Retrospective study subjects were identified according to inclusion and exclusion criteria, and 15 patients were included in the study.

INCLUSION CRITERIA

- Age 18-65 years and met the diagnostic criteria for a distal radius fracture;
- AO type C fractures;
- Patient's informed consent and signed informed consent;
- Patients were treated with arthroscopic treatment with absorbable bone cement;
- A minimum follow-up of 12 months after surgery.

EXCLUDED PATIENTS

- Patients with a history of wrist joint surgery;
- Lost to follow-up.

There were 9 males and 6 females, aged between 40 and 65 years. Causes of injury included falls in 12 cases and collisions in 3 cases. The fractures were classified according to the AO classification as follows: C1 in 4 cases, C2 in 10 cases, and C3 in 1 case. This study was approved by the Medical Research Ethics Board of blinded for review. All patients were informed in detail before surgery, and consent was obtained from both the patients and their families. All surgeries were performed by the author.

Surgical procedure

Patients were administered brachial plexus nerve block anesthesia and placed in a supine position. Closed reduction was performed initially, with traction and compression to achieve closed reduction, aiming to restore radial height and avoid ulnar deviation. Fracture alignment was corrected, and palmar tilt, ulnar variance, and rotational deformity were restored. The affected limb was placed in a wrist arthroscopy tower, providing 3-4 kg of traction force. Dry arthroscopy was used intraoperatively. Arthroscopy was used for exploration, primarily involving intra-articular fracture reduction to eliminate step-off caused by the fracture. Subsequently, the integrity of the triangular fibrocartilage complex (TFCC) and the lunate ligament was further examined. Any injuries were repaired under arthroscopy. Passive wrist joint movement and




Figure 1. **A:** patient, female, 34 years old, suffered from a left distal radius intra-articular fracture due to violence. The fracture was classified as AO type C3 with joint surface depression. **B:** under direct vision of wrist arthroscopy, articular surface fractures with step formation, triangular fibrocartilage complex (TFCC) injury, probe prying reduction, Kirschner wire fixation, reduction of articular surface, and repair of TFCC can be seen. **C:** post-operative fluoroscopy shows anatomical reduction of the joint surface, correction of ulnar variance angle, palmar tilt angle, and radial height. Injectable absorbable bone cement was injected into the fracture gap under fluoroscopic guidance. **D:** 6 weeks after the operation, an X-ray of the wrist joint was taken. The fracture has basically healed, the articular surface has not collapsed, and the bone cement has been partially absorbed.

ulnar-radial and radial-ulnar collision tests were performed during the surgery to assess joint stability. A Kirschner wire or cap screw was used for temporary intraoperative fixation. Once satisfactory exploration was achieved, the wrist joint was positioned under a fluoroscopy machine, and a needle was inserted into the fractured gap under fluoroscopy or arthroscopic guidance. Absorbable injectable calcium sulfate bone cement (STIMULAN, BIOCOSMOS, UK) was prepared with a ratio of 2 g calcium sulfate bone cement to 5 mL of normal saline using a 5 mL syringe. The cement was injected immediately and allowed to naturally solidify, with a solidification time of about 3 min. Postoperatively, the wrist joint was carefully immobilized in a plaster functional position.

Post-operative management and efficacy evaluation

Active and passive functional exercises for interphalangeal and metacarpophalangeal joints were initiated on the first day after surgery. At 3 weeks post-surgery, the plaster was removed, and patients were guided to perform passive wrist flexion-extension training. Resistance and weight-bearing training began at 4 weeks post-surgery. The Kirschner wire was pulled out at 4 weeks post-surgery. These patients began resistance and weight-bearing training at 6 weeks post-surgery.

Follow-up was conducted for at least 1 year. X-ray evaluation was performed to assess fracture healing and the occurrence of arthritis. General wrist conditions were assessed, including stability, range of motion, tendon irritation, wrist pain, and other

complications. The Mayo wrist score and Gartland-Werley score were used to assess wrist pain and joint function.

Results

The average time from injury to operation was 3.5 days (range 2-7 days). All 15 patients in this group were successfully followed up, with a follow-up period ranging from 12 to 23 months, averaging 14.5 ± 4.1 months (Fig. 1). Two patients exhibited TFCC grade I injuries, which were cleaned and repaired intraoperatively. Two patients presented with injuries to the lunotriquetral ligament, which were managed through intraoperative fixation of the lunotriquetral joint. One patient experienced radial nerve symptoms 1 week after surgery, which disappeared after oral administration of methylcobalamin. Minimal cement leakage occurred in one case, requiring no specific treatment. All patients achieved anatomical restoration of the radiocarpal, radioulnar, and distal radioulnar joints. No complications such as infection, non-union, carpal tunnel syndrome, ulnar impaction syndrome, cement intolerance, or tendon rupture were observed.

Fracture healing time ranged from 8 to 15 weeks, with an average of 12.2 ± 2.7 weeks. The bone cement gradually absorbed around 4 weeks after surgery, with complete absorption occurring between 10 and 13 weeks, averaging 13.5 ± 1.9 weeks. At the latest follow-up, all patients exhibited stable wrist joints. Range of motion values were as follows: extension $36-45^\circ$, averaging $38 \pm 3.5^\circ$; flexion $40-52^\circ$, averaging $45.2 \pm 4.1^\circ$; pronation $65-80^\circ$, averaging $75.3 \pm 5.3^\circ$;

supination 60-77°, averaging $69.1 \pm 4.9^\circ$; palmar tilt angle 9-17°, averaging $12.5 \pm 1.7^\circ$; ulnar variance angle 15-27°, averaging $22.5 \pm 3.2^\circ$. In the latest follow-up, 12 cases achieved excellent results in the Mayo score, and 3 cases achieved good results. The Gartland-Werley score showed excellent results in 11 cases and good results in 4 cases.

Discussion

The distal radial articular surface is composed of different combinations of fracture fragments, including the radial column fragment, volar lip fragment of the lunate fossa, ulnar dorsal corner fragment, dorsal wall fragment, and intra-articular fragment. The specific location affected by distal radial fractures corresponds to the arrangement of these five fracture fragments. In addition, soft-tissue structures such as the radial collateral ligament, TFCC, and extensor carpi radialis brevis tendon are often damaged during traumatic events.^{9,10} Achieving anatomical reduction is a treatment principle that can significantly reduce the risk of adverse effects such as traumatic arthritis.

Commonly used distal radius plates are volar plates. Soong reported that placing the distal radius plate at a 2nd position is necessary to fix fractures beyond the watershed line.¹¹ Traditional open surgery entails greater disruption and often requires incisions that sever the pronator quadratus muscle and involve traction of the median nerve. Fixation of dorsal fracture fragments may necessitate additional dorsal incisions, and exploring internal wrist joint structures can be challenging, particularly in cases of TFCC damage and injuries to the lunotriquetral ligament.¹²⁻¹⁴

The technique of arthroscopic-assisted injection of absorbable bone cement for fixation offers advantages such as minimal trauma, excellent outcomes, rapid recovery, and reduced discomfort.¹⁵ Following reduction, step-off and fracture gaps on the joint surface should not exceed 1mm. Otherwise, residual wrist pain or functional impairment may occur postoperatively. However, it has been observed that fluoroscopy during surgery is often inadequate in detecting step-offs or gaps of around 1 mm.¹⁵ Arthroscopy provides a direct view of fractures and soft-tissue injuries, allowing for direct repair of damaged joint surfaces and soft tissue structures. Yin et al.¹⁰ used arthroscopy-assisted Kirschner wire fixation in combination with external fixation for treating fractures of the extreme distal radius in 21 patients. Intraoperative arthroscopy was utilized for initial exploration of the

joint space, revealing TFCC tears in six cases. Arthroscopy can repair joint soft-tissue injuries and effectively address wrist joint dysfunction and persistent pain caused by soft-tissue damage.¹⁰ In our study, two patients experienced TFCC injuries, and two patients had lunotriquetral ligament injuries, all of which were promptly repaired during surgery. Post-operative recovery met expectations, and no significant wrist pain was observed. Arthroscopy facilitates the removal of impurities such as blood clots and bone fragments from gaps while emphasizing the protection of damaged soft tissues. Anatomical reduction of the wrist joint surface was pursued during surgery, and arthroscopy holds unique advantages for certain fractures, such as those within the lunate fossa, dorsal wall fragments, and distal radioulnar joint fractures. In our procedures, we performed closed reduction, followed by meticulous arthroscopic repair and bone cement injection, striving to achieve maximal anatomical reduction. Adhesive fixation with bone cement can reduce the invasiveness of open surgery, aligning more closely with surgical principles¹⁶ established in the field.

Subarticular bone defects often require bone grafting. Calcium sulfate bone cement material has good biocompatibility, degradability, and osteogenesis, and is widely used in the treatment of bone defects caused by vertebral compression fractures and trauma.¹⁷ Calcium sulfate can produce a local micro-acid environment *in vivo*, which can increase the local calcium ion concentration, which is conducive to the aggregation and growth of osteoblasts. During the degradation and absorption, a large number of new bone trabeculae are formed, and the undegraded bone cement acts as a bone conduction scaffold. Bone cement absorption and bone tissue formation are synchronous, and the degradation and absorption rate of calcium sulfate is close to the rate of new bone formation.¹⁸ Previous studies have shown that calcium sulfate bone cement can allow early wrist joint movement after the treatment of distal radius fractures, which can avoid wrist pain, stiffness, traumatic arthritis, and muscle atrophy caused by post-operative fixation, and has a good effect on comminuted fractures of the distal radius, unstable fractures, and osteoporotic fractures in the elderly. In addition, muscle contraction during early post-operative activity can produce axial stress at the fracture end, which can promote bone healing.¹⁸ In this study, the minimally invasive injection method can effectively avoid periosteal injury and facilitate fracture healing. Calcium sulfate cement is used to fill bone

defects, which is an important factor in restoring the articular surface and maintaining radial height. Calcium sulfate is acidic, which can increase the local calcium ion concentration and the activity of osteoblasts. At the same time, bone cement absorption and bone tissue formation are synchronized, and the degradation and absorption rate of calcium sulfate is close to the rate of new bone formation.¹⁹ In this group of cases, the follow-up found that a small amount of bone cement leakage occurred in one case; without special treatment, the bone cement was gradually absorbed after about 4 weeks, and the average time of fracture healing was 13.5 ± 1.9 weeks. The fracture healing time was 8-15 weeks, the average time of fracture healing was 12.2 weeks, and the bone cement absorption and healing time were almost the same, which was basically consistent with the previous reports of healing time. This suggests that cement may not increase the risk of delayed union. The absorbability of bone cement could avoid secondary surgery, and all patients had no rejection reaction. After an average follow-up of 14.5 months, all patients had no chronic wrist pain or wrist arthritis, and wrist joint function recovered well. The authors used manual reduction, wrist arthroscopy, and the use of bone cement for injection to obtain reliable clinical results. Under the arthroscopic, the author examined and repaired TFCC injury, and used bone cement to repair the intra-articular steps and fill the bone defect. The combination of the three can effectively restore the length of the radius, fill the bone defect, and address the problem of articular surface reduction. Subarticular bone defects often require bone grafting. Calcium sulfate bone cement material has good biocompatibility, degradability, and osteogenesis, and is widely used in the treatment of bone defects caused by vertebral compression fractures and trauma.¹⁷ Calcium sulfate can produce a local micro-acid environment *in vivo*, which can increase the local calcium ion concentration, which is conducive to the aggregation and growth of osteoblasts. During the degradation and absorption, a large number of new bone trabeculae are formed, and the undegraded bone cement acts as a bone conduction scaffold. Bone cement absorption and bone tissue formation are synchronous, and the degradation and absorption rate of calcium sulfate is close to the rate of new bone formation.¹⁸ Previous studies have shown that calcium sulfate bone cement can allow early wrist joint movement after the treatment of distal radius fractures, which can avoid wrist pain, stiffness, traumatic arthritis, and muscle atrophy

caused by post-operative fixation, and has a good effect on comminuted fractures of the distal radius, unstable fractures, and osteoporotic fractures in the elderly. In addition, muscle contraction during early post-operative activity can produce axial stress at the fracture end, which can promote bone healing.¹⁸ In this study, the minimally invasive injection method can effectively avoid periosteal injury and facilitate fracture healing. Calcium sulfate cement is used to fill bone defects, which is an important factor in restoring the articular surface and maintaining radial height. Calcium sulfate is acidic, which can increase the local calcium ion concentration and the activity of osteoblasts. At the same time, bone cement absorption and bone tissue formation are synchronized, and the degradation and absorption rate of calcium sulfate is close to the rate of new bone formation.¹⁹ In this group of cases, the follow-up found that a small amount of bone cement leakage occurred in one case; without special treatment, the bone cement was gradually absorbed after about 4 weeks, and the average time of fracture healing was 13.5 ± 1.9 weeks. The fracture healing time was 8-15 weeks, the average time of fracture healing was 12.2 weeks, and the bone cement absorption and healing time were almost the same, which was basically consistent with the previous reports of healing time. This suggests that cement may not increase the risk of delayed union. The absorbability of bone cement could avoid secondary surgery, and all patients had no rejection reaction. After an average follow-up of 14.5 months, all patients had no chronic wrist pain or wrist arthritis, and wrist joint function recovered well. The authors used manual reduction, wrist arthroscopy, and the use of bone cement for injection to obtain reliable clinical results. Under the microscope, the author examined and repaired the scapholunate ligament, TFCC injury, and other soft-tissue injuries, and used bone cement to repair the intra-articular steps and fill the bone defect. The combination of the three can effectively restore the length of the radius, fill the bone defect, and address the problem of articular surface reduction.

Early exercise is key to joint function recovery. Among the 15 cases in this group, active and passive flexion and extension exercises for the fingers and metacarpophalangeal joints were initiated 24 h post-operatively, whereas active wrist flexion, extension, and rotational exercises were started at 4 weeks post-operatively. No complications of refracture or deformity healing were observed. The surgical technique involving a plaster cast combined with injectable

calcium sulfate bone cement not only met the requirements for anatomical reduction of fractures but also fulfilled the need for early rehabilitation exercises. During the procedure, the following points should be noted: (a) for cases where the central column fracture fragment of the radius cannot be reduced by flipping, Kirschner wires should be used to lever the fragment into position under arthroscopy; (b) as a supplementary material, bone cement can adhere to small bone fragments that are difficult to fix or free-floating fragments, such as smaller Die-punch fractures; (c) for cases involving ulnar styloid fractures, TFCC injuries are often present. After reduction, exploration of the TFCC injury can be conducted.

This study has several limitations: a limited number of cases. During the initial exploratory stage of surgery, the operation time was longer, and surgical techniques were less mature, which may have influenced post-operative recovery and data statistics. In addition, all surgeries were performed by one team of doctors, potentially introducing subjective factors influencing therapeutic outcomes. There is limited research on the optimal concentration ratio and injection timing of absorbable bone cement, and there is a lack of quantitative biomechanical studies.

Conclusions

Minimally invasive wrist joint surgery aligns with the concept of rapid recovery. It features small incisions and minimal tissue damage while serving both diagnostic and therapeutic purposes. This approach is particularly effective for repositioning joints with unique anatomical structures. The technique of injectable absorbable calcium sulfate bone cement provides a new perspective for surgeons in choosing treatment options. The combination of these two approaches serves as a beneficial supplement to traditional treatments, offering clinicians new avenues for therapy. In summary, the use of wrist arthroscopy combined with injectable calcium sulfate bone cement for treating distal radius intra-articular fractures results in minimal trauma, satisfactory reduction, and fewer post-operative complications. It allows patients to engage in early flexion-extension exercises, facilitating quicker restoration of wrist joint function.

Funding

Zhengzhou Medical and Health Technology Innovation Guidance Project (2024YLZDJH180 /2025YLZDJH011).

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of humans and animals. The authors declare that the procedures followed complied with the ethical standards of the responsible human experimentation committee and adhered to the World Medical Association and the Declaration of Helsinki. The procedures were approved by the institutional Ethics Committee.

Confidentiality, informed consent, and ethical approval. The authors have followed their institution's confidentiality protocols, obtained informed consent from all patients, and secured approval from the Ethics Committee. SAGER guidelines have been followed as applicable to the nature of the study. This study was approved by the Medical Research Ethics Board of Zhengzhou Orthopedic Hospital.

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing or creation of the content of this manuscript.

References

1. Mauck BM, Swigler CW. Evidence-based review of distal radius fractures. *Orthop Clin North Am.* 2018;49:211-22.
2. Raudasoja L, Vastamäki H, Raatikainen T. The importance of radiological results in distal radius fracture operations: functional outcome after long-term (6.5 years) follow-up. *SAGE Open Med.* 2018;6:1-7.
3. Kreder HJ, Hanel DP, McKee M, Jupiter J, McGilivray G, Swiontkowski MF. X-ray film measurements for healed distal radius fractures. *J Hand Surg Am.* 1996;21:31-9.
4. Mosenthal WP, Boyajian HH, Ham SA, Conti Mica MA. Treatment trends, complications, and effects of comorbidities on distal radius fractures. *Hand (N Y).* 2019;14:534-9.
5. Harness NG, Jupiter JB, Orban JL, Raskin KB, Fernandez DL. Loss of fixation of the volar lunate facet fragment in fractures of the distal part of the radius. *J Bone Joint Surg Am.* 2004;86:1900-8.
6. Lameijer CM, Ten Duis HJ, Dusseldorp IV, Dijkstra PU, Van Der Sluis CK. Prevalence of posttraumatic arthritis and the association with outcome measures following distal radius fractures in non-osteoporotic patients: a systematic review. *Arch Orthop Trauma Surg.* 2017;137:1499-513.
7. Ljungqvist O, Scott M, Fearon KC. Enhanced recovery after surgery: a review. *JAMA Surg.* 2017;152:292-8.
8. Lichtman DM, Bindra RR, Boyer MI, Putnam MD, Ring D, Slutsky DJ, et al. Treatment of distal radius fractures. *J Am Acad Orthop Surg.* 2010;18:180-9.
9. Schmidt V, Gordon M, Petterson A, Buttazzoni C, Seimersson A, Sayed-Noor A, et al. Functional outcomes are restored a decade after a distal radius fracture: a prospective long-term follow-up study. *J Hand Surg Eur Vol.* 2024;49:322-8.
10. Yin F, Rui Y, Li F, Liu J, Wu Y, Lu Y, et al. Effectiveness of arthroscopy-assisted combined fixation of Kirschner wire and external fixator for treating extreme distal radial fractures. *Zhongguo Xiu Fu Chong Jian Wai Ke Za Zhi.* 2017;31(12):1414-1417.
11. Soong M, Earp BE, Bishop G, Leung A, Blazar P. Volar locking plate implant prominence and flexor tendon rupture. *J Bone Joint Surg Am.* 2011;93:328-35.
12. Zhang H, Liu M, Duan SY, Liang HR, Xu RD, Cai ZC. Surgical options for distal radius fractures of type C in elderly patients over 65 years old: a comparison of external fixation with Kirschner wires and volar locking plate. *J Orthop Surg Res.* 2023;18:669.

13. Lamas C, Arenas J, Almenara M, Rojas R, Fa-Binefa M, Toro-Aguilera A. Is pronator quadratus muscle repair required after anterior plate fixation for distal radial fractures? A prospective randomized comparative study. *J Hand Surg Eur Vol.* 2024;49:334-40.
14. Mulders MA, Walenkamp MM, Bos FJ, Schep NW, Goslings JC. Repair of the pronator quadratus after volar plate fixation in distal radius fractures: a systematic review. *Strategies Trauma Limb Reonstr.* 2017;12:181-8.
15. Karjalainen VL, Harris IA, Räisänen M, Karjalainen T. Minimal invasions: is wrist arthroscopy supported by evidence? A systematic review and meta-analysis. *Acta Orthop.* 2023;94:200-6.
16. Leunig M, Hertel R, Siebenrock KA, Ballmer FT, Mast JW, Ganz R. The evolution of indirect reduction techniques for the treatment of fractures. *Clin Orthop Relat Res.* 2000;375:7-14.
17. Winge MI, Røkkum M. CaP cement is equivalent to iliac bone graft in filling of large metaphyseal defects: 2 year prospective randomised study on distal radius osteotomies. *Injury.* 2018;49:636-43.
18. Wolfe SW, Austin G, Lorenze M, Swigart CR, Panjabi MM. A biomechanical comparison of different wrist external fixators with and without K-wire augmentation. *J Hand Surg Am.* 1999;24:516-24.
19. Peltier LF. The use of plaster of Paris to fill large defects in bone: a preliminary report. 1959. *Clin Orthop Relat Res.* 2001;382:3-5.

The effect of ethyl pyruvate administration on intestinal mucosal damage in neonatal rat models of necrotizing enterocolitis

Efecto de la administración de piruvato de etilo sobre el daño de la mucosa intestinal en modelos de enterocolitis necrosante en ratas neonatas

Tugce Merve-Orbay^{1*}, Aysenur Celayir¹, and Hulya Yavuz²

¹Department of Pediatric Surgery; ²Department of Pathology. University of Health Sciences, Istanbul Zeynep Kamil Maternity and Children's Diseases Health Training and Research Center, Istanbul, Turkey

Abstract

Objective: The purpose of this study was to determine whether intraperitoneal administration of ethyl pyruvate (EP) affects intestinal mucosal damage in neonatal rats with a model of necrotizing enterocolitis (NEC). **Methods:** The study comprised 46 Sprague-Dawley newborn rats delivered on the 21st gestational day. The rats were divided into four groups: Sham Group (n = 10), EP Group (n = 10), NEC Group (n = 13), and NEC + EP Group (n = 13). **Results:** The mean body weights of rats in Sham and EP groups significantly increased ($p < 0.05$), whereas those in NEC groups significantly decreased ($p < 0.05$). The changes were similar between groups ($p > 0.05$). There was no significant difference in the clinical sickness score (CSS) between groups ($p > 0.05$). The CSS in the NEC groups showed an increase from day 1 to day 4. Based on the histopathological grading, there was a statistically significant difference ($p < 0.001$) observed between the Sham Group, the NEC Group, and the NEC + EP Group. The differences in biochemical values among the other groups were found to be statistically significant ($p < 0.05$) for all variables. **Conclusions:** The application of EP in high-mortality diseases such as NEC is expected to reduce surgical needs, establishing a basis for future studies on EP as an innovative anti-inflammatory agent.

Keywords: Necrotizing enterocolitis. Neonate. Ethyl pyruvate. Interleukin 6. Malondialdehyde. Sprague-Dawley rat.

Resumen

Objetivo: Determinar si la administración intraperitoneal de piruvato de etilo (PE) afecta al daño de la mucosa intestinal en ratas neonatas con un modelo de enterocolitis necrosante (ECN). **Métodos:** El estudio incluyó 46 ratas recién nacidas Sprague-Dawley entregadas en el día gestacional 21. Las ratas se dividieron en cuatro grupos: grupo simulado (n = 10), grupo PE (n = 10), grupo ECN (n = 13) y grupo ECN + PE (n = 13). **Resultados:** El peso corporal medio de las ratas en los grupos simulado y PE aumentó significativamente ($p < 0.05$), mientras que en los grupos ECN disminuyó significativamente ($p < 0.05$). Los cambios fueron similares entre grupos ($p > 0.05$). No hubo diferencia significativa en el puntuación clínica de enfermedad (PCE) entre grupos ($p > 0.05$). El CSS en los grupos ECN mostró un aumento del día 1 al día 4. Según la clasificación histopatológica, se observó una diferencia estadísticamente significativa ($p < 0.001$) entre el grupo simulado, el grupo ECN y el grupo ECN + PE. Las diferencias en los valores bioquímicos entre los otros grupos resultaron ser estadísticamente significativas ($p < 0.05$) para todas las variables. **Conclusiones:** La aplicación de PE en enfermedades de alta mortalidad como la ECN se espera que reduzca las necesidades quirúrgicas, estableciendo una base para futuros estudios sobre el PE como un innovador agente antiinflamatorio.

Palabras clave: Enterocolitis necrotizante. Neonato. Piruvato de etilo. Interleucina 6. Malondialdehído. Rata Sprague-Dawley.

*Correspondence:

Tugce Merve-Orbay

E-mail: orbaymerve@gmail.com

Date of reception: 06-02-2024

Date of acceptance: 04-04-2024

DOI: 10.24875/CIRUE.M24000943

Cir Cir (Eng). 2026;94(3):255-263

Contents available at PubMed

www.cirugiaycirujanos.com

2444-0507/© 2024 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Necrotizing enterocolitis (NEC) is a pathological condition characterized by inflammation of the intestines that primarily affects neonates. It is sometimes observed in conjunction with sepsis, leading to the occurrence of intestinal perforation, peritonitis, and ultimately, mortality. The etiology of NEC encompasses various factors, including hypoxia, hyperosmolar enteral nutrition, intestinal immaturity, and intestinal dysbiosis.¹ Hypoxia and disruption of the intestinal epithelial barrier due to failure to provide enteral nutrition play an important role in premature infants.²

Studies in the past have shown that ethyl pyruvate (EP) has anti-inflammatory benefits.^{3,4} EP is a derivative of pyruvic acid and serves as an intermediary compound in the process of energy metabolism. Under anaerobic conditions, it facilitates the conversion to lactic acid through the process of dehydrogenation, thereby contributing to energy generation. Consequently, it exhibits antioxidant properties and diminishes the concentration of free radicals.⁵ EP is a chemical compound that serves as a derivative of pyruvic acid. It functions as an intermediary molecule in the process of energy metabolism. In anaerobic conditions, it facilitates the conversion of pyruvate to lactic acid by dehydrogenation, thereby contributing to energy generation. Previous research has demonstrated that EP possesses therapeutic properties in the treatment of intra-abdominal sepsis and colitis by modulating intestinal permeability and fortifying the mucosal barrier systems. According to reports, EP has been found to possess a mitigating impact on intestinal inflammation through the inhibition of toll-like receptor 4 (TLR-4).⁶ The experimental colitis model has demonstrated that EP exhibits beneficial effects by suppressing the "T Helper 1" immune response.⁷ The study investigated the impact of *Escherichia coli* lipopolysaccharide-induced sepsis and septicemia on arterial hypotension in a shock model. The researchers found that the administration of EP solution during resuscitation in rats led to a reduction in arterial hypotension, resulting in prolonged survival. Furthermore, it was observed that the concentrations of interleukin-6 (IL-6) and nitrite/nitrate exhibited a decrease; however, the levels of IL-10 were found to be elevated.⁸⁻¹⁰

The applications of EP are currently being investigated in the experimental study phase, with administration occurring through various channels such as intravenous, intraperitoneal, and intrapulmonary methods.^{9,11-13}

There is no experimental research on the intraperitoneal administration of EP in NEC models in the literature.

The purpose of this study was to determine whether intraperitoneal administration of EP affects intestinal mucosal damage in neonatal rats with a model of NEC.

Methods

After the approval of the Acibadem University Experimental Animals Local Ethics Committee (January 08, 2020–2020/01), the study was approved by the Scientific Research Projects Unit in the University of Health Sciences, Turkey, which is in accordance with legislation for the protection of animals used for scientific purposes.

To create newborn rats, male and female Sprague-Dawley rats were housed together. If the veterinarian observed sperm in the vaginal smear between 08:30 and 09:30 am, this was considered the start of the 1st gestational day. The mother rats were provided with unlimited access to food at a standard room temperature of 24.5°C. They were kept in a controlled environment with a 12-h cycle of darkness and light.¹⁰ On the 21st gestational day, a total of 46 Sprague-Dawley rats were born by cesarean section and were randomly divided into four groups. The study consisted of four groups: Sham Group (n = 10), NEC Group (n = 13), EP Group (n = 10), and NEC + EP Group (n = 13). The rats in the sham group were allowed to stay with their mothers and were given unlimited access to food. To eliminate the protective effect of breast milk against NEC, the rats designated for the NEC model were isolated from their mother. They were then placed in an incubator set at a temperature of 32°C with 60% humidity.¹¹

Sham group

The newborn rats were immediately given to their mother after birth and began to be nourished with breast milk. A total of 0.02 mL of 0.9% NaCl (saline) was administered intraperitoneally into the left lower quadrant at 1:00 p.m. for the initial 4 days.

EP group

Newborn rats, who were being fed breast milk and staying close to their mother for the first 4 days after birth, were given 50 mg/kg of EP intraperitoneally. This was administered as a solution in 0.9% NaCl, with a dosage of 0.02 mL.

NEC group

The newborn rats were fed a hyperosmolar formula consisting of 15 g of formula mixed with 75 mL of puppy or canine milk. This mixture was administered every 4 h using a 24 Gauge orogastric catheter, starting from the 3rd h. The feeding schedule was maintained every 4 h, with an increase of 0.1 mL/day in the subsequent days. The rats used in the NEC model experiment were subjected to specific conditions. They were exposed to hypoxia for a duration of 90 s inside a box filled with nitrogen gas flowing at a rate of 10 L/min. Following this, they were exposed to cold temperatures of 4 °C for a period of 10 min. In addition, the rats were placed in an incubator twice a day, specifically at 9:00 a.m. and 5:00 p.m., for a total of 3 days. A total of 0.02 mL of 0.9% NaCl (saline) was administered intraperitoneally from the left lower quadrant at 1:00 pm for the initial 4 days.¹⁰

NEC + EP group

The initial steps were identical to those in Group C. During the initial 4 days, a dose of 50 mg/kg of EP, in the form of a solution in 0.9% NaCl, was administered intraperitoneally from the left lower quadrant at 1:00 pm.

Histopathological evaluation

Two double-blind observers evaluated the daily weight and clinical sickness scores (CSSs) of rats in all groups.¹⁴ On the 4th day, following hypothermic sedation with an ice pack and 200 mg/kg phenobarbital, the surgical site was cleaned with povidone-iodine. A 3 cm segment of the terminal ileum was extracted through a 2 cm median incision above the umbilicus. Subsequently, all rats were euthanized.

Evaluation of the macroscopic appearance of the intestines was made by double-blind observers according to the “Macroscopic Gut Score”¹² (Table 1). Histopathological evaluation is based on the “NEC Grading Score” after 10% formaldehyde fixation and Hematoxylin and Eosin staining (Intact villus Grade 0, villi deletion 1, mild villus damage 2, complete villus necrosis 3, and transmural necrosis 4) performed by pathologists. Grades 2, 3, and 4 were accepted as NEC.¹³

Tissue samples of 3 cm in length, obtained from the terminal ileum, were subjected to storage at a temperature of –80°C. Subsequently, these samples were homogenized in a phosphate-buffered saline solution at a weight-to-volume ratio of 1/10. The tissue

Table 1. Scoring system for macroscopic gut assessment

Macroscopic gut assessment			
Score	Gut consistency	Gut colour	Gut dilatation
0	Normal	Normal	Normal
1	Moderately Friable	Patchy discoloration	Patchy dilatation
2	Extremely Friable (jelly-like)	Extensive discoloration	Extensive dilatation

Macroscopic Gut assessment, from Zani et al.¹².

homogenates underwent centrifugation at a speed of 3000 revolutions/min (rpm) at a temperature of +4°C. Subsequently, the supernatants obtained were gathered and put into polypropylene tubes. The levels of IL-6 and Malondialdehyde (MDA) were quantified using the Rat-IL-6 Enzyme-Linked Immuno Sorbent Assay (ELISA) kit (Catalog Number E0135Ra) and the Rat-MDA ELISA kit (Catalog Number E0156Ra), respectively.^{14,15}

Statistical analysis

The data was documented using the Microsoft Office 2019 Pro-Excel software. The data were analyzed using the IBM Statistical Package for the Social Sciences 26.0 software (IBM, Armonk, New York, US). The data are presented as means and standard deviations, or as counts and percentages (n%). The Shapiro-Wilk test was used to assess the normality of the data distribution. To assess the distinctions between groups, analysis of variance tests and *post hoc* tests were employed. In order to examine temporal variance, the Friedman and Wilcoxon tests were utilized. The results were assessed using a 95% confidence interval and a statistical significance level of $p < 0.05$.

Results

Nine rats survived in both the sham group and the EP group. One rat from each group could not be found in the cage, leading to the assumption that the mother rat might have eaten the baby rat. Six of 13 rats in the NEC group were euthanized on day 3 due to deterioration in their general condition and increased clinical disease scores. One rat in the NEC + EP group was excluded from the study due to its unfortunate demise in the cage on the 2nd day. Three of them were sacrificed on the 3rd day and included in the study due to

Table 2. Clinical sickness score

Clinical sickness score				
Score	Appearance	Natural activity	Response to touch	Body color
0	Tonic and well hydrated	Moving normally in the cage	Alert (without stimulation)	Pink
1	Slimmer but still tonic and hydrated	Able to wriggle if put supine	Responding to mild stimulation	Pale (just at the extremities)
2	Skinny, floppy, and dehydrated	Not able to wriggle if put supine	Responding to vigorous stimulation	Pale (whole body)
3	Gasping and in agony	Not moving limbs and lying still	Unresponsive notwithstanding vigorous stimulation	Grey

deterioration in general condition and an increase in CSSs on the 3rd day (Table 2).

The mean body weights of rats in Sham and EP groups significantly increased ($p < 0.05$), while those in NEC groups significantly decreased ($p < 0.05$). The changes were similar between groups ($p > 0.05$) (Fig. 1).

There was no statistically significant difference between the groups in terms of the CSS values on the 1st day ($p > 0.05$). The CSS was consistent across all groups on day 1. The results showed that there was no significant difference between the groups on day 1 ($p > 0.05$), but there was a significant difference on days 2-4 ($p < 0.05$). There was no significant difference in the CSS between groups ($p > 0.05$). The CSS in the NEC groups showed an increase from day 1 to day 4 (Fig. 2).


The average macroscopic bowel assessment score for all groups of rats was 1.58. There was no significant difference in macroscopic bowel scoring between the Sham and the EP Group. However, there was a statistically significant difference between all other groups ($p < 0.05$). Figure 3 displays the macroscopic bowel appearances of the different groups.

During the histopathological examination, it was observed that 78% of the rats in the sham Group had grade 0 NEC, while the remaining 22% had grade 1 NEC. Within the NEC group, 7.7% of the rats exhibited grade 1, while another 7.7% displayed grade 2. The majority, 30.7%, had grade 3 NEC, and the highest proportion, 53.8%, exhibited grade 4 NEC. All rats in the EP Group exhibited no signs of NEC, as they were assigned a grade of 0. In the NEC+EP Group, the distribution was as follows: 1 in 8.4%, 2 in 33.3%, 3 in 33.3%, and 4 in 25%. The histopathological microscopic images are displayed in figure 4. Based on the histopathological grading, there was a statistically significant difference ($p < 0.001$) observed between the

Table 3. Histopathological changes

Group (n)	Grade 0	Grade 1	Grade 2	Grade 3	Grade 4
Sham (9)	7 (77.8)	2 (22.2)	0	0	0
EP (9)	9 (100)	0	0	0	0
NEC (9)	0	1 (7.69)	1 (7.69)	4 (30.7)	7 (53.8)
NEC+EP (12)	0	1 (8.3)	4 (33.3)	4 (33.3)	3 (25)

Statistically significant differences were observed between Sham vs NEC, Sham vs NEC + EP, NEC vs EP, and EP vs NEC + EP groups ($p < 0.001$). No statistically significant difference was observed between NEC and NEC + EP groups.
NEC: necrotizing enterocolitis; EP: ethyl pyruvate.

**Figure 1. Time change of mean body weights of the groups.**

Sham Group, the NEC group, and the NEC + EP Group. There was a statistically significant difference ($p < 0.001$) between the NEC Group and the EP Group. A statistically significant difference ($p < 0.001$) was observed between the EP and NEC + EP Group. There was no statistically significant difference observed in the histopathological evaluation between the NEC group and the NEC + EP group. However, when considering the distribution, it was observed that the percentage of grade 3 and 4 rats in the treatment group was lower compared to the untreated NEC group. Table 3 presents the statistical results pertaining to the differences between groups.




Figure 2. Change of clinical sickness scores over time.




Figure 3. Macroscopic bowel appearance. (A) Sham group, (B) necrotizing enterocolitis (NEC) group, (C) ethyl pyruvate (EP) group, (D) NEC + EP group.




Figure 4. Histopathological changes. (A) Sham (Grade 1), (B) necrotizing enterocolitis (NEC) (Grade 3), (C) ethyl pyruvate (EP) (Grade 0), (D) NEC + EP (Grade 1).

There was no statistically significant difference in the biochemical values of the Sham Group and the EP Group when comparing the results of IL-6 level and tissue MDA level. The differences in biochemical values among the other groups were found to be

Table 4. Cytokine levels of groups

Cytokines	Sham (n = 9)	EP (n = 9)	NEC (n = 9)	NEC+EP (n = 12)	p
IL-6 (pg/mg tissue)	0.21 ± 0.02	0.21 ± 0.02	0.43 ± 0.06 ^a	0.28 ± 0.05 ^a	0.0001
MDA (pg/mg tissue)	0.06 ± 0.01	0.05 ± 0.01	0.12 ± 0.02 ^a	0.08 ± 0.02 ^a	0.0001

^aStatistically significant difference between group NEC versus NEC + EP p = 0.001.

NEC: necrotizing enterocolitis; EP: ethyl pyruvate; IL-6: interleukin-6; MDA: malondialdehyde.




Figure 5. Comparison of the cytokine levels.

statistically significant ($p < 0.05$) for all variables (Table 4 and Fig. 5).

Discussion

The advancement of neonatal intensive care units has highlighted NEC as a major concern due to its high rates of illness and death in premature infants. In preventing NEC, administering EP intraperitoneally may play a key role in lessening intestinal harm by reducing intestinal necrosis. This study is the first to examine the impact of EP on neonates with an NEC model. Our research aimed to shed light on NEC's causes and development, as well as on strategies for its prevention. The insights gained from this study lay the groundwork for further research in this field.

Different animal models have been developed for NEC research, but they vary in mechanism and pathophysiology. The human and animal immune systems

differ; for example, humans get immunoglobulins via the placenta, while pigs get them through the gastrointestinal tract. Barlow and Santulli's 1974 NEC model, the first successful one, focuses on altered intestinal bacteria and the lack of breast milk. Its creation involved formula feeding, hypoxia, and cold exposure.¹⁶

Numerous agents have undergone investigation in experimental animal models to explore their potential for preventing and treating NEC.¹⁷⁻¹⁹ EP is a pyruvic acid derivative and has been reported to provide improvement in conditions such as ischemia-reperfusion injury, organ dysfunctions, hemorrhagic shock, sepsis, acute respiratory distress syndrome, acute pancreatitis, and skin burns.⁹ EP has the ability to reduce the action of peritoneal macrophages by decreasing the secretion of HMGB1 protein.⁴ Previous studies have demonstrated that EP possesses therapeutic effects in the treatment of intra-abdominal sepsis and colitis by targeting the intestinal permeability

and mucosal barrier systems.^{6,7} In the colitis model induced by 2,4,6-trinitrobenzene sulfonic acid, it has been demonstrated that intraperitoneal administration of EP at a dose of 50 mg/kg for 1 week improves pathological damage and reduces T-helper-17 cells in the local tissue.⁷

Yurttutan et al.'s study showed newborn rats with an NEC model gained more weight with etanercept, a tumor necrosis factor alpha (TNF- α) blocker.¹⁷ Another study found increased weight in NEC rats fed growth factor-enriched formula.¹⁵ Our study observed normal growth in breastfed control groups and significant weight loss in NEC groups, with no difference in weight gain from intraperitoneal EP application.

The CSS was used to assess the clinical condition of newborn rats with NEC.¹⁴ Korkut et al.'s study showed lower scores in rats treated with obestatin.¹¹ Another study found improvement with mesenchymal stem cell therapy.¹⁸ Our research observed worsening conditions and increased scores in the NEC and NEC + EP groups, with significant deterioration by the 4th day. No difference in scores was noted between these groups, aligning with high morbidity and mortality rates in NEC models.

The macroscopic gut assessment method evaluates bowel appearance in the NEC model newborn rats.¹⁴ A study found that cytidine 50-diphosphocholine significantly reduced macroscopic bowel scores in treated rats.²⁰ Our study, using double-blind observers, showed that EP treatment in NEC rats led to improved bowel appearance and lower scores, indicating potential reduction in intestinal damage.

A grading system has been established to determine the histopathological diagnosis of NEC. The histopathological grading is assessed using the NEC grading score. This scoring system assigns a value of 0 for intact villi, 1 for villi deletion, 2 for mild villus damage, 3 for complete villus necrosis, and 4 for transmural necrosis.¹³ In a study performed by giving pentoxifylline to newborn rats, for which the NEC model was created, it was shown that histopathological NEC findings were milder.²¹ Previous studies similar to ours have reported a morbidity rate of 75% in the experimental NEC model.¹⁰ A study was conducted to investigate the effects of pioglitazone, a peroxisome proliferator-activated receptor- γ agonist, on an NEC model. The study compared a control group with a treatment group. The results showed that the treated group had a lower macroscopic bowel score and exhibited milder intestinal damage in histopathological examinations. The study found that there was no

difference between the two groups in terms of the amount of IL-6 and TNF- α messenger RNA (mRNA) in the tissue.¹⁷ In our study, we observed histopathological findings in newborn rats. Specifically, we found that 53% of rats in the NEC Group had grade 4 NEC, whereas 30% had Grade 3 NEC. In the group treated with NEC+EP, the distribution of rat grades was as follows: 33% were Grade 2, 33% were Grade 3, and 25% were Grade 4 NEC. There was no statistically significant difference between the two groups; distributionally, there was a decrease in the severity of NEC in the NEC + EP Group. In grades 3 and 4, NEC was detected in 83% of the rats in the NEC group. There was a significant difference between the control groups and the NEC groups in relation to the increase in histopathological grading. There was no statistically significant difference observed in histopathological grading between the NEC group and the NEC + EP group. However, the percentage of rats in the treatment group with Grade 3 and 4 distributions was lower compared to the untreated NEC group. In order to achieve statistical significance, it may be necessary to increase the sample size. The results indicate that the therapeutic effect of EP on NEC can be observed through intraperitoneal administration, as seen in the histopathological findings.

NEC is linked with increased levels of IL-6, a proinflammatory cytokine, which correlates with NEC severity. IL-6 stimulates C-reactive protein release and is regulated by nuclear factor kappa-light-chain-enhancer of activated B cells.²² Studies in rats show that interventions can impact IL-6 levels.^{23,24} Fecal microbiota transplantation normalized systemic IL-6, IL-1 β , and TNF- α in a Wistar albino rat NEC model. Another study using an NEC model induced by formula feeding, lipopolysaccharide injection, and hypoxia found that subcutaneously administered glucagon-like peptide-2 significantly lowered TNF- α and IL-6 levels.²⁵ Ulinastatin treatment also significantly decreased tissue IL-6 levels.²⁶ N-Acetyl Cysteine given antenatally and postnatally reduced IL-6 and its mRNA expression in intestinal tissue.^{27,28} Our study focused on the effect of EP on IL-6 levels in NEC rats. While control groups showed no significant IL-6 difference, NEC groups had higher IL-6, which was significantly reduced by EP treatment, suggesting EP's effectiveness in reducing NEC severity by lowering IL-6 in intestinal tissue.

EP has been found to be an effective agent in combating free oxygen radicals during instances of oxidative stress. Previous studies have reported that the

administration of EP reduces the levels of MDA, a free oxygen radical, in conditions such as hemorrhagic shock and ischemia-reperfusion injury.²⁹ In the case of NEC, oxidative stress leads to the formation of free oxygen radicals, which in turn causes an increase in tissue MDA levels.²³ A study was conducted to investigate the effects of quercetin, a flavonoid derivative known for its antioxidant properties, on newborn rats with NEC. After creating the NEC model, quercetin was administered to one group of rats while another group did not receive any treatment. The results revealed a decrease in oxidative stress parameters, specifically MDA levels, in the tissues of the treated group compared to the non-administered group.³⁰ A study conducted by Guven et al. demonstrated that the rats with the NEC model who were administered melatonin exhibited a decrease in the oxidative stress parameter, MDA, compared to the group that did not receive melatonin.³¹ In our study, we did not find any significant difference in MDA levels in intestinal tissues between the two control groups. However, we did observe a significantly higher level of MDA in the NEC groups compared to the control groups. In the NEC group that received EP, it was observed that the levels of tissue MDA decreased significantly compared to the NEC group that did not receive EP.

This study is the first to examine the impact of EP on neonates with an NEC model. The study demonstrated that when EP was administered intraperitoneally to newborn rats with an NEC model, it did not have any impact on the clinical status or body weight of the rats. However, it did lead to an improvement in the macroscopic appearance of the bowel. Although the application of EP did not have a statistically significant effect on the histopathological improvement in the NEC group, it was observed that it improved the overall histopathological condition. It has been found that administering EP intraperitoneally to rats with an NEC model resulted in a reduction in the severity of NEC. This was achieved by decreasing the level of IL-6 in the intestinal tissue and reducing oxidative stress by lowering the MDA level.

This is a pioneering study to investigate the effects of EP on the NF- κ B activation pathway and the TLR-4 inhibition pathway in the pathogenesis of NEC. Due to financial difficulties, this research could not be carried out. It may be necessary to enlarge the sample sizes to obtain more statistically significant results. These situations are the limitations of the study.

Conclusions

It is predicted that the application of EP in diseases with high mortality, such as NEC, can decrease the necessity for surgery. This finding will serve as the foundation for future phase studies exploring the use of EP as a new-generation anti-inflammatory agent.

Funding

This study was financed by the University of Health Sciences Scientific Research Projects Unit at the stage of procurement of materials, procurement of experimental animals, and procurement of laboratory services.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of human subjects and animals. The authors declare that the procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation and with the World Medical Association and the Declaration of Helsinki. The procedures were authorized by the Institutional Ethics Committee.

Confidentiality, informed consent, and ethical approval. This study did not involve human participants. All experimental animal procedures were approved by the Acibadem University Experimental Animals Local Ethics Committee (Approval date: January 08, 2020; Approval number: 2020/01) and were conducted in accordance with institutional and applicable ethical guidelines. The ethics committee approval document is attached for editorial records."

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing or creation of the content of this manuscript.

References

1. Thompson AM, Bizzarro MJ. Necrotizing enterocolitis in newborns: pathogenesis, prevention and management. *Drugs*. 2008;68:1227-38.
2. Terrin G, Scipione A, De Curtis M. Update in pathogenesis and prospective in treatment of necrotizing enterocolitis. *Biomed Res Int*. 2014;2014:543765.
3. Chen B, Na F, Yang H, Li R, Li M, Sun X, et al. Ethyl pyruvate alleviates radiation-induced lung injury in mice. *Biomed Pharmacother*. 2017;92:468-78.

4. Kim YM, Park EJ, Kim JH, Park SW, Kim HJ, Chang KC. Ethyl pyruvate inhibits the acetylation and release of HMGB1 via effects on SIRT1/STAT signaling in LPS-activated RAW264.7 cells and peritoneal macrophages. *Int Immunopharmacol*. 2016;41:98-105.
5. Scrima R, Menga M, Pacelli C, Agriesti F, Cela O, Piccoli C, et al. Para-hydroxyphenylpyruvate inhibits the pro-inflammatory stimulation of macrophage preventing LPS-mediated nitro-oxidative unbalance and immunometabolic shift. *PLoS One*. 2017;12:e0188683.
6. Dong N, Xu X, Xue C, Wang C, Li X, Shan A, et al. Ethyl pyruvate protects against Salmonella intestinal infection in mice through down-regulation of pro-inflammatory factors and inhibition of TLR4/MAPK pathway. *Int Immunopharmacol*. 2019;71:155-63.
7. Guo X, Guo R, Luo X, Zhou L. Ethyl pyruvate ameliorates experimental colitis in mice by inhibiting the HMGB1-Th17 and Th1/Tc1 responses. *Int Immunopharmacol*. 2015;29:454-61.
8. Venkataraman R, Kellum JA, Song M, Fink MP. Resuscitation with Ringer's ethyl pyruvate solution prolongs survival and modulates plasma cytokine and nitrite/nitrate concentrations in a rat model of lipopolysaccharide-induced shock. *Shock*. 2002;18:507-12.
9. Fink MP. Ethyl pyruvate. *Curr Opin Anaesthesiol*. 2008;21:160-7.
10. Zhou W, Zheng XH, Rong X, Huang LG. Establishment and evaluation of three necrotizing enterocolitis models in premature rats. *Mol Med Rep*. 2011;4:1333-8.
11. Korkut S, Özdemir A, Yay AH, Yalçın B, Ceylan M, Korkmaz L, et al. Obestatin reduces intestinal damage in experimental necrotizing enterocolitis in newborn rats. *Am J Perinatol*. 2019;36:1179-87.
12. Zani A, Cordischi L, Cananzi M, De Coppi P, Smith VV, Eaton S, et al. Assessment of a neonatal rat model of necrotizing enterocolitis. *Eur J Pediatr Surg*. 2008;18:423-6.
13. Dvorak B, Halpern MD, Holubec H, Williams CS, McWilliam DL, Dominguez JA, et al. Epidermal growth factor reduces the development of necrotizing enterocolitis in a neonatal rat model. *Am J Physiol Gastrointest Liver Physiol*. 2002;282:G156-64.
14. Harris MC, Costarino AT, Sullivan JS, Dulkerian S, McCawley L, Corcoran L, et al. Cytokine elevations in critically ill infants with sepsis and necrotizing enterocolitis. *J Pediatr*. 1994;124:105-11.
15. Bloom J, Patel N, Wagmister J, Choudhury M, Eshghi M, Konno S. Prophylactic effect of ethyl pyruvate on renal ischemia/reperfusion injury mediated through oxidative stress. *Int Urol Nephrol*. 2019;51:85-92.
16. Barlow B, Santulli TV. Importance of multiple episodes of hypoxia or cold stress on the development of enterocolitis in an animal model. *Surgery*. 1975;5:687-90.
17. Yurttutan S, Ozdemir R, Canpolat FE, Oncel MY, Unverdi HG, Uysal B, et al. Beneficial effects of etanercept on experimental necrotizing enterocolitis. *Pediatr Surg Int*. 2014;30:71-7.
18. Hirfanoglu IM, Helvacioğlu F, Haltas H, Koseoglu B, Tatli MM, Tayman C, et al. Mesenchymal stem cell therapy in necrotizing enterocolitis: a rat study. *Pediatr Res*. 2011;70:489-94.
19. Cetinkaya M, Cansev M, Cekmez F, Tayman C, Canpolat FE, Kafa IM, et al. CDP-choline reduces severity of intestinal injury in a neonatal rat model of necrotizing enterocolitis. *J Surg Res*. 2013;183:119-28.
20. Corsini I, Polvani S, Tarocchi M, Tempesti S, Marroncini G, Generoso M, et al. Peroxisome proliferator-activated receptor- γ agonist pioglitazone reduces the development of necrotizing enterocolitis in a neonatal pre-term rat model. *Pediatr Res*. 2017;81:364-8.
21. Travadi J, Patole S, Charles A, Dvorak B, Doherty D, Simmer K. Pentoxifylline reduces the incidence and severity of necrotizing enterocolitis in a neonatal rat model. *Pediatr Res*. 2006;60:185-9.
22. Shimizu H, Yamamoto K. NF-kappa B and C/EBP transcription factor families synergistically function in mouse serum amyloid A gene expression induced by inflammatory cytokines. *Gene*. 1994;149:305-10.
23. Bazacliu C, Neu J. Pathophysiology of necrotizing enterocolitis: an update. *Curr Pediatr Rev*. 2018;15:68-87.
24. Prado C, Michels M, Ávila P, Burger H, Milioli MV, Dal-Pizzolo F. The protective effects of fecal microbiota transplantation in an experimental model of necrotizing enterocolitis. *J Pediatr Surg*. 2019;54:1578-83.
25. Nakame K, Kaji T, Mukai M, Shinyama S, Matsufuji H. The protective and anti-inflammatory effects of glucagon-like peptide-2 in an experimental rat model of necrotizing enterocolitis. *Peptides*. 2016;75:1-7.
26. Liang S, Lai P, Li X, Xu J, Bao Y, Fang Y, et al. Ulinastatin reduces the severity of intestinal damage in the neonatal rat model of necrotizing enterocolitis. *Med Sci Monit*. 2019;25:9123-30.
27. Tatli Ö, Pasli S, Imamoğlu M, Cicek M, Yadigaroglu M, Sahin A, et al. Potential therapeutic effects of ethyl pyruvate and N-acetyl cysteine in an experimental rat model of corrosive esophageal. *Arab J Gastroenterol*. 2020;21:260-6.
28. Zmora O, Gutzeit O, Segal L, Boulous S, Millo Z, Ginsberg Y, et al. Prophylactic antenatal N-Acetyl cysteine administration combined with post-natal administration can decrease mortality and injury markers associated with necrotizing enterocolitis in a rat model. *PLoS One*. 2020;15:e0233612.
29. Demir S, Kazaz IO, Aliyazicioglu Y, Kerimoglu G, Teoman AS, Yaman SO, et al. Effect of ethyl pyruvate on oxidative state and endoplasmic reticulum stress in a rat model of testicular torsion. *Biotech Histochem*. 2020;95:317-22.
30. Yazlıclı S, Özcan CU, Hismiogullari AA, Sunay FB, Özcan T, Berksoy EA, et al. Protective effects of quercetin on necrotizing enterocolitis in a neonatal rat model. *Am J Perinatol*. 2018;35:434-40.
31. Guven A, Uysal B, Gundogdu G, Oztas E, Ozturk H, Korkmaz A. Melatonin ameliorates necrotizing enterocolitis in a neonatal rat model. *J Pediatr Surg*. 2011;46:2101-7.

Communication as the most important non-technical skill in the operating room

La comunicación como la habilidad no técnica más importante en el quirófano

Miguel Vassallo, Kimberly G. Ordaz-Cedeño*, Rhayniveth Sequera, Gabriel Rondón, and José Morao

Department of Surgery II, Hospital Universitario de Caracas, Universidad Central de Venezuela, Caracas, Venezuela

Abstract

Objective: To carry out a review of the literature about the complementarity between communication in the surgical area as a non-technical skill and its fundamental aspect in surgical practice. **Methods:** A bibliographic search was carried out by consulting the databases of the National Library of Medicine, PubMed, SciELO and Google Scholar. Articles published in the last 10 years, in English and Spanish, were considered as inclusion criteria, according to a series of keywords. **Results:** The initial search provided 2597 titles, which were screened and 22 articles were recovered for study analysis. **Conclusions:** The process of communicating is not so simple and linear; communication must be clarified by the participants in the interaction since they must overcome certain internal and external variables for it to develop properly. Communication in the operating room must be assertive and effective, where everyone feels important and is able to understand what is happening with the ability to present their own opinions as a key factor for patient safety during the surgical time.

Keywords: Non-technical skills. Communication. Surgical competence.

Resumen

Objetivo: Realizar una revisión de la literatura acerca de la complementariedad entre la comunicación en el área quirúrgica como una habilidad no técnica y su aspecto fundamental en la práctica quirúrgica. **Métodos:** Se realizó una búsqueda bibliográfica consultando las bases de datos de la National Library of Medicine, PubMed, SciELO y Google Académico. Se consideraron como criterios de inclusión artículos publicados en los últimos 10 años, en idiomas inglés y español, de acuerdo con una serie de palabras clave. **Resultados:** La búsqueda inicial proporcionó 2597 títulos, los cuales fueron sometidos a cribaje y se recuperaron 22 artículos para el análisis del estudio. **Conclusiones:** El proceso del comunicar no es tan sencillo y lineal; la comunicación debe ser clarificada por los partícipes de la interacción, ya que deben superar ciertas variables internas y externas para que esta logre desarrollarse adecuadamente. La comunicación en el quirófano debe ser asertiva y efectiva, que todos se sientan importantes y sean capaces de comprender lo que está ocurriendo con capacidad de presentar sus propias opiniones como factor clave para la seguridad del paciente durante el tiempo quirúrgico.

Palabras clave: Habilidades no técnicas. Comunicación. Competencia quirúrgica.

*Correspondence:

Kimberly G. Ordaz-Cedeño

E-mail: Kimberlyordaz18@gmail.com

2444-0507/© 2025 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 28-07-2024

Date of acceptance: 16-02-2025

DOI: 10.24875/CIRUE.M25000944

Cir Cir (Eng). 2026;94(3):264-268

Contents available at PubMed

www.cirugiyacirujanos.com

Introduction

Communication is defined as the exchange or feed-back of information, ideas, and feelings that promotes teamwork as a whole, ensuring that all members share the same vision during the effective completion of tasks.¹ Non-technical skills, in turn, are defined as cognitive, social, and personal aspects that complement experience and knowledge. There are five fundamental pillars of non-technical skills: situational awareness, decision-making, leadership, teamwork, and communication.²

Over the last decade, adverse events in health care, especially in the surgical setting, have been considered to arise from failures in behavioural aspects rather than from a lack of expertise. In this regard, it has been reported that up to 86% of adverse events in surgery are related to non-technical skills, with 43% attributed solely to communication failures.³

Through the systematic analysis of the study of human factors related to surgery, it is possible to observe that good-quality care is not limited solely to the technical interaction between the surgeon and the patient on the operating table. Thus, errors can also be explained by understanding the interaction between environmental conditions and personal skills, both technical and non-technical, which allow the surgeon to interact with and respond to this dynamic system.^{3,4}

The objective of this article is to conduct a systematic review of the available literature on the complementarity between communication as a non-technical skill and its fundamental role in surgical practice.

Methods

We conducted a bibliographic search by consulting the databases of the National Library of Medicine, PubMed, SciELO, and Google Scholar. Articles published in the last 10 years (2014-2024), in English and Spanish, were considered as inclusion criteria. The following keywords were used: “non-technical skills”, “communication in surgery”, “medical competencies”, “non-technical skills in surgery”, “communication in surgery”, and their combinations using the Boolean operators AND and OR. Articles whose full text was not available were excluded.

Results

The initial review in the abovementioned databases yielded 2597 results, of which 2403 were excluded

because they were duplicate studies, were unrelated to the selected topic, referred to specific diseases, or focused on another non-technical skill. During title screening, 152 of the 194 studies retrieved were excluded. The abstracts of 42 articles were reviewed, and finally 22 articles related to the study topic were selected.

Discussion

The training of surgeons has always focused on the development of knowledge, clinical experience, and technical skills. Other aspects, such as decision-making, effective communication, and leadership, are developed informally and tacitly rather than being addressed during training. Communication is considered the foundation of teamwork and one of the most important factors to consider in training and assessment.⁵

Communication and its relationship with surgical skill

The operating room is an environment filled with potential situations, many of which cannot be controlled. However, those that can be controlled, including the verbal and gestural language and tone of voice of team members, must be managed. Establishing clear and effective communication ensures that all members of a team share a common vision and, in this way, can complete their tasks appropriately.³

Studies supporting communication in surgery

It has been shown that a large proportion of adverse events occurring in surgery are based on communication problems that tend to recur systematically; therefore, if communication improves, teamwork will improve.⁵ Data collected by the Joint Commission on Accreditation of Healthcare Organizations suggest that poor communication contributed to 70% of the sentinel events reported in 2005.⁶

Lingard et al.⁷ identified 421 communication events and classified almost one third of them as failures. In the analysis by Greenberg et al.⁸ on surgical malpractice claims, recurrent patterns and contributors to miscommunication resulting in patient injury were identified. In a study by Sánchez et al.,⁹ communication failures among professionals during the operative

act were evidenced, with 179 errors occurring in 35 surgical procedures; the number of errors varied individually between 1 (12.8%), 2 (7.8%), 3 (7.8%), and > 3 (71.5%), with different types of causes underlying communication errors by the surgical team.

Currently, we are only beginning to understand the importance of communication and its contribution to the efficient and safe performance of the team during the surgical act.¹⁰

How to achieve communication and its impact on the surgeon's surgical competence

Holding work meetings before and after the surgical act allows team members to discuss intraoperative performance, expected progress, and opportunities for improvement, thereby achieving effective feedback. Similarly, any complication, unexpected finding, or unforeseen event (such as equipment failures) that may have occurred during surgery should be communicated and discussed. This communication routine becomes a tool that will allow the team to maintain a good relationship and prepare for the next surgery, and will help the team jointly develop its skills and prevent the same errors from recurring.⁹

Challenges and recommendations for surgical practice and communication in surgery

An important step in creating highly effective teams lies in teaching staff how to be part of that team. Without training, they will only be groups of people working in the same place, but not a true team. Training will shape this group of people into a single entity that will face problems and work together in patient care.¹¹

Team communication is a key factor for patient safety during the surgical period, since this is a space in which multiple factors intervene, decisions are made, and ideas are shared and discussed in the patient's best interest. However, the communication process is not that simple and linear; communication must be clarified and negotiated by the participants in the interaction, since they are influenced by internal and external variables. Of note, communication can be subdivided into 3 branches: the 1st is verbal communication, which includes words (whether oral or written); the 2nd is non-verbal communication, which comprises the kinesic complex (body language) and

proxemics (associated with body distance); and the 3rd is the paralinguistic line (referring to expression through sounds, but without words).¹²

Team members prioritise familiarity over formality and help one another to avoid mistakes. To encourage this approach in medical care, institutions can follow different actions, such as:¹¹

- Using structured communication tools that eliminate hierarchies.
- Providing feedback to surgeons.
- Training teamwork.
- Addressing conflictive behaviours.

Common obstacles in the pursuit of surgical practice and communication

As with all skills, there are certain minimum parameters that must be met and exceeded to achieve effective communication in the surgical setting; these are known as internal and external barriers.

External barriers consist of noise, low voice volume, distance, time, and even lack of visual expression. Internal barriers, in turn, are differences in language, organisational culture, mood, and affective states among those involved.¹³

Surgical teams usually face challenges imposed by the sophisticated instruments they use, the need to rapidly transmit critical information, the changing nature of the patient's condition, and the uncertainty inherent to all surgery.¹⁴ In addition, team members may know very little about one another or be unaware of the needs of a given patient or procedure. As a consequence, when surgical staff do not communicate efficiently, critical information may be lost, forgotten, or misinterpreted, thereby endangering patient safety.^{14,15}

Strategies to overcome challenges and achieve surgical practice and communication in surgery

Understanding communication as a combined, multidirectional team process, with different senders, messages, and receivers, in which there must be a closed loop, we start from what was intended to be said, then move to what was said, what was heard, and finally what was understood. For communication to be effective, it must be complete, accurate, unambiguous, and clearly understood.¹⁶ Some characteristics or aspects that the communicative process must

fulfil to be effective have been described, such as message clarity, timeliness, assertiveness, and active listening.¹⁷ To meet these goals, the ISBAR strategy has recently been described, consisting of five components: identification, situation, background, assessment, and recommendation; it has been shown to improve communication in work teams, reducing errors and improving the satisfaction of both patients and staff.¹⁸

Conclusions

Traditionally, physicians, especially surgeons, have been trained to think more as individuals than as members of a team. However, patient safety requires an approach in which individuals with different training work together, share information, and coordinate their efforts to provide optimal care.¹⁹ Highly effective teams do not allow hierarchical barriers to endanger safety. When hierarchical differences are emphasised, people at the base of the pyramid often feel uncomfortable or directly afraid to voice problems or concerns. Rude, aggressive, or intimidating behaviours by individuals at the top of the hierarchy endanger communication, making that person appear unapproachable. The team leader is the one who sets the tone for all perioperative interactions.²⁰

Communication, both verbal and non-verbal, takes place in the operating room to ensure adequate performance and is affected by a series of internal and external factors. Factors that facilitate the development of good communication include clarity of expression and listening, respect, situational awareness, experience, and trust.²¹ In addition, communication is not only a process limited to the operating room, since situations must be created in which hierarchies are eliminated both inside and outside surgery and, consequently, individualistic action is rejected. There must be participation not only from the surgical team, but also spaces must be created in which the organisational and administrative areas are integrated to understand errors and areas for improvement, while also recognising the successful actions correctly performed by the team. Thus, communication in the operating room must be assertive and effective, always seeking teamwork, ensuring that everyone feels important and is able to understand what is happening and express opinions about it.²²

Management within the surgical area is a task that requires the collaboration of professionals from

different disciplines; hence the importance of interaction so that it is adequate and maintains a similar perception of the situation, ensuring optimal teamwork and guaranteeing patient safety. The institution may also allocate resources to train members of the surgical team in communication, something that is generally not provided during undergraduate training and whose absence may affect patient outcomes and increase the risk of professional liability claims.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Ethical considerations

Protection of human subjects and animals. The authors declare that no experiments were performed on human subjects or animals for this research.

Confidentiality, informed consent, and ethical approval. The study does not involve personal data, medical records, or human biological samples; therefore, ethical approval is not required. The SAGER guidelines do not apply.

Declaration on the use of artificial intelligence. The authors declare that no type of generative artificial intelligence was used for the writing or creation of content for this manuscript.

References

1. Vega Hurtado C. Importancia de las estrategias de comunicación entre médico y paciente. *Rev Med Inst Mex Seguro Soc.* 2020;58:197-201.
2. Navarro F, González S, Gabrielli M. Evaluación de las habilidades no técnicas en cirugía. *Rev Cir.* 2019;71:359-65.
3. Yule S, Flin R, Paterson-Brown S, Maran N. Non-technical skills for surgeons in the operating room: a review of the literature. *Surgery.* 2016;139:140-9.
4. Berner JE, Ewertz E. Importancia de las habilidades no técnicas en la práctica quirúrgica actual. *Cir Esp.* 2019;97:190-5.
5. Limardo A, Blanco L, Menéndez J, Ortega A. Competencias médicas: evaluación de habilidades no técnicas en cirugía en el proceso de formación continua profesional en un hospital público durante el periodo de un año. *Rev Arg Educ Med.* 2020;9:30-43.
6. The Joint Commission. Sentinel events statistic. 2006. Disponible en: <http://www.jointcommission.org/SentinelEvent/Statistics/>.
7. Lingard L, Espin S, Whyte S, Regehr G, Baker G, Reznick R, et al. Communication failures in the operating room: an observational classification of recurrent types and effects. *Qual Saf Health Care.* 2014;13:330-4.
8. Greenberg CC, Regenbogen SE, Studdert DN. The patterns of communication breakdowns resulting in injury to surgical patients. *J Vasc Surg.* 2017;46:395.

9. Sánchez C, Rodríguez S, Cruz D, Domínguez LC, Vega V. Caracterización de los patrones de comunicación en salas de cirugía, durante procedimientos en un hospital de tercer nivel. *Rev Colomb Cir.* 2013;28:31-8.
10. Allard MA, Blanié A, Brouquet A, Benhamou D. Learning non-technical skills in surgery. *J Visc Surg.* 2020;157(3 Suppl 2):S131-6.
11. Orgambidez-Ramos A, Pérez-Moreno PJ, Borrego-Alés Y. Estrés de rol y satisfacción laboral: examinando el papel mediador del engagement en el trabajo. *Revista de Psicología del Trabajo y de las Organizaciones.* 2015;31(2).
12. Frasier LL, Pavuluri Quamme SR, Ma Y, Wiegmann D, Levenson G, DuGoff EH, et al. Familiarity and communication in the operating room. *J Surg Res.* 2019;235:395-403.
13. Berner JE, Ewertz E. Importancia de las habilidades no técnicas en la práctica quirúrgica actual. *Cir Esp.* 2019;97:190-5.
14. Rosendal AA, Sloth SB, Rölfing JD, Bie M, Jensen RD. Technical, non-technical, or both? A scoping review of skills in simulation-based surgical training. *J Surg Educ.* 2023;80:731-49.
15. Petrone P. Principios de la comunicación efectiva en una organización de salud. *Rev Colomb Cir.* 2021;36:188-92.
16. Guerrero LMA, Enamorado DT. Comunicación en el quirófano. Revisión bibliográfica del estado actual, efectos y factores relacionados. *Tesela, Liderazgo y Gestión.* 2022;30:e13456.
17. Vega-Hurtado C. Importancia de las estrategias de comunicación entre médico y paciente. *Rev Med Inst Mex Seguro Soc.* 2020;58:197-201.
18. Stambulsky E. Comunicación efectiva en el quirófano: clave para la seguridad del equipo y el paciente. *Red de Simulación en Salud.* 2023. Disponible en: <https://redsimulacionensalud.com/simulacion-salud/comunicacion-efectiva-en-el-quirofano-clave-para-la-seguridad-del-equipo-y-el-paciente/>.
19. Leonard M, Graham S, Bonacum D. The human factor: the critical importance of effective teamwork and communication in providing safe care. *Qual Saf Health Care.* 2004;13:85-90.
20. Hu YY, Arriaga AF, Peyre SE, Corso KA, Roth EM, Greenberg CC. Deconstructing intraoperative communication failures. *J Surg Res.* 2012;177:37-42.
21. Arenas-Márquez H, Anaya-Prado R. Estrategias para mejorar la seguridad quirúrgica. *Cir Cir.* 2008;76:355-61.
22. Anaya PR, Medina PJB, Pérez NJV. Comunicación en el equipo quirúrgico y seguridad. *Cir Gen.* 2011;33(Supl 1):96-8.

Incidence of surgical site infection in head and neck surgery: a prospective cohort study

Incidencia de infección del sitio quirúrgico en cirugía de cabeza y cuello: estudio de cohorte prospectivo

Diego Rodríguez-Villar^{1,2}, Enrique Gea-Izquierdo^{3,4}, Pilar Cárdenas-Soriano^{4,5},
M.^a Concepción Villar-del-Campo⁶, Manuel Durán-Poveda⁴, and Gil Rodríguez-Caravaca^{4,5*}

¹Doctorate International School, Rey Juan Carlos University, Madrid, Spain; ²Department of Anaesthesia, Ramón y Cajal Hospital, Madrid, Spain; ³Faculty of Medicine, Pontifical Catholic University of Ecuador, Quito, Ecuador; ⁴Department of Medical Specialties and Public Health, Rey Juan Carlos University, Madrid, Spain; ⁵Preventive Medicine Unit, Alcorcón Foundation University Hospital, Madrid, Spain; ⁶Primary Care Department, Centro de Salud Los Cármenes, Madrid, Spain

Abstract

Objectives: Surgical site infection (SSI) causes high morbidity. Our aim was to study the incidence of SSI and to assess its risk factors in patients undergoing head and neck surgery. **Methods:** A prospective cohort study was conducted, from July 2007 to June 2023, by collecting data on patient, surgery, and infection-related variables, and calculating the incidence of SSI after a maximum follow-up period of 1 year. We assessed the effect of the different risk factors on SSI, using the relative risk adjusted with a logistic regression model. **Results:** We included 423 patients, mean age 65.8 years (standard deviation: 12.4). The most frequent comorbidities were neoplasms (82.1%) and diabetes mellitus (29.3%). The cumulative incidence of surgical infection across the follow-up period was 6.4% (95% confidence interval: 4.1-8.7). Most infections were superficial, with the most common microorganism being *Enterobacter cloacae* (42.8%). Duration of surgery above the 75th percentile was the only risk factor associated with SSI in the multivariate analysis ($p = 0.01$). **Conclusions:** The incidence of infection was low in neck surgery. Infection rates must be assessed continuously, and SSI risk factors identified early through the implementation of prospective hospital surgical-infection surveillance and control programs.

Keywords: Epidemiological surveillance. Head and neck surgery. Surgical site infection. Incidence. Cohort studies. Risk factors.

Resumen

Objetivo: Estudiar la infección del sitio quirúrgico (ISQ) y los factores de riesgo relacionados en pacientes intervenidos de cirugía de cuello. **Métodos:** Se realizó un estudio de cohorte prospectivo desde julio de 2007 hasta junio de 2023. Se recogieron datos de los pacientes, la cirugía y las infecciones quirúrgicas. Se calculó la incidencia de ISQ tras un seguimiento de 1 año. Se evaluó el efecto de los factores de riesgo con el riesgo relativo ajustado con un modelo de regresión logística. **Resultados:** Se incluyeron 423 pacientes, con una edad media de 65.8 años (DE: 1.4). La comorbilidad más frecuente incluyó neoplasias (82.1%) y diabetes mellitus (29.3%). La incidencia acumulada de ISQ fue del 6.4% (IC 95%: 4.1-8.7). La mayoría de las infecciones fueron superficiales y el microorganismo más frecuente fue *Enterobacter cloacae* (42.8%). En el análisis

*Correspondence:

Gil Rodríguez-Caravaca

E-mail: gil.rodriguez@salud.madrid.org

2444-0507/© 2025 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 26-08-2024

Date of acceptance: 12-01-2025

DOI: 10.24875/CIRUE.M25000945

Cir Cir (Eng). 2026;94(3):269-276

Contents available at PubMed

www.cirurgiaycirujanos.com

multivariante, la duración de la cirugía mayor que el percentil 75 se relacionó con la ISQ ($p = 0.01$). Conclusiones: La incidencia de ISQ en la cirugía de cuello fue baja. La incidencia de infección quirúrgica y los factores de riesgo asociados deben ser evaluados de forma continuada con programas de vigilancia y control de las infecciones quirúrgicas.

Palabras clave: Vigilancia epidemiológica. Cirugía de cabeza y cuello. Infección del sitio quirúrgico. Incidencia. Estudio de cohorte. Factores de riesgo.

Introduction

Healthcare-associated infections (HAIs) are those that occur during patients' healthcare processes and were neither present clinically nor incubating at the time of admission.¹ According to the latest Prevalence Study of Nosocomial Infection in Spain (*Estudio de Prevalencia de Infección Nosocomial en España/EPINE*), 6.0% of patients present with one or more HAIs.²

Surgical site infections (SSIs) are infections related to a surgical procedure, which occur in the surgical incision or nearby tissue and whose development is most likely within 15-30 days following surgery. SSIs are classified in three categories: superficial, if they only affect the cutaneous and subcutaneous tissue; deep, if they involve deep soft tissue (fascia and muscle walls); and organ-space, when they affect any distal zone to the sterile incision or cavity.³

SSIs are one of the main causes of morbidity and mortality in hospitalized patients,⁴ prolonging hospital stays, increasing readmissions, reducing the quality of life of the patient intervened, and raising the cost of healthcare processes.⁵⁻⁹ Within HAIs, SSIs lead (26.3%), ranking ahead of urinary infections (16.0%) and bacteremias (14.8%).^{2,10}

Within otorhinolaryngological surgery, head and neck surgery is one of the most important types, mainly encompassing surgical interventions of neoplasms that develop in topographical regions of the mouth, throat, glottis, salivary glands, and skin of the face and neck. These tumors account for around 3% of all cancers, and their incidence is on the rise.¹¹ Treatment of this type of neoplasm is multidisciplinary, with resection of the tumor being practically indispensable for its cure.

The development of SSIs in head and neck surgery is a relatively frequent complication. According to the literature, these figures fluctuate widely, depending on the degree of contamination of surgery, ranging anywhere from 3% to 85%^{7,8,12,13} according to whether it is classed as "clean," "clean-contaminated," "contaminated" or "dirty."

We sought to study the incidence of SSI and assess the risk factors that determine it in patients undergoing head and neck surgery at a University Teaching Hospital.

Methods

Study design

We conducted a prospective cohort study from July 2007 to June 2023, covering patients over 18 years of age undergoing head and neck surgery at a university teaching hospital and excluding those who at the time of surgery presented with suspicion of infection or were receiving antibiotic treatment for confirmed infection. All surgical procedures falling within the CDC's NECK category were included (Table 1). The study was approved by the Clinical Ethics Research Committee of the hospital (number 16/91), and informed consent was obtained from all subjects and/or their legal guardian(s).

Sample size and data collection

Sample size was estimated on the basis of a 95% confidence interval (CI), a precision of 5%, an estimated proportion of infection of 25%, and an envisaged 10% of losses. A necessary sample size of at least 320 patients was thus estimated. The patients were selected from surgical programming and were consecutively recruited throughout the study period.

The research data were collected on a purpose-designed form and entered onto a normalized relational database. The following were collected: sociodemographic variables pertaining to patients and their peri-operative clinical situation, including date of birth, sex, presence of comorbidities (malnutrition, kidney failure, diabetes mellitus, neoplasm, chronic obstructive pulmonary disease (COPD), cirrhosis, obesity and immunodeficiency), need for transfusion or drainage during the intervention; variables relating to hospital stay, that is, date of admission, diagnosis and date of discharge; variables related with presurgical preparation and

Table 1. Surgical procedures studied (ICD-9-CM) of head and neck surgery

ICD-9-CM*	Intervention	n	%
30.1	Hemilaryngectomy	3	0.7
30.21	Epiglottidectomy	0	0
30.22	Vocal cordectomy	168	40
30.29	Other partial laryngectomy	30	7.1
30.3	Total laryngectomy	63	15
30.4	Radical laryngectomy	24	5.7
31.45	Open biopsy of larynx or trachea	24	5.7
40.40	Radical neck dissection, not otherwise specified	12	2.9
40.41	Radical neck dissection, unilateral	87	20.7
40.42	Radical neck dissection, bilateral	6	2.2
Total		423	100

*ICD-9-CM: international classification of diseases, 9th revision, clinical modification.

antibiotic prophylaxis; and surgery-related variables, such as date of intervention, duration of surgery, type of surgical procedure, degree of contamination of surgery (clean, clean-contaminated, contaminated and dirty), American Society of Anesthesiologists (ASA) anesthetic risk and the CDC National Nosocomial Infection Surveillance (NNIS) index. When a patient was diagnosed with SSI, we collected the date of diagnosis, infection site, result of culture (positive or negative), and microorganisms isolated.

SSIs were diagnosed by reference to the CDC's SSI criteria, and taking into account the patient's signs and symptoms, his/her clinical progress, surgical wound assessment, plus results of microbiological cultures where requested. Depth of infection was classified into superficial, deep, and organ-space.³

Patients were clinically followed up by daily visit from the date of admission to discharge, and thereafter through visits to outpatient clinics and emergency services, or from primary care through the review of electronic medical records (Selene®) or clinical information visor HORUS®. The study follow-up and total time were set in accordance with CDC criteria for implant surgery and were 1 year.

Statistical analysis

We used frequency distributions to describe the qualitative variables, which were then compared using

the Pearson Chi-square test or Fisher's exact test, where the conditions of application were not met (expected values < 5). In the case of the quantitative variables, these were described using the mean and standard deviation (SD), or the median and interquartile range (IQR) in those cases where the normality criteria were not fulfilled. To compare the quantitative variables, we used the Student's t-test or the Mann-Whitney U test in those cases where the data were not normally distributed. When the quantitative variables were compared in more than two groups, we used the analysis of variance test if the variables followed a normal distribution, or the Kruskal-Wallis non-parametric test on the contrary.

The incidence of infection was estimated by calculating the cumulative incidence of surgical infection across the follow-up period. SSI incidence was calculated by stratifying the risk according to the NNIS index. We standardized the incidence of infection by the indirect method, taking into account the cumulative incidence of SSIs in our region, in our country, and those reported by the CDC. The different risk factors were evaluated by calculating the relative risk (RR) of infection, both individually and grouped, with explanatory logistic regression models being built with the "backstep" method to control for confounding and interaction among different factors. In the internal calibration of the model (goodness-of-fit), we used the Hosmer-Lemeshow test.

The sample size was calculated using the Epidat 4.2 epidemiological software program, and the analyses were performed using the Statistical Package for the Social Sciences v.27 statistical software program, with statistical significance set at $p < 0.05$.

Results

The study covered a total of 423 surgical interventions of the head and neck. The breakdown showed that the patients were 81.4% men and 18.6% women, with a mean age of 65.8 years (SD = 12.4), 68.1 years in men and 55.5 years in women ($p < 0.01$). Table 2 shows the data on the patients' perioperative clinical situation and comorbidities.

The interventions were distributed as shown in Table 1: vocal cordectomy was the most frequent type (40%), followed by unilateral radical dissection of the neck (20.7%). The median overall hospital stay was 3 days (IQR = 13): the median stay among patients who did not develop SSIs was 3 days (IQR = 12) and 26 days (IQR = 26) among those with surgical

Table 2. Preoperative clinical situation and comorbidities of patients intervened

Comorbidity	n	%
Drainage	201	47.9
Shaving	18	4.3
Transfusion	9	2.1
Kidney failure	18	4.3
Malnutrition	18	4.3
Diabetes mellitus	123	29.3
Neoplasm	345	82.1
COPD	78	18.6
Cirrhosis	15	3.6
Obesity	36	8.6
Immunodeficiency	3	0.7
Total	423	100

COPD: chronic obstructive pulmonary disease.

infection ($p < 0.001$). The mean time from admission until diagnosis of infection was 10.6 days (SD = 6.9). The earliest case was diagnosed 2 days after surgery, and the latest, 24 days after surgery.

A total of 86.7% of intervened patients received appropriate presurgical preparation (95% CI: 83.8-89.1), with the most common cause of inappropriateness being non-application of mouthwash (67.1%). Of the 228 patients who required antibiotic prophylaxis, 189 received it, and among these, it was appropriate in 162 (85.7%; 95% CI: 80.8-90.1). The principal cause of inappropriateness was the choice of antibiotic (65.0%).

For the calculation of the NNIS index, we used the variables of classification of ASA anesthetic risk, degree of contamination of surgery, and duration of surgery (75th percentile, 225 min). While most patients (47.1%) had an NNIS of 0, no patient with an NNIS of 3 underwent surgery. Table 3 shows the mean stay of patients according to the NNIS index.

There were a total of 27 infections across the follow-up period, amounting to an overall SSI incidence of 6.4% (95% CI: 4.1-8.7). Most of the infections (44.4%) occurred in patients with an NNIS index of 1. Of the 27 surgical infections, 18 were superficial (66.7%) and 9 deep (33.3%). There were no organ-space SSIs within the 16 years analyzed.

A microbiological culture was performed in all infections and tested positive in 15. The microorganism most frequently found was *Enterobacter cloacae*

Table 3. Mean stay of patients according to the NNIS index (n = 423)

NNIS index	n (%)	Days (SD)
0	199 (47.1)	3.4 (5.8)
1	160 (37.9)	11.1 (12.6)
2	64 (15.0)	19.9 (14.2)
3	0 (0)	-

NNIS: National Nosocomial Infections Surveillance System; SD: standard deviation.

(43.0%). In six cases, two microorganisms were isolated, with the microorganisms responsible for the infections being shown in figure 1.

Cumulative SSI incidence at our hospital was compared to cumulative incidences, regionally and nationwide, issued by the INCLIMECC work group (Clinical Continuous Quality Improvement Indicators) and stratified according to NNIS risk (Table 3). Furthermore, these figures were also compared to the cumulative incidence rates issued by the CDC. Indirect standardization was performed by calculating the standardized infection ratio (SIR). The SIR at the hospital was: 1.48 (95% CI: 0.51-2.45) with respect to the Region; 1.94 (95% CI: 0.67-3.21) with respect to the country as a whole; and 2.07 (95% CI: 0.72-3.42) with respect to the CDC.

SSI risk factors were studied univariate. The statistically significant variables ($p < 0.05$) were as follows: malnutrition; duration of surgery above the 75th percentile; and the presence of drainage. Table 4 shows the univariate analysis of the various SSI risk factors: it omits the risk factors which had very low prevalence and those that did not present with infection in some of the groups.

The multivariate analysis (Table 4) studied the independent SSI risk factors shown to be statistically significant by the univariate analysis. Duration of surgery above the 75th percentile was the sole factor to prove statistically significant ($p = 0.01$).

Discussion

Overall SSI incidence in our study was 6.4% (95% CI: 4.1-8.7), a figure lower than that of most of the papers reviewed.¹⁴⁻¹⁶ There were no significant differences after standardization with existing data, neither for the region nor the country, nor those reported by the CDC.¹²

Surgery of the head and neck is a complex surgery that encompasses many procedures and different




Figure 1. Etiology of surgical site infection in head and neck surgery (%), n = 15.

Table 4. Univariate and multivariate analysis for SSI risk factors

Univariate analysis					
Risk factors	SSI (n: 9) n (%)	No SSI (n: 131) n (%)	RR ¹	95% CI ²	p
Mean age (SD ³)	65.7 (9.3)	65.8 (12.7)	1	0.9-1.1	0.99
Diabetes mellitus	3 (33.3)	38 (29.0)	1.22	0.3-5.1	0.52
COPD ⁴	1 (11.1)	25 (19.1)	0.53	0.1-4.4	0.48
Malnutrition	2 (22.2)	4 (3.1)	9.1	1.4-58.3	0.048
Cirrhosis	1 (11.1)	4 (3.1)	4.0	0.4-39.8	0.29
NNIS ⁵ index					
0	2 (22.2)	64 (48.9)	Ref.		
1	4 (44.4)	49 (37.4)	2.61	0.5-14.9	0.28
2	3 (33.3)	18 (13.7)	5.33	0.8-34.4	0.08
ASA ⁶ > 2	4 (44.4)	57 (43.5)	1.04	0.3-4.0	0.61
Duration of surgery (> P75)	6 (66.7)	28 (21.4)	7.36	1.7-31.3	0.007
Inappropriate prophylaxis	2 (28.6) of 7	7 (12.5) of 56	2.8	0.4-17.3	0.26
Drainage	7 (77.8)	60 (45.8)	4.14	0.8-20.7	0.06
Multivariate analysis					
Risk factors	RR	95% IC	p		
Malnutrition	6.57	0.9-49.2	0.07		
Duration of surgery (> P75)	6.52	1.5-28.6	0.01		

¹Relative risk.

²Confidence interval.

³Standard deviation.

⁴Chronic obstructive pulmonary disease.

⁵National nosocomial infections surveillance system index.

⁶American Society of Anesthesiologists.

different complications might arise as a result of partial/total flap compromises or vascular failures, which could account in part for the wide variability in SSI incidence rates found in the literature.^{7,8,12,13}

In our study, patients with SSIs had hospital stays longer than patients who did not present with surgical infections, in line with the literature.^{17,18} The 23-day increase in hospital stay was higher than that reported by Al-Qurayshi et al.⁷ for surgeries of the head and neck, estimated at 8 days, with our case being more in line with that described by Badia et al.,⁵ who found wide variability in the six countries included in their systematic review, with hospitalization always being longer in patients with surgical infection.

The microorganisms isolated in infections of head and neck surgery vary widely, particularly the common bacterial colonization of the respiratory tract. In our study, there were a very small number of infections, with the pathogens most frequently encountered being *E. cloacae* and gram-negative bacteria, a finding that agrees with other studies of surgical infection at this site.¹⁹⁻²²

The risk of developing a surgical infection depends on various factors, some of them of intrinsic to the patient him/herself (age, sex, obesity, nutritional status, or comorbidities), and others extrinsic (presurgical preparation, antimicrobial prophylaxis, use of drainage or transfusion, duration of intervention, appropriate instrumental sterilization or surgical technique used).^{14,23} Updated antibiotic prophylaxis and surgical preparation protocols based on the existing literature have been implemented at our teaching hospital and are periodically reviewed by the Infections Committee. Based on these, prophylaxis and preparation were rated appropriate or inappropriate in each case, citing the reason for inappropriateness where this existed.

Antibiotic prophylaxis failed to prove statistically significant in terms of reducing SSIs, probably due to the negligible number of infections found. We found studies that did not reflect a reduction in incidence of infection, despite appropriate antibiotic prophylaxis, such as those by Mitchell et al.²⁴ and Sepehr et al.,²⁵ though the majority tended to corroborate the existence of a lower incidence of SSIs in head and neck surgery when such prophylaxis was performed correctly,²⁶⁻²⁸ including systematic reviews and meta-analyses in clean-contaminated surgeries.²⁹

When it came to the percentage of presurgical preparation of patients, this was also very high (86.7%), as in the case of prophylaxis, with the main cause of inappropriateness being the non-administration of

surgical techniques. It should be stressed here that there is great variability in the diagnosis of SSIs in this type of patient, for example, due to the possibility that

mouthwash, which prevents post-operative pneumonia but has no influence on the development or lack of development of surgical infection.

While ASA anesthetic risk,³⁰ age,³¹ obesity¹⁴ and diabetes³²⁻³⁴ are risk factors described in the literature for this type of surgery, in our study, they failed to achieve statistical significance. Other risk factors for the development of SSIs, such as the patient's sex, use of drainage, or comorbidities such as COPD and cirrhosis, or described in other types of interventions³⁵⁻³⁹ were also not statistically significant in our study.

The study's univariate analysis displayed two significant variables: duration of surgery above the 75th percentile; and presence of malnutrition in the patient. Duration of surgery is a risk factor reported in other studies for this same surgical site;^{8,31} in our study, its RR was 6.5 (95% CI: 1.5-28.6) following the multivariate analysis. The rise in risk among such patients may be due to a more prolonged exposure to the wound's contaminants, longer duration of oropharyngeal secretions in the field, or increased difficulty of the surgical act. In this respect, an important factor for the appearance of SSIs is the surgeon's experience, as highlighted by McHugh et al.⁴⁰ or Mallol et al.⁴¹ This variable was not evaluated in the study, since ours is a university teaching hospital where there is thus a higher turnover of surgeons and physicians in training during interventions.

In the multivariate logistic regression analysis, the RR associated with malnutrition was 6.6 (95% CI: 0.9-49.2). Malnutrition was assessed at our hospital using body mass index (BMI), with malnutrition being deemed to exist if BMI < 18. In this study, the percentage of patients who presented with malnutrition was lower than in others consulted;^{21,42} this may be due to the fact that malnutrition was measured with the BMI, whereas other authors considered malnutrition to exist if there was a loss of weight > 10% in the 3 months prior to surgery, or with measurement of serum albumin. Our study shows a higher risk of suffering SSIs if the patient is malnourished, a finding in line with other papers.^{21,30}

There are studies in which the association between malnutrition and SSIs has been seen, and hence Bassett and Dobie⁴³ report the relationship between malnutrition and worse prognosis in neck cancer. In this type of neoplasm, there may be a deficit of food intake due to gastrointestinal tract occlusion, plus the placement of tracheotomies for lengthy periods of time, which could entail severe malnourishment. Moreover,

there are tumor factors that promote endocrine alterations and systemic responses, such as the release of cytokines, which trigger the protein catabolism and stages of cachexia. Malnutrition also affects the wound-healing process, favoring the development of septic complications.

Accordingly, we feel that a prior nutritional assessment should be made of all patients undergoing head and neck surgery, due to its close relationship with states of malnutrition, and that other anthropometric and specific biochemical parameters should be assessed, and not only BMI. A number of studies have sought to counteract malnutrition by means of enteral formulae (with arginine, glutamine, omega-3 fatty acids, and nucleotides), with evidence of a reduction in SSI and hospital-stay figures following the use of such formulae.⁴⁴⁻⁴⁶

One of our study's possible limitations is that the incidence of infection may have been underestimated due to some mild cases of superficial SSIs with minimal symptoms having been missed, where the patient might not have made use of the emergency, outpatient ear, nose, and throat, or primary care services. We would expect the infections missed to be negligible, in view of the fact that patients were followed up after discharge and over the course of a year. Similarly, the identification of some risk factors for surgical infection, detected through the patient's clinical history, may not be wholly accurate: for instance, if there are different forms of measurement (such as malnutrition) that pose problems when it comes to making comparisons with other studies in the literature which use different measurements. Finally, though the sample size was calculated for the assessment of SSI incidence, some of the risk factors studied may not be represented because they are factors with a very low prevalence. We endeavored to control for losses to follow-up by estimating a percentage of possible losses in the pre-calculation of sample size, in order not to lose statistical power.

Conclusions

Incidence of SSIs in head and neck surgery at our hospital during the study period was low. According to the results of this study, in order to reduce the risk of infection among the patients intervened, special attention should be paid to optimizing surgical time and ensuring an improvement in the nutritional status of patients who undergo these types of interventions.

A low SSI incidence rate is a good measure for assessing surgical practice and, by extension, quality in healthcare; hence, it is essential for infection rates to be evaluated and the possible risk factors identified. In this regard, prospective hospital surgical infection surveillance and control programs should be implemented, thereby allowing for continuous assessment of SSI incidence and the adoption of early preventive measures targeting potentially modifiable risk factors.

Funding

This study was funded by the European Regional Development Fund (ERDF) and the Health Research Fund (Fondo de Investigación Sanitaria/FIS), supporting the research projects PI11/01272, PI14/01136, and PI19/00987, which enabled the completion of this study. E.G.I. received a María Zambrano Program fund from the European Union-Spain.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of humans and animals. The authors declare that the procedures followed complied with the ethical standards of the responsible human experimentation committee and adhered to the World Medical Association and the Declaration of Helsinki.

Confidentiality, informed consent, and ethical approval. The authors have followed their institution's confidentiality protocols, obtained informed consent from patients. The study was approved by the Clinical Ethics Research Committee of the Fundación Alcorcón University Teaching Hospital (number 16/91). All research activities conformed to the principles embodied in the Declaration of Helsinki.

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing of this manuscript.

References

- Horan TC, Andrus M, Dudeck MA. CDC/NHSN surveillance definition of health care-associated infection and criteria for specific types of infections in the acute care setting. *Am J Infect Control*. 2008;36:309-32.
- Parra LM, Cantero M, Orti-Lucas RM, Salcedo-Leal I, Asensio A, EPINE study Group. Evaluation of infection prevention and control programmes according to the European centre for disease prevention and control and the world health organization in Spain 2012-2022: indicators of core component 1. *J Hosp Infect*. 2024;147:17-24.
- Yin V, Perren Cobb J, Wightman SC, Atay SM, Harano T, Kim AW. Centers for disease control (CDC) wound classification is prognostic of 30-day readmission following surgery. *World J Surg*. 2023; 47:2392-400.
- Klevens RM, Edwards JR, Richards CL Jr., Horan TC, Gaynes RP, Pollock DA, et al. Estimating health care-associated infections and deaths in U.S. Hospitals, 2002. *Public Health Rep*. 2007;122:160-6.
- Badia JM, Casey AL, Petrosillo N, Hudson PM, Mitchell SA, Crosby C. Impact of surgical site infection on healthcare costs and patient outcomes: a systematic review in six European countries. *J Hosp Infect*. 2017; 96:1-15.
- Rizki Purba AK, Luz CF, Wulandari RR, Van Der Gun I, Dik JW, Friedrich AL, et al. The impacts of deep surgical site infections on readmissions, length of stay, and costs: a matched case-control study conducted in an academic hospital in the Netherlands. *Infect Drug Resist*. 2020;13:3365-74.
- Al-Qurayshi Z, Walsh J, Owen S, Kandil E. Surgical site infection in head and neck surgery: a national perspective. *Otolaryngol Head Neck Surg*. 2019;161:52-62.
- Karakida K, Aoki T, Ota Y, Yamazaki H, Otsuru M, Takahashi M, et al. Analysis of risk factors for surgical-site infections in 276 oral cancer surgeries with microvascular free-flap reconstructions at a single university hospital. *J Infect Chemother*. 2010;16:334-9.
- Alfonso JL, Blasco Pereperez S, Moreno Canoves J, Melgar Martinez M, Martinez Martinez I, Martin-Moreno JM. Are we really seeing the total costs of surgical site infections? A Spanish study. *Wound Repair Regen*. 2007;15:474-81.
- Allegranzi B, Nejad SB, Combescure C, Graafmans W, Attar H, Donaldson L, et al. Burden of endemic health-care-associated infection in developing countries: systematic review and meta-analysis. *Lancet*. 2011; 377:228-41.
- Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer statistics, 2022. *CA Cancer J Clin*. 2022;72:7-33.
- Edwards JR, Peterson KD, Mu Y, Banerjee S, Allen-Bridson K, G Morrell, et al. National healthcare safety network (NHSN) report: data summary for 2006 through 2008, issued December 2009. *Am J Infect Control*. 2009;37:783-805.
- Simo R, French G. The use of prophylactic antibiotics in head and neck oncological surgery. *Curr Opin Otolaryngol Head Neck Surg*. 2006; 14:55-61.
- Khariwala SS, Le B, Pierce BH, Vogel RI, Chipman JC. Antibiotic use after free tissue reconstruction of head and neck defects: short course vs. Long course. *Surg Infect (Larchmt)*. 2016;17:100-5.
- Bourget A, Chang JT, Wu DB, Chang CJ, Wei FC. Free flap reconstruction in the head and neck region following radiotherapy: a cohort study identifying negative outcome predictors. *Plast Reconstr Surg*. 2011; 127:1901-8.
- Durand ML, Yarlagadda BB, Rich DL, Lin DT, Emerick KS, Rocco JW, et al. The time course and microbiology of surgical site infections after head and neck free flap surgery. *Laryngoscope*. 2015;125:1084-9.
- De Lissovoy G, Fraeman K, Hutchins V, Murphy D, Song D, Vaughn BB. Surgical site infection: incidence and impact on hospital utilization and treatment costs. *Am J Infect Control*. 2009;37:387-97.
- Lebo NL, Quimby AE, Caulley L, Thavorn K, Kekre N, Brode S, et al. Surgical site infection affects length of stay after complex head and neck procedures. *Laryngoscope*. 2020;130:E837-42.
- Park SY, Kim MS, Eom JS, Lee JS, Rho YS. Risk factors and etiology of surgical site infection after radical neck dissection in patients with head and neck cancer. *Korean J Intern Med*. 2016;31:162-9.
- Hirakawa H, Hasegawa Y, Hanai N, Ozawa T, Hyodo I, Suzuki M. Surgical site infection in clean-contaminated head and neck cancer surgery: risk factors and prognosis. *Eur Arch Otorhinolaryngol*. 2012; 270:1115-23.
- Cannon RB, Houlton JJ, Mendez E, Futran ND. Methods to reduce postoperative surgical site infections after head and neck oncology surgery. *Lancet Oncol*. 2017;18:e405-13.
- Skitarelic N, Morovic M, Malestar D. Antibiotic prophylaxis in clean-contaminated head and neck oncological surgery. *J Craniomaxillofac Surg*. 2007;35:15-20.
- Xu Z, Qu H, Kanani G, Guo Z, Ren Y, Chen X. Update on risk factors of surgical site infection in colorectal cancer: a systematic review and meta-analysis. *Int J Colorectal Dis*. 2020;35:2147-56.
- Mitchell RM, Mendez E, Schmitt NC, Bhargava AD, Futran ND. Antibiotic prophylaxis in patients undergoing head and neck free flap reconstruction. *JAMA Otolaryngol Head Neck Surg*. 2015;141:1096-103.
- Sepehr A, Gutierrez Santos BJ, Chou C, Karimi K, Devic Z, Oels S, et al. Antibiotics in head and neck surgery in the setting of malnutrition, tracheotomy, and diabetes. *Laryngoscope*. 2009;119:549-53.
- Alonso-García M, Toledano-Muñoz A, Aparicio-Fernández JM, De-la-Rosa-Astacio FM, Rodríguez-Villar D, Gil-De-Miguel A, et al. Adequacy of antibiotic prophylaxis and incidence of surgical site infections in neck surgery. *Sci Rep*. 2021;11:16413.

27. Velanovich V. A meta-analysis of prophylactic antibiotics in head and neck surgery. *Plast Reconstr Surg*. 1991;87:429-34; discussion 435.
28. Bartella AK, Kamal M, Teichmann J, Kloss-Brandstätter A, Steiner T, Hölzle F, et al. Prospective comparison of perioperative antibiotic management protocols in oncological head and neck surgery. *J Craniomaxillofac Surg*. 2017;45:1078-82.
29. Vander Poorten V, Uytendaele S, Robbins ST, Rodrigo JP, De Bree R, Laenen A, et al. Perioperative antibiotics in clean-contaminated head and neck surgery: a systematic review and meta-analysis. *Adv Ther*. 2020;37:1360-80.
30. Kamizono K, Sakuraba M, Nagamatsu S, Miyatomo S, Hayashi R. Statistical analysis of surgical site infection after head and neck reconstructive surgery. *Ann Surg Oncol*. 2014;21:1700-5.
31. Cohen LE, Finnerty B, Golas AR, Ketner JJ, Weinstein A, Boyko T, et al. Perioperative antibiotics in the setting of oropharyngeal reconstruction: less is more. *Ann Plast Surg*. 2016;76:663-7.
32. Chiu TH, Tsao CK, Chang SN, Lin JW, Hwang JJ. Clinical consequences of head and neck free-flap reconstructions in the DM population. *Sci Rep*. 2021;11:6034.
33. Lo SL, Yen YH, Lee PJ, Liu CC, Pu CM. Factors influencing postoperative complications in reconstructive microsurgery for head and neck cancer. *J Oral Maxillofac Surg*. 2017;75:867-73.
34. Raikundalia MD, Fang CH, Spinazzi EF, Vazquez A, Park RC, Baredes S, et al. Impact of diabetes mellitus on head and neck cancer patients undergoing surgery. *Otolaryngol Head Neck Surg*. 2016;154:294-9.
35. Shao J, Zhang H, Yin B, Li J, Zhu Y, Zhang Y. Risk factors for surgical site infection following operative treatment of ankle fractures: a systematic review and meta-analysis. *Int J Surg*. 2018;56:124-32.
36. Colás-Ruiz E, Del-Moral-Luque JA, Gil-Yonte P, Fernández-Cebrián JM, Alonso-García M, Villar-Del-Campo MC, et al. Incidencia de infección de sitio quirúrgico y factores de riesgo en cirugía de recto. Estudio de cohortes prospectivo. *Cir Esp*. 2018;96:640-7.
37. Del-Moral-Luque JA, Sánchez-Santana T, Gil-Yonte P, Fernández-Cebrián JM, Hijas-Gómez AI, Rodríguez-Caravaca G. Effect of a plan for quality improvement and clinical safety in the incidence of surgical site infections in appendectomy. A quasi-experimental study. *Cir Cir*. 2018;86:437-45.
38. Mesolella M, Allosso S, M Di Lullo A, Ricciardiello F, Motta G. Postoperative infectious complications in head and neck cancer surgery. *Ann Ital Chir*. 2022;93:637-47.
39. Mellinshoff SC, Bruns C, Al-Monajjed R, Cornely FB, Grosheva M, Hampf JA, et al. Harmonized procedure coding system for surgical procedures and analysis of surgical site infections (SSI) of five European countries. *BMC Med Res Methodol*. 2022;22:225.
40. McHugh SM, Hill AD, Humphreys H. Intraoperative technique as a factor in the prevention of surgical site infection. *J Hosp Infect*. 2011;78:1-4.
41. Mallol M, Sabaté A, Kreisler E, Dalmau A, Camprubi I, Trenti L, et al. Incidencia de la infección de la herida quirúrgica en cirugía colorrectal electiva y su relación con factores perioperatorios. *Cir Esp*. 2012;90:376-81.
42. Martín-Villares C, San-Román-Carabajo J, Fernández-Pello ME, Tapia-Risueño M, Domínguez-Calvo J. El estado nutricional en pacientes con cáncer de cabeza y cuello: implicaciones pronósticas. *Nutr Hosp*. 2003;18:91-4.
43. Bassett MR, Dobie RA. Patterns of nutritional deficiency in head and neck cancer. *Otolaryngol Head Neck Surg*. 1991;91:119-25.
44. Waitzberg DL, Saito H, Plank LD, Jamieson GG, Jagannath P, Hwang TL, et al. Postsurgical infections are reduced with specialized nutrition support. *World J Surg*. 2006;30:1592-604.
45. Marik PE, Zaloga GP. Immunonutrition in high-risk surgical patients: a systematic review and analysis of the literature. *JPEN J Parenter Enterol Nutr*. 2010;34:378-86.
46. Durán Poveda M, Suárez-De-La-Rica A, Cancer Minchot E, Ocón Bretón J, Sánchez Pernaute A, Rodríguez Caravaca G, et al. The prevalence and impact of nutritional risk and malnutrition in gastrointestinal surgical oncology patients: a prospective, observational, multicenter, and exploratory study. *Nutrients*. 2023;15:3283.

Ethical considerations in dietary supplement consumption behaviors

Consideraciones éticas en los comportamientos de consumo de suplementos dietéticos

İlknur Arslan-Aras^{1*} and Aykut Ekiyor²

¹Department of Health Management, Faculty of Health Sciences, Yüksek İhtisas University; ²Department of Health Management, Faculty of Economics and Administrative Sciences, Ankara Hacı Bayram Veli University, Ankara, Turkey

Abstract

Objective: This study examines the dietary supplement consumption behaviors and ethical attitudes aiming to contribute to a more conscientious and ethical consumption culture within the health industry. **Methods:** A descriptive cross-sectional study was conducted from October to December 2019 with a sample size of 384 individuals aged 20-65 in Turkey. Analysis was performed using the Statistical Package for the Social Sciences 22.0, including exploratory factor analysis and statistical tests. **Results:** About 55.5% of participants used dietary supplements, with Vitamin D, Vitamin B, Omega-3, and multivitamin complexes being common. Ethical concerns included inadequate label information, price discrepancies, and legal oversight issues. Statistically significant differences were observed in gender regarding the dimensions of meeting needs and promotion, in age regarding the legal dimension, and in the reasons for visiting a dietitian in the other four dimensions, except for the legal dimension ($p < 0.05$). **Conclusions:** This study revealed that factors such as gender, age, and dietary preferences influence individuals' supplement choices. Men were found to be more inclined toward the meeting needs and promotion aspects of dietary supplement purchases than women. In addition, age group and dietary preferences were found to affect the sensitivity of individuals in their supplement choices.

Keywords: Dietary supplements. Consumer behavior. Health ethics. Supplement regulation.

Resumen

Objetivo: Examinar los comportamientos de consumo de suplementos dietéticos y las actitudes éticas con el objetivo de contribuir a una cultura de consumo más consciente y ética dentro de la industria de la salud. **Métodos:** Se realizó un estudio descriptivo transversal de octubre a diciembre de 2019 con un tamaño de muestra de 384 individuos de entre 20 y 65 años, en Turquía. El análisis se realizó utilizando SPSS 22.0, incluido el análisis factorial exploratorio y las pruebas estadísticas. **Resultados:** El 55.5% de los participantes utilizaron suplementos dietéticos, siendo comunes la vitamina D, la vitamina B, los ácidos grasos omega-3 y los complejos multivitamínicos. Las preocupaciones éticas incluían información inadecuada en las etiquetas, discrepancias en los precios y problemas de supervisión legal. Se observaron diferencias estadísticamente significativas en el sexo en relación con las dimensiones de satisfacción de las necesidades y promoción, en la edad en relación con la dimensión legal, y en los motivos para acudir al dietista en las otras cuatro dimensiones, excepto en la dimensión legal ($p < 0.05$). **Conclusiones:** Este estudio reveló que factores como el sexo, la edad y las preferencias dietéticas influyen en la elección de suplementos por parte de los individuos. Se observó que los hombres se inclinaban más que las mujeres por los aspectos de satisfacción de las necesidades y promoción de la compra de suplementos dietéticos. Además, se encontró que el grupo de edad y las preferencias dietéticas influían en la sensibilidad de los individuos en el momento de elegir suplementos.

Palabras clave: Suplementos dietéticos. Comportamiento del consumidor. Ética de la salud. Regulación de suplementos.

*Correspondence:

İlknur Arslan-Aras
E-mail: ilknurarslan@yiu.edu.tr

Date of reception: 09-06-2024

Date of acceptance: 12-11-2024

DOI: 10.24875/CIRUE.M24000946

Cir Cir (Eng). 2026;94(3):277-286

Contents available at PubMed

www.cirugiaycirujanos.com

2444-0507/© 2024 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Today, people in pursuit of healthy living and longevity are turning to various food supplements to support their diets. Food supplements are generally defined as products containing nutrients such as vitamins, minerals, amino acids, herbs, or herbal dietary ingredients. The use of these products has increased due to successful sales strategies, advertisements, and the widespread use of the internet.¹ However, the ethical use of food supplements is an important issue in terms of public health and safety. One area of particular focus for ethical questions about food supplements is their impact on intake behavior. Factors such as consumers' reasons for using food supplements, their level of trust in these products, and their reactions to health claims are important factors shaping consumer purchasing behavior. Therefore, ethically evaluating the purchasing behavior of food supplements is an important step that can affect both the health and well-being of individuals and the overall health of society. In this context, food safety and ethical principles should be active at every stage of agriculture and food systems.²

Based on this idea, the aim of this study was to determine the intake behaviors of individuals taking food supplements and to examine their ethical attitudes in this field. The results of this study aim to contribute to the creation of a more conscious and ethical consumption culture in the health industry by shedding light on the ethical evaluations of consumers regarding the use of food supplements. In addition, the results of this research will be a valuable source of information for health policy makers and marketing experts.

Conceptual framework

Food supplement use in other countries

Food supplements are carbohydrates, protein, fat, minerals, vitamins, herbs, and nectars. Many studies have shown that the characteristics of people who use food supplements are that they tend to be older, have a low body mass index, exercise, do not smoke, and have a high level of education.³⁻⁹ and are more preferred by those who strongly believe in the importance of treating certain diseases and athletes.¹⁰⁻¹²

In the last 30 years, there has been a great increase in the types and sales of food supplements. The Ministry of Food, Agriculture, and Livestock is the main

regulator in Turkey, and the total sales of food supplements in 2020 are estimated to be 885 million TL. The National Health and Nutrition Examination Survey began to investigate motivations for the use of food supplements for the first time in 2007. The purpose of this analysis was to examine the types of food used as well as the motivations for food supplement use by adults.¹³ A search of the Web of Science database with the keyword "Dietary Supplements" yielded 2,666 publications. The distribution of these publications by country is shown in the figure below.

Figure 1 shows that 1,320 of the 2,666 studies on dietary supplements were published in the USA. This is followed by the UK with 154 publications, Germany with 111 publications, Australia with 107 publications, and Italy with 98 publications.

As a result of the search of the Web of Science database on food supplements, the top 10 most cited studies were analyzed. Among these studies, the study "Standard methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements" by Prior et al. in 2005 ranks first with 2579 citations. This is followed by "Cellular antioxidant activity assay for evaluating antioxidants, foods and dietary supplements" by Wolfe and Liu in 2007, with 571 citations, and "Adverse cardiovascular and central nervous system events associated with dietary supplements containing ephedra alkaloids" by Haller and Benowitz in 2000, with 557 citations.

In Turkey, in a study conducted by Çoşkun and Turan in Istanbul on 1000 people representing four different sociocultural groups, 34.6% reported using dietary supplements. In 2017, in the study on the use of food supplements published by the Food Supplement and Nutrition Association (GTBD) and Ipsos Social Research Institute, the rate of use of food supplements was reported as 13% after a survey of 1750 people over the age of 18 with telephone support in 81 provinces. In 2019, in the study titled "Evaluation of Food Supplement Intake Behaviors and Related Factors of Individuals Between the Ages of 18-65 Applying to the Family Health Center" conducted by Karadağ, 32.3% of the participants reported that they took supplements.

Ethical dimension of food supplements use behaviors

Ethics are principles, values, and standards that define how individuals behave correctly toward other




Figure 1. Number of studies on dietary supplements by countries.

beings¹⁴ Professional ethics, which is a sub-branch of ethics, can be expressed as the sum of ethical principles and standards that guide behaviors in professional life and guide what to do and what not to do. Businesses feel the need to behave ethically and make decisions in order to increase their reputation and thus gain profit in the medium and long term, and it is beneficial to see the expenditures made for this purpose as one of the most valuable asset items of the business.¹⁵

Related to this issue, Torok and Murray in their 2008 study “Using the sword of professional ethics against misleading dietary supplements,” tried to understand how dietary supplements are currently regulated for scientists who want to take a more prominent role in protecting the integrity of the interests of science and society. In general, the Food and Drug Administration (FDA) oversees safety, manufacturing, labeling, and product literature for drugs and medical devices, while the Federal Trade Commission (FTC) regulates how products are marketed. Mannatech Inc. is a publicly traded multinational multi-level marketing company founded in 1993 by Samuel Caster that markets dietary supplements and personal care products.¹⁶ Due to consumer complaints filed with the FDA and unethical use of credit advertising, Mannatech’s sales declined by 50% after 2006. Scientists should provide informational materials to consumers to prevent misuse and unethical production. The FTC and agencies such as the FDA and the Office of Dietary

Supplements should have supplement manufacturers and contact information on their websites, and consumers should be able to check whether a supplement has been tested for safety and efficacy through the government.

Proactive steps are needed to prevent public mistrust of food supplements. One such measure could be to develop educational materials and training activities that can be used by medical schools and professional associations, such as the American Medical Association, to ensure that doctors and medical students are effective in countering suspicious claims. Doing so would not only address the lack of reliable information on dietary supplements available to the public, but would also give the science community a level of respectability to other professionals. In establishing standards for the proper labeling and manufacture of food supplements, the FDA has been confronted with the lack of information on the efficacy of some supplements and the lack of data to determine adequate dosage for children. Children are not just small adults in terms of dosage; untested mold supplements pose the potential for harm when administered to children. Moreover, it is clear that even benign substances such as vitamins taken in excessive amounts can be harmful.¹⁷

Methods

In this section of the study, the purpose and importance of the study, the data collection tool, and the analysis of the data will be discussed.

Purpose and hypothesis of the study

The aim of this study is to determine the intake behaviors of individuals taking food supplements to examine their ethical attitudes in this field. For this purpose, the following hypotheses were tested to determine the ethical dimension of dietary supplement use behaviors in the areas of needs satisfaction, ethics, monitoring, promotion, and legal areas.

- Hypothesis 1. Gender differs according to (a) Meeting needs (b) Ethical dimension (c) Monitoring dimension (d) Promotion dimension (e) Legal dimension of individuals' food supplement use behaviors
- Hypothesis 2. Age differs according to (a) Meeting needs (b) Ethical dimension (c) Monitoring dimension (d) Promotion dimension (e) Legal dimension of individuals' food supplement use behaviors
- Hypothesis 3. Dietitian Visiting Status differs according to (a) Meeting needs (b) Ethical dimension (c) Monitoring dimension (d) Promotion dimension (e) Legal dimension of individuals' food supplement use behaviors.

Population and sample of the study

The descriptive and cross-sectional study was conducted in October 2019 and December 2019. The research population consisted of individuals between the ages of 20 and 65 residing in Turkey. At the end of 2017, the population between the ages of 20 and 65 in Turkey was 48,355,065 (TÜİK, 2017); the sample size that can represent the research population was determined as 384 people with a 95% confidence interval. Convenience sampling method was used among the sampling methods.

Data collection tool of the study

While collecting the research data, the “Knowledge, attitude and usage behavior about food supplements” scale taken from the thesis of Karadağ, 2019, was used. After interviewing with medical representatives and pharmacies within the scope of expert opinion, the scale was answered as “I think ethical values are observed in food supplement information, I think ethical values are observed in pricing policies, I think ethical values are observed when obtaining legal permissions of food supplements, I think ethical values

are observed when determining the amount of use, I think that there is not enough supervision in the production and distribution processes of food supplements, I am hesitant that the prices on the internet and pharmacy prices are not the same, and I do not find it right to advertise food supplements in print and visual media” were added and the question “Food supplements are supervised by the Ministry of Food, Agriculture and Livestock” was removed.

After the questions to be asked in the questionnaire were determined, it was determined whether the questions in the draft questionnaire were compatible with the application, the aims, and assumptions of the application. Then, the questionnaire was tested by 10 pharmacists and 3 medical representatives before it was applied to the individuals and the questions that caused misunderstandings were rearranged. The questionnaire form can be classified as follows; in the first 24 questions of the questionnaire, it is aimed to obtain information about the general health status of individuals, such as gender, body mass index, sleep levels, chronic disease status, and quality of life.

Questions 24-30 of the questionnaire aimed to find out which supplements individuals who use food supplements take, the reasons for taking supplements, where they obtain supplements, and how much they spend on supplements in a year. Finally, in the 31st question, a 3-point Likert-type scale was used to determine the behaviors and attitudes of individuals toward food supplements. The answering scale for each statement of the 31st question in the scale is a 3-point Likert; 1 = No Idea, 2 = No, 3 = Yes.

Information was collected from social media and social media platforms (LinkedIn, Facebook, etc.) through the Google form. The survey method was chosen due to its ability to collect a wide range of information, speed, and cost advantages, while the face-to-face and social media survey method was chosen due to its high response rates and the ability to eliminate misunderstandings. A total of 474 questionnaire forms were collected.

Data analysis

The data obtained at the end of the research were analyzed with the Statistical Package for the Social Sciences 22.0 package program. First, frequencies and percentages were given according to the individual and demographic characteristics of the participants. Exploratory factor analysis was applied to the scale. At this stage, the Independent Samples t-test

Table 1. Factor analysis table

Ethical	Meeting needs	Ethics	Monitoring	Promotion	Legal
I believe ethical values are observed when obtaining legal permissions for dietary supplements	0.839				
I believe ethical values are observed in pricing policies.	0.748				
I believe ethical values are considered when determining the usage amount.	0.684				
I believe ethical values are observed in dietary supplement information.	0.675				
Since I started using dietary supplements, I feel healthier.		0.814			
I am aware of which dietary supplements I need for my health.		0.788			
Dietary supplements can meet all nutritional needs.		0.543			
The information on the labels of dietary supplements is sufficient.		0.445			
I am concerned about the side effects of dietary supplements.			0.781		
Dietary supplements should be prescribed and monitored by a physician.			0.597		
Taking dietary supplements along with prescribed medications can cause drug interactions.			0.591		
Vitamin and mineral levels should be regularly monitored with laboratory tests.			0.799		
I believe there is not enough supervision in the production and distribution processes of dietary supplements.				0.818	
I do not approve of dietary supplement advertisements in print and visual media.				0.720	
The fact that internet prices and pharmacy prices are not the same makes me skeptical.				0.565	
Dietary supplements are regulated by the Ministry of Health.					0.751
Physicians should have more information about dietary supplements.					0.632
Dietary supplements should be covered by health insurance.					0.557

and One-way analysis of variance (ANOVA) test were applied since the data were normally distributed and the variances were homogeneous. After conducting the ANOVA analysis, *post hoc* tests (Tukey) were utilized to determine between which groups the differences originated.

Validity and reliability of the research scale

In the study, factor analysis with varimax rotation was applied to 18 statements to reveal the construct validity of the ethical scale of dietary supplement intake behaviors. As a result of the analysis, five dimensions were reached as meeting needs, ethical dimension, monitoring dimension, promotion dimension, and life dimension. The rate of explaining the total variance of the five factors obtained was 69.543%,

the Kaiser-Meyer-Olkin value was 0.810, and the Bartlett Sphericity test value was 1240.994 and was significant ($p < 0.05$). The distribution table of the factor analysis according to the dimensions is given in table 1.

There are many methods to measure the reliability of scales. Cronbach's Alpha coefficient, which is one of them, is a method frequently used in reliability calculations of scales based on total score, such as Likert-type scales.¹⁸ The Cronbach's Alpha value used for the research was found to be quite reliable, with 0.73.

Findings

A total of 474 people participated in this study conducted on individuals living in Turkey. Of the 474

participants, 49.6% were female, and 50.4% were male. 47.6% of the individuals participating in the study were between the ages of 21 and 30. 62% of the participants are married, and 60.3% of them live with their spouse and children. When the participants are analyzed in terms of education and occupational groups, 41.6% are university graduates, 34% are public sector employees, and 32.9% are private sector employees. 53% of the participants are Social Security Institution (SSI/SSK) members, and 37.1% of them have an income level between 2000 and 4000, 27.4% between 4000 and 6000.

When the healthy living behaviors of the individuals participating in the study were evaluated, 47.26% were normal weight, and 37.97% were slightly overweight. While 38.80% of the participants do not exercise, 38.60% do mild exercise, and 46.60% do not go to general health check-ups. The proportion of the participants who consulted a nutrition and diet specialist was 26.2%, and 56.6% of them went to lose weight. When the nutritional status of the participants is analyzed; 51.70% of them apply equal weight from carbohydrates. 36.50% of the participants consume 1-15 L of water/day, 26.60% sleep between 7 and 8 h. In addition, 67.9% do not smoke, and 84% do not consume alcohol. Only 26.2% of the participants reported having a chronic disease. The most common diseases were hypertension and migraine, with 18%, followed by asthma with 14%. Among the participants with chronic diseases, 41.9% used medication when necessary, while 39.5% reported that they used medication every day. 55.5% of the participants reported taking food supplements. Among these, 57.8% use them every day. 41.8% spend 200 TL or less on food supplements in a year, while 38.8% spend between 200 and 500 TL. 70.7% of the participants obtain food supplements from pharmacies.

Table 2 shows that 15.8% of the participants take vitamin D, 12.8% vitamin B, 11.3% Omega 3, and 9.30% multivitamin complex.

When the reasons for using food supplements are analyzed in table 3, 17.60% of the participants use food supplements to "Increase body performance and energy during the day," 16.50% to "Promote general health," 16.20% to "Treat specific health problems," and 14.70% to "Increase immune power." Weight gain (0.40%) and weight loss (0.10%) are the least preferred reasons for use.

In table 4, when the participants' food supplement purchasing behaviors were examined in terms of ethics, 78.7% stated that physicians should have

Table 2. Distribution of food supplements used by the participants

Supplements	n	%	Supplements	n	%
Vitamin D	126	15.80	Coenzyme Q10	23	2.90
Vitamin B	102	12.80	Sports drinks and foods	18	2.30
Omega-3	90	11.30	Calcium	15	1.90
Multivitamin/multimineral complex	74	9.30	Selenium	14	1.80
Vitamin C	53	6.60	Alpha lipoic acid	9	1.10
Iron	53	6.60	Melatonin	9	1.10
Herbal supplements	52	6.50	Glucosamine	8	1.00
Probiotic	41	5.10	Protein Bar	5	0.60
Magnesium	39	4.90	Other*	4	0.50
Vitamin K	33	4.10	Amino Acid	3	0.40
Zinc	26	3.30	Protein Powder	2	0.30

*Indicates statistically significant difference (p < 0.05).

Table 3. Distribution of reasons for use of food supplements by participants

Reasons for use	n	%
To enhance daytime physical performance and energy	121	17.60
To promote general health	113	16.50
To treat specific health issues	111	16.20
To boost immunity	101	14.70
As recommended by a doctor	81	11.80
To improve nutrition	62	9.00
As recommended by a nutrition and diet specialist	25	3.60
To reduce stress	20	2.90
As recommended by a pharmacist	14	2.00
As recommended by a life/sports coach	14	2.00
Based on family, friends, and community recommendations	13	1.90
After seeing on social media (TV/radio/internet/phone/ads)	7	1.00
To gain weight	3	0.40
To lose weight	1	0.10

more information about food supplements, 76.8% stated that vitamin and mineral levels should be regularly monitored with laboratory tests, 72.6%

Table 4. Results of the ethical analysis of participants' food supplement intake behavior scale

Variables	n	%	n	%	n	%
Food supplements should be prescribed and monitored by a doctor.	176	66.9	62	23.6	25	9.5
Doctors should have more knowledge about food supplements.	207	78.7	34	12.9	22	8.4
Food supplements should be covered by health insurance.	191	72.6	46	17.5	26	9.9
Vitamin and mineral levels should be regularly monitored with laboratory tests.	202	76.8	39	14.8	22	8.4
Food supplements are regulated by the Ministry of Health.	128	48.7	76	28.9	59	22.4
I am concerned about the side effects of food supplements.	151	57.4	77	29.3	35	13.3
The information on the labels of food supplements is sufficient.	75	28.5	149	56.7	39	14.8
Food supplements may cause drug interactions when taken with prescribed medications.	141	53.6	42	16	80	30.4
I am aware of which food supplements I need for my health.	172	65.4	51	19.4	40	15.2
I feel healthier since I started using food supplements.	171	65	54	20.5	38	14.4
All nutritional needs can be met with food supplements.	57	21.7	166	63.1	40	15.2
I believe ethical values are considered in the information provided about food supplements.	81	30.8	115	43.7	67	25.5
I believe ethical values are considered in pricing policies.	58	22.1	136	51.7	69	26.2
I believe ethical values are considered when obtaining legal permits for food supplements.	77	29.3	104	39.5	82	31.2
I believe ethical values are considered when determining the dosage.	96	36.5	91	34.6	76	28.9
I believe there is not enough control in the production and distribution processes of food supplements.	127	48.5	74	28.2	61	23.3
The discrepancy between internet and pharmacy prices makes me hesitant.	189	71.9	50	19	24	9.1
I do not think it is correct to advertise food supplements in print and visual media.	142	54	89	33.8	32	12.2

stated that food supplements should be covered by health insurance, and 71.9% stated that they were hesitant about the fact that internet prices and pharmacy prices were not the same. 63.1% of the participants think that food supplements do not meet all nutritional needs, 56.7% think that the information written on the labels of food supplements is insufficient, and 51.7% think that ethical values are not observed in pricing policies.

According to this table, statistically significant differences were not found between the sub-dimensions of the scale. According to table 5, Hypotheses 1a, 1d, 2e, 3a, 3b, 3c, and 3d are accepted while the other hypotheses are rejected.

Discussion

The purpose of this study is to determine the purchasing behaviors of individuals who take dietary supplements to examine their ethical attitudes in this field. The finding that men pay more attention to the meeting

needs and promotion processes when buying dietary supplements than women is supported by various literature studies. This difference reflects differences in men's and women's behaviors and motivations for supplement use. First, women generally use supplements at higher rates than men. According to a study conducted in the USA, 63.8% of women used any food supplement in the past 30 days, compared to 50.8% of men.¹⁹ Although women have a higher propensity to use food supplements, men generally use supplements more strategically for specific health goals (e.g., increasing muscle mass or preventing a specific disease). Men generally appear to be more motivated to use supplements to achieve specific health goals. In this context, men are more careful in their choice of supplements to meet their needs and to ensure continuity of use. For example, men place more emphasis on muscle health and performance-enhancing supplements, whereas women generally prefer supplements that focus on general health and bone health.²⁰ Furthermore, among the factors that increase men's motivation

Table 5. Hypothesis test results

Category	Meeting needs (a)	Ethical dimension (b)	Monitoring dimension (c)	Promotion dimension (d)	Legal dimension (e)
Women Avg	2.30	1.96	2.50	2.37	2.53
Women SD	0.40	0.58	0.44	0.59	0.45
Men Avg	2.31	2.10	2.46	2.51	2.53
Men SD	0.52	0.58	0.47	0.51	0.51
T	-0.283	-1.938	0.695	-2.105	0.142
p	0.000*	0.623	0.089	0.047*	0.267
Hypothesis accepted	1a Accepted	1d Accepted			
20-30 Avg (1)	2.25	1.94	2.50	2.38	2.59
20-30 SD	0.49	0.57	0.44	0.60	0.47
31-40 Avg (2)	2.33	2.00	2.47	2.38	2.60
31-40 SD	0.43	0.58	0.43	0.53	0.38
41-50 Avg (3)	2.39	2.17	2.48	2.54	2.42
41-50 SD	0.44	0.70	0.58	0.47	0.51
51-65 Avg (4)	2.30	2.09	2.46	2.55	2.36
51-65 SD	0.43	0.46	0.41	0.54	0.52
F	1.092	1.875	0.065	1.727	3.866
p	0.35	0.13	0.98	0.16	0.010*
<i>Post hoc</i> test					1-4 p = 0.03 2-4 p = 0.04
Hypothesis accepted	2e Accepted				
Weight loss Avg (1)	2.38	2.05	2.50	2.54	2.41
Weight loss SD	0.37	0.55	0.37	0.48	0.53
Weight gain Avg (2)	1.86	1.48	1.93	1.94	2.33
Weight gain SD	0.55	0.39	0.57	0.70	0.47
Healthy eating Avg (3)	2.28	2.19	2.42	2.31	2.56
Healthy eating SD	0.45	0.50	0.49	0.48	0.38
Doctor's recommendation Avg (4)	2.44	1.84	2.28	2.00	2.71
Doctor's recommendation SD	0.42	0.63	0.49	0.69	0.28
F	4.83	5.25	5.16	5.8	1.64
p	0.000*	0.000*	0.000*	0.000*	0.19
<i>Post hoc</i> test	1-2 p = 0.002 2-3 p = 0.029 2-4 p = 0.021	1-2 p = 0.008 2-3 p = 0.001	1-2 p = 0.001 2-3 p = 0.013	1-2 p = 0.004 1-4 p = 0.037	
Hypothesis accepted	3a Accepted	3b Accepted	3c Accepted	3d Accepted	

*Indicates statistically significant difference ($p < 0.05$)

Avg: average; SD: standard deviation.

and commitment to supplement use, scientific evidence and the desire to be informed about the efficacy of products play an important role. Men often conduct extensive research on the health effects and scientific basis of a particular supplement before choosing it.²¹ This attitude increases their tendency to use supplements regularly and consistently.

Another result of this study shows that individuals aged 31-40 years are more concerned about the legal regulations of dietary supplements than other age groups. This age group may focus on legal concerns. Individuals in this age group are generally more conscious about protecting their health and improving their quality of life. This awareness is coupled with a desire to learn more about the safety and efficacy of the food supplements consumed.²² People in the 31-40 age group are aware that legal regulations are important to ensure the quality and safety of supplements.

The study found that individuals who go on a diet to gain weight are more sensitive to the meeting of needs, ethical standards, monitoring processes, and promotion methods when choosing dietary supplements. These individuals place greater importance on these factors, which play a decisive role in their diet choices. The literature supports this finding. The study by Cialdini and Trost emphasizes the important role of social norms and the meeting of individual needs in people's diet choices.²³ Individuals who start a diet to gain weight place more importance on the meeting of their needs because it increases their motivation and adherence to the diet. The study by Puhl and Brownell shows that ethical standards play a significant role in individuals' attitudes toward diet programs and dietary supplements. Wing and Phelan highlight the importance of monitoring processes in the success of diet programs. Individuals participating in weight gain programs are more sensitive to regular monitoring and feedback processes because these processes ensure healthy and sustainable weight gain.²⁴ Jeffrey and Wing demonstrate that incentive methods increase individuals' adherence to diet programs. The weight gain group is more sensitive to incentive methods because they increase their motivation and strengthen their adherence to the diet.²⁵

In addition, the most frequently used food supplements in the study were Vitamin D, Vitamin B, Omega 3, and multivitamin complex. A study by Bailey et al. found that multivitamin-mineral products and fish oils were the most commonly reported food supplements.¹³ Supplement use was also found to be related to more appropriate health and lifestyle choices, and less than a quarter of supplements were recommended by a

doctor or dietitian. For this reason, it is recommended that food supplements should be used consciously in line with the recommendation of a doctor or dietitian. Similarly, Çoşkun and Turhan argued that if individuals do not have sufficient knowledge about vitamins, they should act by getting information from experts.²⁶

In the information on food supplements, pricing and obtaining legal permissions, not observing ethical values, pharmacy prices not being the same as internet prices, being sold without a prescription, and insufficient information written on it are the results reached within the scope of this study and are not ethically appropriate. Acting in line with ethical principles in this regard will prevent both individual and social damages that may occur.

Conclusions

This study revealed that factors such as gender, age, and dietary preferences influence individuals' supplement choices. Men were found to be more inclined toward the meeting needs and promotion aspects of dietary supplement purchases than women. In addition, factors such as age group and dietary preferences were found to affect the sensitivity of individuals in their supplement choices. It was found that individuals between the ages of 31-40 gave more importance to legal regulations, and those who were trying to gain weight were more sensitive to need fulfillment, ethical standards, monitoring, and promotion processes. In conclusion, a conscious and ethical use of dietary supplements should be encouraged. This can help adopt a credible approach to improving individuals' health and quality of life. The findings may also contribute to health management practices by supporting consumer awareness, ethical marketing strategies, and regulatory policy development regarding dietary supplements.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of human and animal subjects. The authors declare that no experiments involving humans or animals were conducted for this research.

Confidentiality, Informed Consent, and Ethical Approval. The study does not involve patient personal data nor requires ethical approval. The SAGER guidelines do not apply.

Use of artificial intelligence for generating text. The authors declare that no generative artificial intelligence was used in the writing of this manuscript.

References

1. 'Atalay D, Erge HS. Gıda takviyeleri ve sağlık üzerine etkileri. Food Health. 2018;4:98-111.
2. Ataman RP, Evren M. Gıda güvenliği ve etik. Türk Biyoetik Derg. 2020;7:169-77.
3. Briefel RR, Johnson CL. Secular trends in dietary intake in the United States. Annu Rev Nutr. 2004;24:401-31.
4. Radimer K, Bindewald B, Hughes J, Ervin B, Swanson C, Picciano MF. Dietary supplement use by US adults: data from the national health and nutrition examination survey, 1999-2000. Am J Epidemiol. 2004;160:339-49.
5. Bailey RL, Gahche JJ, Lentino CV, Dwyer JT, Engel JS, Thomas PR, et al. Dietary supplement use in the United States, 2003-2006. J Nutr. 2011;141:261-6.
6. Picciano MF, Dwyer JT, Radimer KL, Wilson DH, Fisher KD, Thomas PR, et al. Dietary supplement use among infants, children, and adolescents in the United States, 1999-2002. Arch Pediatr Adolesc Med. 2007;161:978-85.
7. Foote JA, Murphy SP, Wilkens LR, Hankin JH, Henderson BE, Kolonel LN. Factors associated with dietary supplement use among healthy adults of five ethnicities: the multiethnic cohort study. Am J Epidemiol. 2003;157:888-97.
8. Rock CL. Multivitamin-multimineral supplements: who uses them? Am J Clin Nutr. 2007;85:277-9S.
9. Harrison RA, Holt D, Pattison DJ, Elton PJ. Are those in need taking dietary supplements? A survey of 21 923 adults. Br J Nutr. 2004;91:617-23.
10. Ersoy G. Spor Performansı İçin Ergojenik Yardım Ve Besin Destekleri. VI Uluslararası Beslenme Ve Diyetetik Kongresi. Antalya; 2008.
11. Erden BF, Tanyeri P. Ülkemizde vitamin ve mineral eklentilerin akılcı kullanımı. Sürekli Tıp Eğitim Dergisi. 2004;13:411-4.
12. Karadağ NS. Aile Sağlığı Merkezine Başvuran 18-65 Yaş Arası Bireylerin Gıda Takviyesi Alım Davranışları Ve İlişkili Etmenlerin Değerlendirilmesi. Turkey: Çelebi University Faculty of Medicine; 2019.
13. Bailey RL, Gahche JJ, Miller PE, Thomas PR, Dwyer JT. Why US adults use dietary supplements. JAMA Int Med. 2013;173:355-61.
14. Bayram F, Güngör T, Bayram SS. Meslek olarak tıbbi mümessilliğin seçiminde etik bağlılık: tıbbi tanıtım ve pazarlama öğrencileri üzerine bir uygulama. Third Sect Soc Econ Rev. 2018;53:201-16.
15. Büyüktanır BÖ, Birinci G, Ömürgönülşen U. TRIPS sözleşmesi ve ilaç firmalarının sağlık hakkına ilişkin etik sorumlulukları. Hacettepe Üniv İktisadi İdari Bilimler Fak Derg. 2012;30:1-26.
16. Mannatech. Company overview [Internet]. Coppell (TX): Mannatech Inc.; 2018 [cited May 20, 2018]. Available from: <https://www.mannatech.com/welcome.php>
17. Torok CB, Murray TH. Wielding the sword of professional ethics against misleading dietary supplement claims. Glycobiology. 2008;18:660-3.
18. Alpar R. Uygulamalı İstatistik Ve Geçerlik-Güvenlilik: Spor, Sağlık Ve Eğitim Bilimlerinden Örneklerle. Turkey: Detay Yayıncılık; 2010.
19. Wang Y, Neilson LC, Ji S. Why and how do consumers use dietary supplements? A systematic review and thematic analysis. Health Promot Int. 2023;38:daac197.
20. Djaoudene O, Romano A, Bradai YD, Zebiri F, Ouchene A, Yousfi Y, et al. A global overview of dietary supplements: regulation, market trends, usage during the COVID-19 pandemic, and health effects. Nutrients. 2023;15:3320.
21. Mahdavi-Roshan M, Rezazadeh A, Joukar F, Khorshidi Y, Naghipour M, Mansour-Ghanaei F. Dietary supplements consumption and its association with socioeconomic factors, obesity and main non-communicable chronic diseases in the north of Iran: the PERSIAN guilan cohort study (PGCS). BMC Nutr. 2021;7:84.
22. Dores AR, Peixoto M, Castro M, Sá C, Carvalho IP, Martins A, et al. Knowledge and beliefs about herb/supplement consumption and herb/supplement-drug interactions among the general population, including healthcare professionals and pharmacists: a systematic review and guidelines for a smart decision system. Nutrients. 2023;15:2298.
23. Cialdini RB, Trost MR. Social Influence: Social Norms, Conformity and Compliance. New York: McGraw Hill; 1998.
24. Wing RR, Phelan S. Long-term weight loss maintenance. Am J Clin Nutr. 2005;82:222-5S.
25. Jeffrey RW, Wing RR. Long-term effects of interventions for weight loss using food provision and monetary incentives. J Consult Clin Psychol. 1995;63:793-6.
26. Çoşkun F, Turhan H. İstanbul'da vitamin kullanım alışkanlıkları ve bu alışkanlıkları etkileyen faktörler üzerine bir araştırma. Marmara Pharm J. 2010;14:21-8.

Is there any effect of lycopene's preventing peritoneal adhesion formation in rats: an experimental study

¿Tiene el licopeno algún efecto en la prevención de la formación de adherencias peritoneales en ratas? Un estudio experimental

Ahmet Bilgi¹, Mustafa C. Terek², Gürkan Yiğittürk³, Dilek Taşkıran⁴, Orkun İlgen⁵, İsmet Hortu², Bulent Yilmaz⁶, Cavit Kart⁷, Fazıl Avcı^{1*}, and Mehmet Kulhan¹

¹Department of Gynecology and Obstetrics, Selçuk University Faculty of Medicine, Konya; ²Department of Gynecology and Obstetrics, Ege University Faculty of Medicine, İzmir; ³Department of Histology and Embryology, Mugla Sıtkı Kocman University Faculty of Medicine, Mugla; ⁴Department of Physiology, Ege University Faculty of Medicine, İzmir; ⁵Department of Gynecology and Obstetrics, Dokuz Eylül University Faculty of Medicine, İzmir; ⁶Department of Gynecology and Obstetrics, Recep Tayyip Erdogan University Faculty of Medicine, Rize; ⁷Department of Gynecology and Obstetrics, Karadeniz Technical University Faculty of Medicine, Trabzon. Turkey

Abstract

Objective: Effects of lycopene in prevention of intra-abdominal adhesion formation in a rat model. **Methods:** Twenty-eight rats were divided into four groups consisting of 7 rats each. Group 1 (only adhesion), Group 2 (adhesion + corn oil), Group 3 (adhesion + 5 mg/kg lycopene), Group 4 (adhesion + 20 mg/kg lycopene). Adhesion score, histopathology, vascular endothelial growth factor (VEGF) H-score, malondialdehyde (MDA), total antioxidant capacity, and VEGF values were measured. **Results:** There were significantly higher extent, severity, degree, and total adhesion scores in the control and corn-oil group than in the low lycopene and high lycopene group. VEGF H-scores were significantly lower in lycopene-given groups, regardless of dose. When the low lycopene and high lycopene groups were compared in terms of anti-VEGF H-score, no significant difference was observed. Malondialdehyde levels were statistically significantly lower in the control and high lycopene group. **Conclusions:** Biochemical parameters, histopathological examination, and adhesion scoring revealed that lycopene significantly reduced adhesion formation.

Keywords: Adhesion. Lycopene. Peritoneum. Rat. Vascular endothelial growth factor.

Resumen

Objetivo: Evaluar los efectos del licopeno en la prevención de la formación de adherencias intraabdominales en un modelo con ratas. **Métodos:** Veintiocho ratas se dividieron en cuatro grupos de siete cada uno: grupo 1, solo adhesión; grupo 2, adhesión más aceite de maíz; grupo 3, adhesión más 5 mg/kg de licopeno; y grupo 4, adhesión más 20 mg/kg de licopeno. Se midieron la puntuación de adherencia, la histopatología, la puntuación H del factor de crecimiento endotelial vascular (VEGF), el malondialdehído, la capacidad antioxidante total y los valores de VEGF. **Resultados:** Hubo puntuaciones de extensión, gravedad, grado y adhesión total significativamente más altas en los grupos control y con aceite de maíz que en el grupo con bajo y alto licopeno. Las puntuaciones VEGF H fueron significativamente más bajas en los grupos que recibieron licopeno, independientemente de la dosis. Cuando se compararon los grupos con niveles bajos y altos de licopeno en términos de puntuación H anti-VEGF, no se observaron diferencias significativas. Los niveles de malondialdehído fueron más bajos en los grupos de control y alto en licopeno, con una diferencia estadísticamente significativa. **Conclusiones:** Los parámetros bioquímicos, el examen histopatológico y la puntuación de adherencias revelaron que el licopeno redujo significativamente la formación de adherencias.

Palabras clave: Adhesión. Licopeno. Peritoneo. Rata. Factor de crecimiento endotelial vascular.

*Correspondence:

Fazıl Avcı

E-mail: fazilavci01@hotmail.com

2444-0507/© 2024 Academia Mexicana de Cirugía. Published by Permalyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 12-08-2024

Date of acceptance: 06-11-2024

DOI: 10.24875/CIRUE.M24000947

Cir Cir (Eng). 2026;94(3):287-294

Contents available at PubMed

www.cirugiaycirujanos.com

Introduction

Although our knowledge about peritoneal cavity physiology and peritoneal healing mechanisms is increasing, post-operative adhesions continue to be a problem for surgeons from different disciplines. In different studies, the direct relationship between intra-abdominal surgical interventions and adhesion formation has been revealed, and it has been stated that the most important cause of intestinal obstruction is adhesions due to previous surgeries.^{1,2} Post-operative adhesions are fibrous connections that can be seen between organs that are not normally combined with each other and are surrounded by serous membranes following injury or surgical operations.¹ Approximately 30% of intestinal obstructions develop due to intra-abdominal adhesion, and post-operative peritoneal adhesion occurs in more than 90% of all laparotomies. In addition, they may lead to post-operative mortality, morbidity, and cost increase due to their ability to extend the length of hospital stay.³ Despite advanced surgical techniques and medical treatment options, post-operative intra-abdominal adhesions are still an unsolved problem. When the literature is searched, it is seen that many experimental, clinical studies, and theoretical reports on adhesion prevention have been published since the beginning of the century. Various chemical agents used as adhesion inhibitors prevent fibrin organization by inhibiting fibroblastic proliferation. Therefore, many agents such as non-steroidal anti-inflammatory drugs, corticosteroids, calcium channel blockers, histamine antagonists, antibiotics, fibrinolytic drugs, antioxidants, and vitamins have been tried to inhibit this proliferation.⁴ Lycopene is a pigment belonging to the carotenoid family, naturally found in vegetables and fruits. Lycopene is a potent antioxidant substance; besides protecting cells from free radical damage, it strengthens the bonds between cells and improves cell metabolism.⁵⁻¹⁰ There have been reports supporting lycopene's effect on reduced risk of many diseases, including Alzheimer's disease, peripheral nerve damages,¹¹ cardiovascular disease, skin health,^{12,13} and even some types of cancer (breast and uterine).¹⁴ Lycopene also slows down the aging process with its antioxidant properties.¹⁵ In this observational rat study, we investigated the effectiveness of lycopene use in preventing adhesions after gynecological surgery.

Methods

Ethical statement

This study was approved by our University Animal Experiments Local Ethics Committee (April 26, 2017, Protocol No: 2017/011).

Study design

Twenty-eight female Wistar albino rats were included in the study. Inclusion criteria in the study were as follows: the animals were female Wistar albino species, weighing between 160 and 250 g, and survived from the beginning to the end of the study. The exclusion criteria were as follows: the animals were out of the female Wistar albino species, their weight was not between 160 and 250 g, and they died before the study was completed. Experimental animals were kept in our University Animal Laboratory at a room temperature of 24°C and a 12/12 h day and night cycle. Fed with standard rat food and water without restrictions. Each rat was anesthetized with ketamine hydrochloride (40 mg/kg IV). Before surgery, the abdominal area of the rats was shaved and wiped with 1% povidone iodine and prepared for the operation. Approximately 4 cm lower midline laparotomy was performed on the umbilical region. Before surgery, the rats were randomly divided into four groups each consisting of 7 rats. The operations performed on groups of rats were as follows:

- Control group (CG): a standard adhesion was created; no adjuvant was given. Corn oil group (CoG): After the injury, 1 mL of corn oil was administered daily by direct intraperitoneal injection method. The same dose was continued intraperitoneally for 14 days
- Low-dose lycopene group (LLG): After the injury, 5 mg/kg lycopene was administered daily by direct intraperitoneal injection method. The same dose was continued intraperitoneally for 14 days
- High-dose lycopene group (HLG): After injury, only a single dose of 20 mg/kg lycopene was administered intraperitoneally.

Lycopene is a powder compound. It has been used in a liquid-based substance, corn oil, to make lycopene suitable for intraperitoneal injection. To determine the effect of corn oil alone, the CoG was created as a separate group. The dose of lycopene was chosen based on a previous study.¹⁶

After a 4 cm abdominal incision, the uterine horns were exposed and a lesion was created on the antimesenteric surface of each uterine horn (free part of the bicornual rat uterus in the pelvic area) surface with the help of a 10-volt bipolar cautery as previously described.¹⁷ At the same time, extra adhesion was created by scraping operations on the visceral surface until serosal bleeding occurred. After the procedure was completed, the abdomen was closed continuously using 4.0 Vicryl. The rats were then allowed to recover for 2 weeks. A total of 4 animals died of lung infection before starting the experiment. Animals that died were excluded from the study. After the recovery period, the animals were sacrificed and evaluated for adhesion formations. The researchers who evaluated the adhesions had no prior knowledge of which group the rat belonged to. Adhesion scoring based on extent, severity, and resistance to applied force is summarized in table 1.^{18,19}

The sum of the three parameters was used as the total score for each group. Tissue samples taken from the peritoneum and adhesion area were sent for histopathological examination. Adhesion examples are shown in figure 1.

Histopathological examination

Tissues were fixed with formaldehyde for 48 h and washed with phosphate-buffered saline at pH 7.4. Dehydration was achieved by passing the tissues through increasing concentrations of ethyl alcohol (from 80%, 90%, 90%, 96% and to 96%) for 45 min in each ethanol series. After dehydration, specimens were cleared in xylene and embedded in paraffin wax.²⁰ Paraffin tissue blocks (5- μ m thickness) were cut using a rotary microtome. Tissue sections were stained with hematoxylin and eosin staining for histomorphological analysis.²¹ Tissue slides were visually assessing with a Nikon Eclipse 80i image analysis system. A total of 100 fields were scored per tissue in 10 random slide of view taken with a $\times 10$ objective for evaluation of tissue inflammation and edema.

Anti-vascular endothelial growth factor (VEGF) immunohistochemical staining

For immunohistochemistry (IHC), sections were incubated with H₂O₂ (3%) for 30 min to eliminate endogenous standard activity and blocked with protein

Table 1. Macroscopic adhesion scoring method

Score	Extend	Severity	Degree
0	No	No	No
1	< 25%	Filmy avascular	Detached with gentle traction
2	Between 25 and 50%	Opaque, translucent, avascular	Detached with moderate traction
3	Between 50 and 75%	Vascular or opaque	Detached with sharp traction
4	More than 75%	Opaque, thick veins available	-

The adhesion score is equal to the sum of the scores from each part of the adhesion. The highest possible score is 11




Figure 1. Adhesion examples. A: adhesions between bowel and uterine horn. **B:** fibrous adhesion between bowel and uterine horn. **C and D:** complete adhesion to the anterior abdominal wall and bowel.

block for 20 min at room temperature. Later, sections were incubated with primary antibody (anti-VEGF, Santa Cruz Biotechnology, Inc.; dilution 1/100) overnight at 4°C. Antibody detection was performed with the UltraTek HRP Anti-Polyvalent Staining System (ScyTec Inc., USA), and 3,30-diaminobenzidine was used to visualize the final product.²²

Immunostaining was evaluated semiquantitatively using H-score analysis. Immunostaining intensity was categorized under the following scores: 0 (no staining), 1 (weak staining), 2 (moderate staining), and 3 (intense staining). H-score value was derived for each specimen by calculating the sum of the percentage of cells for the cytoplasmic and nuclear immunoreaction

of the sections that were stained at each intensity category multiplied by its respective score, by means of the formula $H\text{-score} = \sum P_i (i + 1)$, where i = intensity of staining with a value of 1, 2, or 3 (weak, moderate, or strong, respectively) and P_i is the percentage of stained cells for each intensity, varying from 0% to 100%. For each slide, 10 different fields were microscopically evaluated at $\times 20$ magnification. H-score evaluations were independently performed by at least two investigators blinded to the source of the samples as well as to each other's results.

Biochemical analysis

MEASUREMENT OF LIPID PEROXIDE LEVEL

Measurement of lipid peroxides in tissue samples was made by the Uchiyama and Mihara methods.²³ Malondialdehyde (MDA) levels in the samples were calculated from the calibration curve prepared from standard solutions. Results are given as nmoL/mg protein.

Measurement of VEGF level

Measurement of VEGF levels in tissue samples was carried out with an enzyme-linked immunosorbent assay (ELISA) kit (Sun Red, Shanghai Sunred Biological Technology Co. Ltd.). First of all, standard samples were prepared. Standard prefixes of 50, 100, 200, 400, 800, and 1600 ng/L were prepared from the stock solution. 0.05 mL standard and 0.05 mL HRP-streptavidin were added to the wells determined in the ELISA plate. Samples were pipetted. The plate was covered and incubated at 37°C for 90 min. For tissue samples, 0.04 mL of sample, 0.01 mL VEGF antibody, and 0.05 mL of the HRP-streptavidin were added. The ELISA plate was incubated at 37°C for 60 min. At the end of the period, all samples were removed from the ELISA plate with a pipette and washed 3 times with washing buffer. At the end of the washing process, chromogen A (0.05 mL) and B solutions (0.05 mL) were added to the wells, and the plate was incubated at 37°C for 15-30 min. At the end of the period, the reaction was stopped by adding 0.05 mL of stop solution to the wells. The optical density of the resulting yellow color was read in a microplate reader (Thermo Scientific® Multiskan Go) at 450 nm wavelength. Results were evaluated according to the standard calibration curve.

Measurement of total antioxidant capacity (TAC)

Measurement of TAC in tissue samples was performed spectrophotometrically with a commercial kit (Rel Assay Diagnostics). According to the principle of this method, the antioxidant molecules in the studied sample cause the formation of a lighter colored compound by reducing the dark blue-green colored ABTS radical. The absorbance of the resulting color is read in the spectrophotometer at a wavelength of 660 nm.²⁴ The level of antioxidant molecules in the sample is calibrated with the stable antioxidant compound Trolox Equivalent (a vitamin E analog) present in the kit.

Statistical analysis

The Statistical Package for the Social Sciences 25 (SPSS 25) (IBM Corp. Released 2017. IBM SPSS Statistics for Windows, Version 25.0. Armonk, NY: IBM Corp. The statistical package program was used to evaluate the data. Histopathological changes between experimental groups were analyzed comparatively. Experiments were done independently in 6 replicates. Variables are expressed using mean $\pm$ standard deviation, percentage, and frequency values. Variables were evaluated after checking the preconditions for normality and homogeneity of variances (Shapiro Wilk and Levene test). For comparison of three or more groups, the one-way ANOVA test and the Tukey Honestly Significant Difference test, one of the multiple comparison tests, were used. For the significance level of the tests, $p < 0.05$ and $p < 0.01$ values were accepted.

Results

The results obtained from the present study are shown in table 2. The total adhesion scores were 5.86 ± 1.21 for the CG, 6.29 ± 1.49 for the CoG, 2.6 ± 1.14 for the LLG, and 2 ± 1.41 for the HLG. There were statistically significant differences between macroscopic adhesion scores between groups. A statistically significant decrease was found in all adhesion scores in rats given lycopene compared to groups not given lycopene, regardless of dose. There were significantly higher extent ($p < 0.05$), severity ($p < 0.05$), degree ($p < 0.05$), and total adhesion ($p < 0.05$) scores in the CG and CoG than in the LLG

Table 2. Macroscopic adhesion score, biochemical, histopathological findings, and VEGF H-score results

Variable	Mean $\pm$ standard deviation				p*	p**
	CG	CoG	LLG	HLG		
VEGF (ng/L)	283.75 $\pm$ 69.88	351.42 $\pm$ 96.47	313 $\pm$ 45.4	295 $\pm$ 86.44	p = 0.427	
TAC (mmol trolox Eq/L)	0.77 $\pm$ 0.32	0.58 $\pm$ 0.13	0.68 $\pm$ 0.37	0.74 $\pm$ 0.32	p = 0.636	
MDA (nmol/mL)	13 $\pm$ 3.82 ^a	20.48 $\pm$ 5.29 ^{a, b}	16.08 $\pm$ 4.27	11.47 $\pm$ 1.47 ^b	p = 0.005***	p ^a = 0.014*** p ^b = 0.007***
Adhesion score						
Extend	2.71 $\pm$ 0.48 ^{b, c}	2.85 $\pm$ 0.37 ^{d, e}	1.4 $\pm$ 0.54 ^{c, d}	1.4 $\pm$ 0.54 ^{b, e}	p = 0.000***	p ^b = 0.001*** p ^c = 0.001*** p ^d = 0.001*** p ^e = 0.001***
Severity	1.71 $\pm$ 0.48 ^b	2 $\pm$ 0.81 ^{d, e}	0.8 $\pm$ 0.44 ^d	0.4 $\pm$ 0.54 ^{b, e}	p = 0.001***	p ^b = 0.007*** p ^d = 0.015*** p ^e = 0.001***
Degree	1.42 $\pm$ 0.78 ^b	1.42 $\pm$ 0.78 ^e	0.4 $\pm$ 0.54	0.2 $\pm$ 0.44 ^{b, e}	p = 0.007***	p ^b = 0.029*** p ^e = 0.029***
Total	5.85 $\pm$ 1.21 ^{b, c}	6.28 $\pm$ 1.49 ^{d, e}	2.6 $\pm$ 1.14 ^{c, d}	2 $\pm$ 1.41 ^{b, e}	p = 0.000***	p ^c = 0.002*** p ^b = 0.000*** p ^d = 0.001*** p ^e = 0.000***
Histopathological findings and VEGF H-score						
Inflammation	4.21 $\pm$ 1.3	4.01 $\pm$ 2.30	3.9 $\pm$ 1.36	1.10 $\pm$ 2.1 ^f		p ^f = 0.005***
Edema	2.42 $\pm$ 0.2	2.33 $\pm$ 0.26	2.212 $\pm$ 0.72	0.11 $\pm$ 0.64 ^f		p ^f = 0.002***
VEGF H-Score	271.30 $\pm$ 2.41	265.12 $\pm$ 0.96	152.23 $\pm$ 1.10 ^g	132.56 $\pm$ 1.12 ^g		p ^g = 0.01***

*Data were analyzed using one-way ANOVA test.

**Data were analyzed using the Tukey HSD test.


***p-value was considered statistically significant.

p^a: comparison between CG and CoG.p^b: comparison between CG and HLG.p^c: comparison between CG and LLG.p^d: comparison between CoG and LLG.p^e: comparison between CoG and HLG.p^f: comparison between CG, CoG, and LLG.p^g: comparison between CG and CoG.

CG: control group; CoG: corn oil group; LLG: low lycopene group; HLG: high lycopene group; VEGF: vascular endothelial growth factor; TAC: total antioxidant capacity; MDA: malondialdehyde.

and HLG. Although some adhesion scores (severity, degree, and total adhesion score) were lower in the HLG compared to the LLG, these differences were not statistically significant. P-values were 0.73, 0.97, and 0.89, respectively. There was no significant difference between the CG and CoG in terms of all scoring. The line charts of the groups according to the adhesion score are shown in figure 2.

When histopathological findings are evaluated, a high level of inflammation was observed between the perimetrium and smooth muscles in the CG and CoG groups. Hyperemia and intense inflammatory cells have been detected in vascular structures. Moderate edema and thickening were also detected in the perimetrium. Similar to CG and CoG groups, high-level inflammation, hyperemia, and moderate edema were observed in the

**Figure 2.** Line charts according to the adhesion score of the groups.




Figure 3. A: control group (CG), black arrow; high level of inflammation in the perimetrium and between smooth muscles, arrowhead; moderate edema, h; hyperemia magnification $\times 4$. B: CG black arrow; high level of inflammation in the perimetrium and between smooth muscles, arrowhead; moderate edema, magnification $\times 10$. C: corn oil group (CoG), black arrow; high level of inflammation in the perimetrium and between smooth muscles, h; hyperemia magnification $\times 4$. D: CoG arrowhead; moderate edema, magnification $\times 10$. E: low-dose lycopene group (LLG), black arrow; high level of inflammation in the perimetrium, h; hyperemia magnification $\times 4$. F: LLG black arrow; high level of inflammation in the perimetrium, arrowhead; medium edema magnification $\times 10$. G: high-dose lycopene group (HLG) black arrow; low level inflammation in the perimetrium $\times 4$. H: HLG black arrow; low-level inflammation in perimetrium, magnification $\times 10$, hematoxylin and eosin staining.

perimetrium in the LLG. Inflammation ($p = 0.005$) and edema ($p = 0.002$) decreased statistically significantly ($p < 0.05$) in the HLG group compared to the other groups, including the LLG group. Symptoms of hyperemia almost disappeared (Table 2 and Fig. 3).

Anti-VEGF involvement was observed around the vascular structures and perimetrium in the anti-VEGF IHC staining of the uterine sections of the experimental groups. It was observed that VEGF H-score values of the CG and CoG groups were statistically significantly different from LLG ($p = 0.01$) and HLG ($p = 0.01$) groups ($p < 0.05$). VEGF H-scores were significantly lower in lycopene-given groups, regardless of dose. When LLG and HLG were compared in terms of anti-VEGF H-score, no significant difference was observed (Table 2 and Fig. 4). There was no significant difference between the groups in terms of VEGF and TAC in biochemical parameters except for MDA (Table 2). Compared to the CoG group, MDA levels were statistically significantly lower in the CG and HLG ($p < 0.05$). Line charts of the results of biochemical TAG, MDA, and VEGF markers according to the groups are shown in figure 5.

Discussion

Post-operative adhesions are a result of the cellular and biochemical response that occurs after peritoneal trauma while attempting to repair the peritoneum. Many methods have been used to reduce adhesion




Figure 4. Anti-vascular endothelial growth factor immunohistochemical staining. A: control group. B: corn oil group. C: low-dose lycopene group. D: high-dose lycopene group, magnification of large pictures is $\times 10$, thumbnails represent vascular structures belonging to groups.

formation, such as reducing the initial inflammatory response, preventing fibrin formation, increasing fibrinolysis, preventing collagen deposition, and using a barrier against adhesion formation, and many studies have been conducted on them²⁵ but today there is still no drug or method that can be used alone that can prevent adhesion formation satisfactorily. Many studies have shown that inflammatory cells and reactive oxygen species play a role in the formation of adhesion. The anti-adhesion effects of antioxidants have been tested in various animal models.²⁶ Heparin application before and after the operation has been shown




Figure 5. Line charts of the results of biochemical TAG, MDA, and VEGF markers according to groups. MDA: malondialdehyde; TAC: total antioxidant capacity and VEGF: vascular endothelial growth factor.

to have anti-adhesive efficacy in a rat adhesion model made by Başbuğ et al.²⁷ In another anti-adhesion model with quercetin, the adhesion scores of the Quercetin + Surgicel group were reported to be lower than the quercetin group.¹⁷ In this study, we tried to examine the potential effect of lycopene, a potent antioxidant, antiproliferative, anticarcinogenic, and anti-inflammatory, on adhesion formation in rats with traumatized uterine serosa.¹⁵ To our knowledge, the present study is the first study that used lycopene as an adhesion inhibitor in an animal model.

MDA is produced by cells involved in the inflammatory response. It is a by-product formed by oxygen radicals breaking down lipid-containing structures such as plasma and cell membranes. It is a parameter used in evaluating both tissue damage and inflammation severity.²⁸ In our study, a significant difference was observed between the HLG and CG in terms of MDA levels ($p = 0.007$). Although MDA levels were found to be lower in the HLG than in the LLG, this difference was not statistically significant.

VEGF is an angiogenic cytokine that participates in the process of adhesion formation through the formation of new vessels.²⁹ In this study, although there was no statistical difference between the groups in terms of VEGF release, numerically, more VEGF-positive cells were encountered in the treatment groups compared to the CG. We think that the biochemical and histopathological VEGF analysis differences between the experimental groups are due to the difference in the number of samples evaluated statistically.

Substances that prevent oxidation caused by free radicals and have the ability to capture and stabilize free radicals are called “antioxidants.” The total effect of all antioxidants in body fluids is called total TAC.³⁰ In this study, there was no significant difference between the groups in terms of TAC. Higher TAC levels were detected in the HLG but it was statistically insignificant.

Various methods have been used for grading intra-abdominal adhesions. The systems that are made according to the percentage of traumatized adherent area used by Linsky et al.³¹ and Leach et al.¹⁸ systems, consisting of three different parameters, are the most commonly used methods.³¹ We used the system of Leach et al.¹⁸ In this system, adhesions are scored in three separate categories according to type, prevalence, and easy or difficult separation. In our study, adhesion scores were significantly lower in the lycopene groups compared to the CG, which indicates the effectiveness of lycopene alone in reducing adhesion formation, in line with other studies.³² In *in vitro* studies, it was found that vitamin E has antioxidant, anti-inflammatory, anticoagulant, and antifibroblastic effects to a similar effect of Lycopene. Yetkin et al. demonstrated that intraperitoneal vitamin E injection and human amniotic membrane separately reduced post-operative adhesion in rats; however, a synergistic increase in their effects could not be demonstrated with their co-administration.³³ The results of our study are consistent with these studies.

Our study had some limitations. First, it was the route of application of the anti-adhesion agent. We chose this method in our study because the intraperitoneal route is generally used in the literature. Another limitation is that the dose of intraperitoneal lycopene required to prevent adhesion is unknown. In this study, we used the doses used in animal models of oxidative stress.

Conclusions

Histological and mechanical parameters obtained in our study suggest that lycopene, which we use as an anti-adhesion agent, is effective in preventing intra-abdominal adhesions and does not negatively affect wound healing.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of humans and animals. The authors declare that the procedures followed complied with the ethical standards of the responsible human experimentation committee and adhered to the World Medical Association and the Declaration of Helsinki.

Confidentiality, informed consent, and ethical approval. The study does not involve patient personal data. The SAGER guidelines were followed according to the nature of the study. This study was approved by our University Animal Experiments Local Ethics Committee (April 26, 2017, Protocol No: 2017/011).

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing of this manuscript.

References

- Diamond MP. Incidence of Postsurgical Adhesions. Peritoneal Surgery. Berlin: Springer; 2000. p. 217-20.
- Mavigök E, Bakacak M, Yazar FM, Bakacak Z, Yaylalı A, Boran ÖF, et al. A comparison in an experimental rat model of the effects on adhesion formation of different hemostatic methods used in abdominopelvic surgery. Ginekol Pol. 2019;90:507-12.
- Schreinemacher MH, Richard P, Bakkm EA, Van Goor H, Bouvy ND. Adhesion awareness: a national survey of surgeons. World J Surg. 2010; 34:2805-12.
- Risberg B. Adhesions: preventive strategies. Eur J Surg Suppl. 1997; 577:32-9.
- Yue YS, Song X, Ma C, Li D, Hu X, Chen F, et al. Lycopene ameliorated DSS-induced colitis by improving epithelial barrier functions and inhibiting the *Escherichia coli* adhesion in mice. J Agric Food Chem. 2024; 72:5784-96.
- Yang PM, Chen HZ, Huang YT, Hsieh CW, Wung BS. Lycopene inhibits NF- κ B activation and adhesion molecule expression through Nrf2-mediated heme oxygenase-1 in endothelial cells. Int J Mol Med. 2017;39:1533-450.
- Zamani M, Behmanesh Nia F, Ghaedi K, Mohammadpour S, Amirani N, Goudarzi K, et al. The effects of lycopene and tomato consumption on cardiovascular risk factors in adults: a grade assessment systematic review and meta-analysis. Curr Pharm Des. 2023;29:1671-700.
- Long Y, Paengkoum S, Lu S, Niu X, Thongpea S, Taethaisong N, et al. Physicochemical properties, mechanism of action of lycopene and its application in poultry and ruminant production. Front Vet Sci. 2024; 11:1364589.
- Gao X, Lin B, Chen C, Fang Z, Yang J, Wu S, et al. Lycopene from tomatoes and tomato products exerts renoprotective effects by ameliorating oxidative stress, apoptosis, pyroptosis, fibrosis, and inflammatory injury in calcium oxalate nephrolithiasis: the underlying mechanisms. Food Funct. 2024;15:4021-36.
- Salunke MR, Kala K, Mandlik DS, Ganeshpurkar A, Kulkarni R, Shinde V. Lycopene potentiates wound healing in streptozotocin-induced diabetic rats. J Diabetes Metab Disord. 2024;23:1359-70.
- Farjah GH, Mohammadzadeh S, Zarak Javanmard M. The effect of lycopene in egg shell membrane guidance channel on sciatic nerve regeneration in rats. Iran J Basic Med Sci. 2020;23:527-33.
- Zhang X, Zhou Q, Qi Y, Chen X, Deng J, Zhang Y, et al. The effect of tomato and lycopene on clinical characteristics and molecular markers of UV-induced skin deterioration: a systematic review and meta-analysis of intervention trials. Crit Rev Food Sci Nutr. 2024;64:6198-217.
- Shih CM, Hsieh CK, Huang CY, Huang CY, Wang KH, Fong TH, et al. Lycopene inhibit IMQ-induced psoriasis-like inflammation by inhibiting ICAM-1 production in mice. Polymers (Basel). 2020;12:1521.
- Shafe MO, Gumede NM, Nyakudya TT, Chivandi E. Lycopene: a potent antioxidant with multiple health benefits. J Nutr Metab. 2024;2024: 6252426.
- Kulhan NG, Kulhan M, Turkler C, Ata N, Kiremitli T, Kiremitli S, et al. Effect of lycopene on oxidative ovary-damage induced by cisplatin in rats. Gen Physiol Biophys. 2019;38:253-8.
- Burgess LC, Rice E, Fischer T, Seekins JR, Burgess TP, Sticka SJ, et al. Lycopene has limited effect on cell proliferation in only two of seven human cell lines (both cancerous and noncancerous) in an *in vitro* system with doses across the physiological range. Toxicol In Vitro. 2008;22:1297-300.
- Güney G, Kaya C, Oto G, Yıldırım S, Özdemir H, Tokmak A. Effects of quercetin and surgical for preventing adhesions after gynecological surgery: a rat uterine horn model. J Obstet Gynaecol Res. 2017;43:179-84.
- Leach RE, Burns JW, Dawe EJ, SmithBarbour MD, Diamond MP. Reduction of postsurgical adhesion formation in the rabbit uterine horn model with use of hyaluronate/carboxymethylcellulose gel. Fertil Steril 1998; 69: 415-418.
- Hooker GD, Taylor BM, Driman DK. Prevention of adhesion formation with use of sodium hyaluronate-based bioresorbable membrane in a rat model of ventral hernia repair with polypropylene mesh—a randomized, controlled study. Surgery. 1999;125:211-6.
- Çakalağaoğlu F. Tissue processing-Doku takibi. J Aegean Pathol. 2005;2:29-34.
- Ekicioğlu G, Özkan N, Şalvaaz E. Hematoksilin-eozin (hematoxylin-eosin)(H&E). Aegean Pathol J. 2005;2:58-61.
- Goldstein M, Watkins S. Immunohistochemistry. Curr Protoc Mol Biol. 2008;81:14.6.1-23.
- Mihara M, Uchiyama M. Determination of malonaldehyde precursor in tissues by thiobarbituric acid test. Anal Biochem. 1978;86:271-8.
- Erel O. A novel automated direct measurement method for total antioxidant capacity using a new generation, more stable ABTS radical cation. Clin Biochem. 2004;37:277-85.
- Moris D, Chakedis J, Rahnamai-Azar AA, Wilson A, Hennessy MM, Athanasiou A, et al. Postoperative abdominal adhesions: clinical significance and advances in prevention and management. J Gastrointest Surg. 2017;21:1713-22.
- Portz D, Elkins T, White R, Warren J, Adadevoh S, Randolph J. Oxygen free radicals and pelvic adhesion formation: I. Blocking oxygen free radical toxicity to prevent adhesion formation in an endometriosis model. Int J Fertil. 1991;36:39-42.
- Başbuğ M, Aygen E, Tayyar M, Kaya E, Narin F, Oktem O. Hyaluronic acid plus heparin for improved efficacy in prevention of adhesion formation in rat uterine horn model. Eur J Obstet Gynecol Reprod Biol. 1998; 78:109-12.
- Farhangi MA, Tajmiri S. The effects of powdered black cumin seeds on markers of oxidative stress, intracellular adhesion molecule (ICAM)-1 and vascular cell adhesion molecule (VCAM)-1 in patients with Hashimoto's thyroiditis. Clin Nutr ESPEN. 2020;37:207-12.
- Marino M, Del Bo C, Tucci M, Klimis-Zacas D, Riso P, Porrini M. Modulation of adhesion process, e-selectin and VEGF production by anthocyanins and their metabolites in an *in vitro* model of atherosclerosis. Nutrients. 2020;12:655.
- Roohbakhsh Y, Rahimi VB, Silakhori S, Rajabi H, Rahmanian-Devin P, Samzadeh-Kermani A, et al. Evaluation of the effects of peritoneal lavage with rosmarinus officinalis extract against the prevention of postsurgical-induced peritoneal adhesion. Planta Med. 2020;86:405-14.
- Linsky CB, Diamond MP, Cunningham T, Constantine B, DeCherney AH, diZerega GS. Adhesion reduction in the rabbit uterine horn model using an absorbable barrier, TC-7. The Journal of Reproductive Medicine. 1987;32(1):17-20.
- Fletcher NM, Awonuga AO, Saeed MG, Abu-Soud HM, Diamond MP, Saeed GM. Lycopene, a powerful antioxidant, significantly reduces the development of the adhesion phenotype. Syst Biol Reprod Med. 2014; 60:14-20.
- Yetkin G, Uludag M, Citgez B, Karakoc S, Polat N, Kabukcuoglu F. Prevention of peritoneal adhesions by intraperitoneal administration of vitamin E and human amniotic membrane. Int J Surg. 2009;7:561-5.

Machine learning for COVID-19 mortality prediction: enhancing cart models with node-specific odds ratios

Aprendizaje automático para la predicción de mortalidad por COVID-19: mejora de los modelos CART con razones de probabilidad específicas por nodo

Argelia Pérez-Pacheco^{1,3}, Alejandra E. Herrera-Suárez¹, Jonathan Vélez-Mata¹,
Liza M. Magdaleno-Fourlong¹, Amy D. Avendaño-Carrera¹, Catalina Casillas-Suárez²,
and Adolfo Pérez-García^{3*}

¹Research and Technological Development Unit; ²Pulmonology Service; ³Department of Research. Hospital General de México "Dr. Eduardo Liceaga", Mexico City, Mexico

Abstract

Objective: The objective of the study is to evaluate a classification and regression tree (CART) model combined with odds ratio (OR) to identify key predictors of COVID-19 mortality. Data from 1,432 patients hospitalized at Hospital General de México during the pandemic's 1st year were analyzed. **Methods:** A CART model was constructed using demographic, clinical, and laboratory data collected at admission. The model's performance was evaluated using multiple criteria, and ORs were calculated for each node to measure mortality. **Results:** Mechanical ventilation emerged as the strongest predictor of mortality. Patients intubated at admission with hospital stays under 17.5 days had the highest mortality rate (99%, OR = 80). Conversely, non-intubated patients hospitalized over 5.5 days with glomerular filtration rates above 66.5 had the lowest mortality (5%, OR = 0.26). The model showed excellent performance, with an F1 score of 0.918, accuracy of 0.903, and area under the curve of 0.955. **Conclusions:** The CART model, combined with OR calculations, offers a reliable and comprehensible tool for predicting COVID-19 mortality risk. This model provides practical utility in various healthcare settings, including resource-limited contexts, by focusing on readily available clinical parameters. The results emphasize the significant influence of mechanical ventilation on patient outcomes and the need for timely interventions in high-risk patients.

Keywords: COVID-19. Mortality prediction. Classification and regression tree model. Mechanical ventilation. Machine learning. Odds ratio.

Resumen

Objetivo: Evaluar un modelo CART con razones de probabilidad para identificar predictores de mortalidad por COVID-19 en 1432 pacientes hospitalizados. **Métodos:** Se construyó un modelo CART con datos demográficos, clínicos y de laboratorio al ingreso. Se evaluó el rendimiento y se calcularon las razones de probabilidad para medir el riesgo de mortalidad. **Resultados:** La ventilación mecánica fue el predictor más fuerte. Los pacientes intubados al ingreso con estancias de 5.5 días con filtración glomerular > 66.5 tuvieron la menor mortalidad (5%; OR: 0.26). El modelo mostró un excelente rendimiento (F1 = 0.918, precisión de 0.903, AUC = 0.955). **Conclusiones:** El modelo CART con razones de probabilidad ofrece una herramienta confiable para predecir la mortalidad por COVID-19, útil en diversos entornos, incluso con recursos limitados. Enfatiza la influencia de la ventilación mecánica y la necesidad de intervenciones oportunas en pacientes de alto riesgo.

Palabras clave: COVID-19. Predicción de mortalidad. Modelo CART. Ventilación mecánica. Aprendizaje automático. Razones de probabilidad.

*Correspondence:

Adolfo Pérez-García

E-mail: adolfo.perez@salud.gob.mx

2444-0507/© 2025 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 20-02-2025

Date of acceptance: 23-04-2025

DOI: 10.24875/CIRUE.M25000948

Cir Cir (Eng). 2026;94(3):295-303

Contents available at PubMed

www.cirugiaycirujanos.com

Introduction

In December 2019, Wuhan, China, became the epicenter of a global health crisis because of the emergence of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) virus. This novel virus, which causes coronavirus disease 2019 (COVID-19), belongs to the coronavirus family – a group of viruses capable of infecting both humans and animals. Coronaviruses are associated with several illnesses, from common colds to severe respiratory conditions, such as severe acute respiratory syndrome (SARS) and Middle East Respiratory Syndrome.¹ A frequent and severe complication of these infections is acute hypoxemic respiratory failure or acute respiratory distress syndrome (ARDS).² ARDS is a diffuse inflammatory lung injury causing low blood oxygen levels; it is defined as a PaO₂ under 60 mmHg or a SaO₂ under 88%.³ Pulmonary inflammation cytokines and other proinflammatory molecules damage the capillary endothelium and alveolar epithelium, disrupting the barrier between the capillaries and air spaces. As a result, patients with ARDS suffer severe respiratory distress that requires oxygenation therapy and respiratory support.^{4,5}

On September 25, 2022, 612 million confirmed cases and 6.5 million deaths were recorded due to COVID-19 worldwide. In Mexico alone, the global report included 7,702,809 confirmed cases and 334,958 deaths.⁶ Between 2020 and 2021, mortality in Mexico secondary to COVID-19 accounted for 47% of all deaths registered in the country, according to data from the National Institute of Statistics and Geography.

Globally, the most important risk factors influencing the progression and severity of infection include advanced age and comorbidities, such as diabetes, hypertension, cardiovascular disease, chronic lung disease, and chronic kidney disease.^{7,8} The global prevalence of overweight and obesity remains alarmingly high, affecting 62.5% of adults, with 64.1% of men and 60.9% of women falling into this category.^{9,10}

During the COVID-19 pandemic, a body mass index (BMI) > 35 kg/m² was strongly associated with an increased risk of intensive care unit (ICU) admission. Studies have noted that this BMI threshold not only heightened the likelihood of ICU admission but also increased the need for mechanical ventilation and prolonged hospital stays. Furthermore, overweight

and obese individuals were found to have a greater likelihood of developing ARDS, accompanied by higher mortality rates.¹¹⁻¹³

At the peak of the pandemic, most patients hospitalized for COVID-19 and/or ARDS required ventilatory support because of advanced disease progression. Ventilatory support is a critical intervention for managing respiratory failure by ensuring adequate oxygen delivery and gas exchange.¹⁴ Various ventilatory support methods exist, including non-invasive ventilation and invasive mechanical ventilation, each with distinct benefits and risks.¹⁵

The decision between non-invasive and invasive ventilation strategies is influenced by several factors, such as the severity of respiratory failure, patient response to initial treatment, and the presence of comorbidities. Although intubation and invasive mechanical ventilation remain contentious due to their association with high morbidity and mortality rates (reported between 81% and 97%), they often play a pivotal role in managing severe cases.^{5,16-19}

Another critical predictor of COVID-19 mortality was the duration of hospitalization. Prolonged hospital stay was found to triple the risk of mortality compared with shorter stays.²⁰ However, Sousa Neto et al.²¹ observed that longer ICU stays paradoxically served as a protective factor for certain patients. This finding suggests that extended ICU care may benefit those admitted with extreme disease severity, improving outcomes despite the high initial risk.

The rapid spread of the virus, its capacity for mutation, patient-specific factors, and the natural history of the disease have led to a severe shortage of medical resources and trained personnel. This crisis underscores the urgent need for risk stratification models to optimize healthcare delivery.

Predictive models provide a means to identify patients at higher risk of mortality, enabling timely and targeted interventions. Machine learning (ML) classifiers have proven to be valuable tools for supporting clinical decision-making. As a branch of artificial intelligence, ML leverages large raw datasets to develop high-quality predictive models. Among these, Classification and Regression Trees (CART) are ML-based AI tools that utilize statistical, probabilistic, and optimization techniques to integrate risk factors and generate profiles of patients at high risk of mortality.^{22,23} CART models have been applied in various medical domains, including survival analysis,²⁴ dyslipidemia classification,²⁵ and prognosis of heart failure.²⁶ CART can handle non-linear relationships through recursive

partitioning, creating splits based on different thresholds of the predictor variables.²⁷ Importantly, CART can manage both categorical and continuous variables without requiring extensive preprocessing, which enhances its usability in diverse contexts.²⁸

This study aimed to identify the key predictors of mortality in patients admitted to the Hospital General de México “Dr. Eduardo Liceaga” (HGMEL) during the 1st year of the COVID-19 pandemic. Using available clinical and laboratory data, the research employs an easy-to-use and interpretable model developed with accessible software capable of presenting results in a user-friendly, visual format. This approach ensures its applicability in real-world healthcare settings. Identifying predictors of mortality in hospitalized patients with COVID-19 is crucial for improving patient survival and optimizing healthcare resources. By focusing on actionable insights, this study seeks to improve healthcare service delivery and patient outcomes.

Methods

This retrospective study was conducted at the HGMEL. A total of 1432 patients of both sexes, older than 18 years, positive for SARS-CoV-2 by polymerase chain reaction, hospitalized from the second quarter of 2020 to the fourth quarter of 2021, were included. Data were collected upon admission. Demographical data and laboratory test results were registered; the latter consisted of complete blood counts, blood gases, and chemical blood analyses.

The subjects were divided into two groups according to the main outcome: survivors and non-survivors. For the comparison of the variables between groups, the appropriate statistical test was used according to the normality of the data, taking a significant difference $p > 0.05$.

Research involving human subjects complied with all relevant national regulations and institutional policies, which are in accordance with the tenets of the Helsinki Declaration (as revised in 2013), and was approved by the Research Ethics Committee of the HGMEL (Reg. No. DMC-3369-20-20-1).

CART model

The variables to introduce in the model were selected following a literature survey and consensus among experts regarding the most pertinent patient characteristics that could predict mortality. The variables introduced were sex, age, days of hospital stay,

BMI, type 2 diabetes mellitus, systemic arterial hypertension, glucose (mg/dL), cardiac rate in beats per minute (CR), respiratory rate in breaths per minute (RR), oxygen saturation in percentage (SpO₂), lactate dehydrogenase (LDH) in IU/L, glomerular filtration rate (GFR), hemoglobin (g/dL), leukocytes ($\times 10^3$), neutrophil/leukocyte ratio (NLR), invasive ventilatory support, and rejection of invasive ventilatory support. Subsequently, the variables underwent an analysis for multicollinearity employing the variance inflation factor approach, with all scoring between 1 and 5. Three variables, specifically BMI, HR, and RR rate, exhibited more than 10% of missing values (10.2, 12.9, and 12.8, respectively). These missing values amounts were considered acceptable on the one hand because even with some missing values, the sample size maintains sufficient statistical power for analysis, and on the other hand because CART models can handle missing data through surrogate splits, a mechanism that uses alternative splitting rules when the primary splitting variable is missing, and maintains all observations in the analysis, even those with missing values,²⁹ therefore identifying important relationships between variables that could be obscured by imputation.³⁰ The tree was built considering *deceased* as the dependent variable. The growth method selected was CART with a 10-fold cross-validation as the training method. The maximum tree depth was three levels, with 100 minimal subjects in parent nodes and 50 minimal subjects in child nodes. The Gini impurity measure was used as the splitting criterion, with 0.0001 as the minimum improvement required for a split. The prognostic value for mortality was recorded as a new variable to build a receiver operating characteristic curve. Tetrachoric tables with true/false positives and negatives were used to calculate the Odds Ratio (OR) of each node and the overall sensitivity, specificity, F1 score, positive and negative predictive values (PPV and NPV), positive and negative likelihood ratios (LR+ and LR-), area under the curve (AUC), and accuracy of the model.

All statistical data were analyzed using IBM Statistical Package for the Social Sciences (SPSS)[®] version 27.

Data availability statement

The data that support the findings of this study are openly available in Figshare at DOI: 10.6084/m9.figshare.28256057.³¹ The dataset consists of a csv file. The data can be accessed without restrictions

and are provided under the Creative Commons Attribution 4.0 International License. Any additional information required to reanalyze the data reported in this paper is available from the corresponding author upon request.

Results

A total of 1432 data from confirmed patients of COVID-19 were evaluated, among whom 583 (41%) survived and 849 (59%) died. The sex ratio differed significantly between sexes, with 936 males (66%) and 496 females (34%), $p < 0.001$. The mean age $\pm$ standard deviation of patients who died was higher than that of survivors. SpO_2 on admission was significantly lower among patients who died compared to survivors ($p < 0.001$). Among the biochemical findings, elevated LDH levels were observed in patients who died compared with those who survived ($p \leq 0.001$). The NLR was significantly higher in patients who died than in survivors ($p \leq 0.001$). Regarding the number of days of hospital stay, there was no significant difference between the groups. Interestingly, BMI showed no difference between survivors and non-survivors ($p = 0.167$). Demographic data are presented in table 1.

Figure 1 shows the obtained classification tree, which identified patients with a high-risk profile for mortality based on the risk factors present upon admission.

Patients intubated on admission who spent < 17.5 days (Node 9) had the highest mortality (466/470; 99%), with an OR of 80 (confidence interval [CI] 95% = 12.84-32.82, $p < 0.001$). Contrastingly, non-intubated patients who stayed more than 5.5 days and had a $\text{GFR} > 66.5$ (Node 14) had the lowest mortality (5/95; 5%). Moreover, this profile served as a protective factor (OR = 0.26, CI 95% = 0.22-0.32, $p < 0.001$).

The overall performance achieved by the tree was F1 score = 0.918, ACC (accuracy) = 0.903, AUC = 0.955, CI 95% = 0.944-0.965, $p < 0.001$ (Fig. 2). The other performance metrics were PPV = 0.917, NPV = 0.883, $\text{LR}^+ = 7.686$, and $\text{LR}^- = 0.091$.

Discussion

CART models provide a means of identifying high-risk subgroups to whom prevention and intervention efforts can be targeted, making them useful tools for addressing research questions that cannot be answered by traditional regression methods.³²

Our model effectively captures the complex, non-linear relationship between length of stay and survival, particularly when considering ventilation status and subsequent physiological parameters.

Similar studies

Previous studies have compared ML algorithms to predict mortality in COVID-19 patients. Zakariaee et al.³³ used Weka 3.9.2 to compare various algorithms for mortality prediction in an 815-subject database, finding that random forest performed best with 97.2% accuracy, 100% sensitivity, 94.8% precision, 94.5% specificity, 97.3% F-score, and 99.9% AUC. Other ML algorithms with an AUC from 81.2% to 93.9% also showed good performance.

Mohammadi-Pirouz et al.³⁴ developed three classification tree algorithms (Chi-squared Automatic Interaction Detector [CHAID], C5.0, and CART) for 5080 COVID-19-positive subjects using R 4.2.1 and SPSS 26. All models indicated that factors such as ICU hospitalization, intubation, age, kidney disease, BUN, CRP, WBC, NLR, SpO_2 , and hemoglobin influenced mortality rates. CART performed best, achieving 0.75 sensitivity, 0.91 specificity, 0.36 precision, 0.90 accuracy, 0.41 F-score, and 0.92 AUC.

A Mexican study³⁵ evaluated four ML algorithms in 580,570 COVID-19 patients, namely logistic regression, Naive Bayes, decision tree, and random forest, obtaining accuracy values of 0.91, 0.89, 0.91, and 0.91, respectively. An AUC of 0.92 was reported without specifying the algorithm.

These studies undoubtedly provide valuable insights into the application of ML models for COVID-19 prognosis. However, many are limited by small sample sizes and lack comprehensively report of all performance metrics.

Huyut and Üstündağ studied 686 patients in Turkey focused on COVID-19 diagnosis and prognosis using blood-gas data and the CHAID tree algorithm.³⁶ They achieved a total accuracy of 65.0% in predicting the prognosis of the disease and 68.2% in diagnosing the disease. Other performance metrics were not reported. In this work, the model was developed in SPSS version 25.

ORs

Previous studies using CART trees have used ORs for single features.³⁷ However, no similar study

Table 1. Demographics all figures represent means $\pm$ SD compared by Mann-Whitney's U except where indicated

Variable	Overall (n = 1347)	Survivors (n = 583)	Non-survivors (n = 764)	p
*Sex Female, n (%)	458 (34)	211 (36)	247 (32)	< 0.001
AGE	55.31 $\pm$ 14.51	50.72 $\pm$ 13.54	58.82 $\pm$ 14.25	< 0.001
**Days of hospital stay	11 (6-17)	11 (7-15)	11 (5-18)	0.865
BMI	28.22 $\pm$ 5.42	28.26 $\pm$ 5.14	28.18 $\pm$ 5.62	0.436
*DM, n (%)	505 (40)	176 (39)	329 (61)	0.003
*SAH, n (%)	778 (60)	336 (43)	442 (57)	< 0.001
Glucose	173.41 $\pm$ 135.53	146.02 $\pm$ 128.99	194.40 $\pm$ 136.75	< 0.001
CR	94.01 $\pm$ 19.31	89.17 $\pm$ 16.3	96.69 $\pm$ 20.3	< 0.001
RR	26.07 $\pm$ 7.37	23.14 $\pm$ 4.87	27.7 $\pm$ 8.0	< 0.001
SpO ₂	75.52 $\pm$ 15.82	81.94 $\pm$ 10.30	70.75 $\pm$ 17.44	< 0.001
Serum LDH IU/L	506.74 $\pm$ 390.26	415.80 $\pm$ 170.44	577.74 $\pm$ 487.04	< 0.001
GFR	86.63 $\pm$ 53.19	101.68 $\pm$ 47.41	75.54 $\pm$ 54.51	< 0.001
HB	14.03 $\pm$ 3.1	14.42 $\pm$ 3.04	13.74 $\pm$ 3.11	0.021
Leukocytes	11.02 $\pm$ 6.67	9.02 $\pm$ 5.55	12.51 $\pm$ 7.04	< 0.001
NLR	14.36 $\pm$ 16.34	8.84 $\pm$ 7.84	18.58 $\pm$ 19.57	< 0.001
*Invasive ventilatory support, n (%)	708 (52)	59 (8)	649 (92)	< 0.001

*Chi-squared.

**Medians (Interquartile range) compared by the median test.

BMI: body mass index; DM: diabetes mellitus; SAH: systemic arterial hypertension; CR: cardiac rate; RR: respiratory rate; SpO₂: oxygen saturation; LDH: lactate dehydrogenase; GFR: glomerular filtration rate; HB: hemoglobin; NLR: neutrophil-to lymphocyte ratio; SD: standard deviation.

(supervised ML for COVID-19 mortality) combines the calculation of the OR of each profile. This combination significantly enhances the model's utility by:

- Risk quantification, as each patient profile has a precise OR with CIs
- Terminal nodes with specific risk profiles: For non-ventilated patients hospitalized for 5.5 days or less, OR was 0.56, thus acting as a protective factor (Node 7), while for those with 80.5 or less SpO₂ (Node 11), the OR for mortality was 3.36 (77% probability of dying).

While the standard statistical comparison of days of hospital stay found no differences between survivors and non-survivors, the tree and OR calculation showed this variable's importance. For intubated subjects with hospital stays $\leq$ 25.5 days, the OR was 20.53 (CI 95% = 12.84-32.82, $p < 0.001$), whereas for those with longer stays, the OR reduced significantly (1.36, CI 95% = 0.92-2.00, $p < 0.001$). This represents a 58% difference in the probability of dying.

This OR-enhanced analysis strengthens the model's value as a clinical decision support tool, providing clinicians with precise risk estimates for different patient subgroups.

Variables

While other studies found laboratory values, comorbidities, or demographic factors as primary predictors, this model identified mechanical ventilation as the strongest discriminator, even when glucose, hemoglobin, LDH, neutrophils, and lymphocytes were introduced. Prior research has also recognized mechanical ventilation as a critical factor in predicting mortality among patients with COVID-19.^{21,38,39} Our study quantifies the dramatic impact of mechanical ventilation, with an OR of 7.55 for ventilated patients compared to 0.26 for non-ventilated patients.

The model's reliance on easily available parameters (mechanical ventilation, days of stay, GFR, SpO₂)




Figure 1. Classification and regression tree obtained. Improvement is represented as the Gini index. OR is shown for each profile.

represents a significant practical advantage, as all variables are routinely monitored during standard care without requiring highly specialized tests or extensive

patient histories. This emphasis on easily obtainable clinical data, combined with the model's strong performance metrics, makes it valuable for real-world




Figure 2. Receiver operating characteristic curve obtained. Area under curve = 0.955, confidence interval 95% (0.944-0.965), $p < 0.001$. Sensitivity = 0.91. Specificity = 0.88.

implementation across various healthcare settings, including resource-limited environments.

Voluntarily rejected mechanical ventilation

We observed that during the period of study, certain patients (or their responsible relatives on their behalf) rejected mechanical ventilation, mainly because of beliefs about COVID-19 and its treatments, shaped by misinformation or distrust in healthcare systems. Research has shown that health misinformation negatively affects individuals' decisions, leading to poor health outcomes and continued viral spread.⁴⁰ These beliefs may be influenced by fear and uncertainty in the context of the pandemic and the impact of social media, where even physicians posted misleading information.⁴¹ In our study, a small number of patients rejected mechanical ventilation (86/1432; 6%); however, this decision had a striking impact on their outcome, given that none of them survived (Node 4, 0/86; OR not calculable). Recognizing this finding is crucial for evaluating the impact of misinformation and its outcomes within the context of pandemic outbreaks and could facilitate better-informed discussions on shared decision-making with patients and their families.

Metrics

The model exhibited strong discriminative performance, with balanced accuracy in both positive and

negative predictions. It achieved a high sensitivity of 91.97% and specificity of 88.03%, reflecting its excellent capability to accurately identify both mortality and survival cases. Furthermore, the model's PPV of 91.76% and NPV of 88.34% demonstrate its reliability in predicting outcomes. Overall, the model's predictions are accurate approximately 90% of the time, regardless of the outcome.

The LR+ value of 7.686 indicates that a positive prediction is nearly 8 times more likely in true positive cases, whereas the very low LR- value of 0.0912 suggests that false negatives are rare. These values demonstrate strong diagnostic performance.

The F1 score (91.86%) and overall accuracy (90.36%) were fairly high, indicating that the model performed consistently well across all prediction aspects.

Although most published models limit their reporting to basic metrics such as accuracy, sensitivity, and specificity, our model provides a complete performance assessment through nine distinct metrics. Furthermore, the combination of node-specific ORs with these metrics (PPV, NPV, AUC) provides a thorough validation framework that surpasses the partial reporting common in published studies.

Compared with other clinical prediction models, these metrics indicate superior performance, suggesting that the CART model, which also accounts for intubation refusal, could be a highly reliable tool for COVID-19 mortality prediction, risk stratification, and resource allocation planning.

Limitations

This model needs to be tested using data from different centers with lower mortality rates. Potential temporal variations in COVID-19 treatment should be considered. Decisions on ventilation are not exempt from bias. Ultimately, the potential for selection bias must be acknowledged, as the study aimed to identify high-risk profiles during the initial phase of the pandemic; however, the selected participants may not accurately reflect the wider population due to the specific criteria set for inclusion. Finally, there exist confounding variables that may not have been considered in this investigation, including socioeconomic factors or discrepancies in healthcare practices.

Conclusions

The early identification of patients at high risk of mortality poses a significant challenge in medical

practice, especially in the context of SARS-CoV-2 infection, where timely intervention is critical.

By combining the CART model with the OR calculation of the profiles, our approach offers simple performance and interpretation, quickness, accessibility, and reliability, rendering it particularly suitable for implementation in resource-constrained healthcare settings, where the usage of perhaps more precise but indeed more complex tools for AI models (e.g. Python or R) represent a substantial complication, as they require much more specialized knowledge for installation, programming, and interpretation. By promptly identifying high-risk individuals, healthcare professionals can prioritize interventions and allocate resources more effectively, ultimately improving patient outcomes in the face of SARS-CoV-2 infection. Furthermore, insight into patients who voluntarily rejected mechanical ventilation provides real-world evidence of the impact of misinformation and supports clinical teams in counseling patients and families about prognosis.

Acknowledgments

Raúl Serrano-Loyola for the resources provided and Lucía Monserrat Pérez-Navarro, Diana Sofia Cabrera-Orejuela, Daniel E. Colli-Trejo, Melanie A. Mafara-Castillejos, Karla R. Rodríguez-Francisco, Carlos E. Maldonado-Barrientos, and Abraham Meneses-Juarez for data handling.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of humans and animals. The authors declare that no experiments involving humans or animals were conducted for this research.

Confidentiality, informed consent, and ethical approval. The authors have obtained approval from the Ethics Committee for the analysis of routinely obtained and anonymized clinical data, so informed consent was not necessary. Relevant guidelines were followed.

Declaration on the use of artificial intelligence.

The authors declare that artificial intelligence was used in the writing of this manuscript. Paperpal was used throughout the manuscript to correct and improve English writing.

References

1. Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med*. 2020;382:727-33.
2. Meng L, Qiu H, Wan L, Ai Y, Xue Z, Guo Q, et al. Intubation and ventilation amid the COVID-19 outbreak: wuhan's experience. *Anesthesiology*. 2020;132:1317-32.
3. Lagina M, Valley TS. Diagnosis and management of acute respiratory failure. *Crit Care Clin*. 2024;40:235-53.
4. Yang X, Yu Y, Xu J, Shu H, Xia J, Liu H, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med*. 2020;8:475-81.
5. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA*. 2020;323:1061-9.
6. Zhang J, Dong X, Liu G, Gao Y. Risk and protective factors for COVID-19 morbidity, severity, and mortality. *Clin Rev Allergy Immunol*. 2022;64:90-107.
7. Najera H, Ortega-Avila AG. Health and institutional risk factors of COVID-19 mortality in Mexico, 2020. *Am J Prev Med*. 2021;60:471-7.
8. Casillas N, Ramón A, Torres AM, Blasco P, Mateo J. Predictive model for mortality in severe COVID-19 patients across the six pandemic waves. *Viruses*. 2023;15:2184.
9. Okunogbe A, Nugent R, Spencer G, Powis J, Ralston J, Wilding J. Economic impacts of overweight and obesity: current and future estimates for 161 countries. *BMJ Glob Health*. 2022;7:e009773.
10. Murray CJ, Aravkin AY, Zheng P, Abbafati C, Abbas KM, Abbasi-Kangevari M, et al. Global burden of 87 risk factors in 204 countries and territories, 1990-2019: a systematic analysis for the global burden of disease study 2019. *Lancet*. 2020;396:1223-49.
11. Földi M, Farkas N, Kiss S, Zádori N, Váncsa S, Szakó L, et al. Obesity is a risk factor for developing critical condition in COVID-19 patients: a systematic review and meta-analysis. *Obes Rev*. 2020;21:e13095.
12. Lighter J, Phillips M, Hochman S, Sterling S, Johnson D, Francois F, et al. Obesity in patients younger than 60 years is a risk factor for COVID-19 hospital admission. *Clin Infect Dis*. 2020;71:896-7.
13. Gong MN, Bajwa EK, Thompson BT, Christiani DC. Body mass index is associated with the development of acute respiratory distress syndrome. *Thorax*. 2010;65:44-50.
14. Abd-Elsayed, Alaa, Mena Abdelmalak, Kevin W. Chung, 'Mechanical Ventilation', in Alaa Abd-Elsayed (ed.), Miscellaneous, in Alaa Abd-Elsayed (ed.), *Advanced Anesthesia Review* (New York, 2023; online edn, Oxford Academic, 1 Mar. 2023), <https://doi.org/10.1093/med/9780197584521.003.0359>
15. Pierson DJ. Complications associated with mechanical ventilation. *Crit Care Clin*. 1990;6:711-24.
16. Weiss P, Murdoch DR. Clinical course and mortality risk of severe COVID-19. *Lancet*. 2020;395:1014-5.
17. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020;395:1054-62.
18. Du RH, Liu LM, Yin W, Wang W, Guan LL, Yuan ML, et al. Hospitalization and critical care of 109 decedents with COVID-19 pneumonia in Wuhan, China. *Ann Am Thorac Soc*. 2020;17:839-46.
19. Namendys-Silva SA. Respiratory support for patients with COVID-19 infection. *Lancet Respir Med*. 2020;8:e18.
20. Chalkias A, Huang Y, Ismail A, Pantazopoulos I, Papagiannakis N, Bitterman B, et al. Intubation decision based on illness severity and mortality in COVID-19: an international study. *Crit Care Med*. 2024;52:930-41.
21. Sousa Neto AL, Mendes-Rodrigues C, Pedrosa RD, Röder DV. Revisiting the COVID-19 pandemic: mortality and predictors of death in adult patients in the intensive care unit. *Life (Basel)*. 2024;14:1027.
22. Singhal A, Shukla G. Decision Trees: Classification and Regression Trees (CART). Bhubaneswar: National Institute of Science Education and Research (NISER); 2023. p. 16.
23. Karthikeyan A, Garg A, Vinod PK, Priyakumar UD. Machine learning based clinical decision support system for early COVID-19 mortality prediction. *Front Public Health*. 2021;9:626697.
24. Dauda K. Development and modeling of decision tree for survival data with multiple events using deviance and cox-snell residuals within node homogeneity technique. *Int J Adv Heal Sci Technol*. 2022;2:183-90.

25. Costanza MC, Paccaud F. Binary classification of dyslipidemia from the waist-to-hip ratio and body mass index: a comparison of linear, logistic, and CART models. *BMC Med Res Methodol.* 2004;4:7.
26. Umehara T, Katayama N, Kaneguchi A, Iwamoto Y, Tsunematsu M, Kakehashi M. Models to predict prognosis in older patients with heart failure complicated by pre-frailty and frailty: a pilot prospective cohort study. *Hong Kong Med J.* 2022;28:356-66.
27. Orfanoudaki A, Chesley E, Cadisch C, Stein B, Nouh A, Alberts MJ, et al. Machine learning provides evidence that stroke risk is not linear: the non-linear Framingham stroke risk score. *PLoS One.* 2020;15:e0232414.
28. Shim EJ, Yoon MA, Yoo HJ, Chee CG, Lee MH, Lee SH, et al. An MRI-based decision tree to distinguish lipomas and lipoma variants from well-differentiated liposarcoma of the extremity and superficial trunk: classification and regression tree (CART) analysis. *Eur J Radiol.* 2020;127:109012.
29. Hyunjoong K, Loh W. Classification trees with unbiased multiway splits. *J Am Stat Assoc.* 2001;96:589-604.
30. Twala BE, Jones MC, Hand DJ. Good methods for coping with missing data in decision trees. *Pattern Recognit Lett.* 2008;29:950-6.
31. Pérez-Pacheco A, Herrera-Suárez EA, Vélez Mata J, Avendaño-Carrera AD, Casillas-Suárez C, Pérez García A. Dataset of 1432 COVID-19 patients from Hospital General de México "Dr. Eduardo Liceaga" https://figshare.com/articles/dataset/Dataset_of_1432_COVID-19_patients_from_Hospital_General_de_Mexico_Dr_Eduardo_Liceaga/28256057?file=51881855
32. Esteban C, Arostegui I, Moraza J, Aburto M, Quintana JM, Pérez-Izquierdo J, et al. Development of a decision tree to assess the severity and prognosis of stable COPD. *Eur Respir J.* 2011;38:1294-300.
33. Zakariaee SS, Naderi N, Ebrahimi M, Kazemi-Arpanahi H. Comparing machine learning algorithms to predict COVID19 mortality using a data-set including chest computed tomography severity score data. *Sci Rep.* 2023;13:11343.
34. Mohammadi-Pirouz Z, Hajian-Tilaki K, Sadeghi Haddad-Zavareh M, Amoozadeh A, Bahrami S. Development of decision tree classification algorithms in predicting mortality of COVID-19 patients. *Int J Emerg Med.* 2024;17:126.
35. Sánchez-Gálvez AM, Sánchez-Gálvez S, Álvarez-González R, Rojas-Alarcon F. Covid-19 mortality risk prediction model using machine learning. *Comput Sist.* 2023;27:881-8.
36. Huyut MT, Üstündağ H. Prediction of diagnosis and prognosis of COVID-19 disease by blood gas parameters using decision trees machine learning model: a retrospective observational study. *Med Gas Res.* 2022;12:60-6.
37. Zimmerman RK, Balasubramani GK, Nowalk MP, Eng H, Urbanski L, Jackson ML, et al. Classification and regression Tree (CART) analysis to predict influenza in primary care patients. *BMC Infect Dis.* 2016;16:503.
38. Soares TH, De Moraes NH, Soares KP, Carvalho MM, Holanda AS, Rodrigues LF, et al. Factors associated with mortality of patients with COVID-19 on invasive mechanical ventilation: a retrospective cohort study in a university hospital in Northeastern Brazil. *An Acad Bras Cienc.* 2024;96:e20231355.
39. Li H, Ashrafi N, Kang C, Zhao G, Chen Y, Pishgar M. A machine learning-based prediction of hospital mortality in mechanically ventilated ICU patients. *PLoS One.* 2024;19:e0309383.
40. Caceres MM, Sosa JP, Lawrence JA, Sestacovschi C, Tidd-Johnson A, UI Rasool MH, et al. The impact of misinformation on the COVID-19 pandemic. *AIMS Public Health.* 2022;9:262-77.
41. Sule S, DaCosta MC, DeCou E, Gilson C, Wallace K, Goff SL. Communication of COVID-19 misinformation on social media by physicians in the US. *JAMA Netw Open.* 2023;6:e2328928.

The importance of neutrophil-to-lymphocyte ratio and monocyte-to-lymphocyte ratio to prediction severity of esophageal injury after corrosive substance ingestion

La importancia del INL y el IML para predecir la gravedad de la lesión esofágica tras la ingestión de sustancias corrosivas

Ayşe Kefeli¹  and Feyyaz Onayrılı² 

¹Department of Gastroenterology, Faculty of Medicine, Tokat Gaziosmanpaşa; ²Service of Internal Medicine, Tokat Gaziosmanpaşa University, Tokat, Turkey

Abstract

Objectives: The ability to predict the severity of corrosive injury in patients remains challenging. This study aims to explore the potential of the neutrophil-to-lymphocyte ratio (NLR) and the monocyte-to-lymphocyte ratio (MLR) as novel diagnostic markers to predict the severity of esophageal injury following corrosive substance ingestion (CSI). **Methods:** We conducted a retrospective cohort analysis on patients with CSI. Demographic data, complete blood counts, and endoscopic findings were collected and analyzed. Esophageal injuries were graded using Zargar's classification. The NLR and MLR were calculated and evaluated as potential predictors of injury severity. **Results:** The mean age, NLR, MLR, red distribution width (RDW), and platelet count were increased in the hospitalized patients. Patients with severe esophageal injury also had higher mean age, NLR, MLR, and RDW. In receiver operating characteristic analyses for patients with severe esophageal injury, the area under the curve for NLR and MLR in predicting injury severity and hospitalization was statistically significant. **Conclusions:** NLR, MLR, and RDW hold promise as easily accessible and cost-effective inflammatory markers for predicting esophageal injury severity in patients with CSI. These markers could assist healthcare providers in assessing injury extent, avoiding unnecessary esophagogastroduodenoscopy, and guiding treatment decisions.

Keywords: Caustic injury. Neutrophil-to-lymphocyte ratio. Monocyte-to-lymphocyte ratio. Red cell distribution. Esophagogastroduodenoscopy.

Resumen

Objetivo: Explorar el potencial del índice neutrófilos-linfocitos (INL) y del índice monocitos-linfocitos (IML) como nuevos marcadores diagnósticos para predecir la gravedad de la lesión esofágica tras la ingestión de sustancias corrosivas. **Métodos:** Se realizó un estudio de cohortes retrospectivo de pacientes con ingestión de sustancias corrosivas. Se recopiló y analizaron datos demográficos, hemogramas completos y hallazgos endoscópicos. Las lesiones esofágicas se calificaron utilizando la clasificación de Zargar. El INL y el IML se calcularon y evaluaron como predictores potenciales de la gravedad de la lesión. **Resultados:** La edad media, el INL, el IML, la amplitud de distribución eritrocitaria y el recuento de plaquetas aumentaron en los pacientes hospitalizados. La edad media, el INL, el IML y la amplitud de distribución eritrocitaria aumentaron en los pacientes con lesión esofágica grave. En los análisis ROC realizados en pacientes con lesiones graves en el esófago, el AUC del INL y del IML para predecir la gravedad de la lesión y la hospitalización fue estadísticamente significativo. **Conclusiones:** El INL, el IML y la amplitud de distribución eritrocitaria se muestran prometedores como marcadores inflamatorios de fácil acceso y rentables para predecir la gravedad de la lesión esofágica en pacientes con ingestión de sustancias corrosivas.

*Correspondence:

Ayşe Kefeli

E-mail: ayse.kefeli@gop.edu.tr

Date of reception: 23-09-2023

Date of acceptance: 10-12-2023

DOI: 10.24875/CIRUE.M23000949

Cir Cir (Eng). 2026;94(3):304-309

Contents available at PubMed

www.cirugiaycirujanos.com

2444-0507/© 2023 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Estos marcadores podrían ayudar a los proveedores de atención médica a evaluar la extensión de la lesión, evitar una esofagogastroduodenoscopia innecesaria y guiar las decisiones de tratamiento.

Palabras clave: Lesión cáustica. Índice linfocitos-neutrófilos. Índice monocitos-linfocitos. Distribución eritrocitaria. Esofagogastroduodenoscopia.

Introduction

Corrosive substance ingestions (CSIs) constitute a significant portion of emergency cases and pose a major health challenge, especially in developing countries, but also in developed ones. CSIs can occur either intentionally, as suicide attempts (often in adolescents and adults), or accidentally (mainly in children). The consequences can be devastating, resulting in individual suffering, disability, resource utilization, and costs. Common complications include the formation of gastrointestinal tract strictures, perforation, and hemorrhage. In addition, systemic effects such as disseminated intravascular coagulation, multi-organ system failure, and sepsis may occur. However, predicting which patients will develop such complications is a challenging task. Patient history and clinical findings are insufficient to reliably predict the severity of corrosive injury. Several studies have shown that clinical signs are not always reliable indicators of the degree of injury and the potential outcome of stricture formation.¹ Given that patient follow-up and treatment strategies depend on the severity of the injury, accurately determining the extent of damage is crucial. While esophagogastroduodenoscopy (EGD) is considered the gold standard for assessing the severity of caustic injury, it is invasive, expensive, and yields negative results in 60-82% of cases.^{2,3} Therefore, there is a pressing need for a quantifiable, rapid, inexpensive, and reproducible indicator to predict the severity of corrosive injury.

The neutrophil-to-lymphocyte ratio (NLR) and the monocyte-to-lymphocyte ratio (MLR) are inexpensive and easily accessible inflammatory indicators that can be calculated using a simple blood count. These ratios, especially MLR, reflect the balance between innate and adaptive immunity, offering a straightforward measure of immune status and inflammation level. Numerous studies in the literature have demonstrated associations between NLR and MLR with various inflammatory diseases, several types of cancer, and various liver conditions.⁴⁻⁷

However, the predictive value of NLR and MLR in assessing the severity of esophageal injury in patients

with CSIs has not yet been explored. In this study, we introduce novel diagnostic markers, NLR and MLR, for predicting the severity of CSIs.

Methods

This retrospective cohort analysis consecutively enrolled a series of patients with CSIs who were admitted to the emergency department and underwent EGD between May 2016 and December 2022. Data from medical records included patients' age, sex, duration between corrosive ingestion and hospital presentation, treatment modality, and complications. A complete blood count (CBC) was obtained upon admission, and EGD was performed within 48 h of the initial injury by experienced endoscopists. Esophageal injuries were graded using the method of Zargar et al. as follows: grade 0: normal; grade 1: mucosal edema and hyperemia; grade 2a: hemorrhagic, bullous mucosa, exudates, fibrinous membranes, or superficial ulceration; grade 2b: circumferential ulceration in addition to grade 2a; grade 3: scattered small necrotic areas, multiple and deep ulcerations; grade 3b: extensive necrosis; and grade 4: perforations.⁸ Grades 1 and 2a were defined as mild-moderate lesions, while grades 2b and 3 were defined as severe lesions. Subsequently, patients were divided into two groups: mild-moderate injury and severe injury.

General information was collected, including gender, age, and comorbidity. Laboratory blood results were obtained, including hemoglobin, platelet count (PLT), neutrophil count, lymphocyte count, monocyte count, mean platelet volume (MPV), and red distribution width (RDW). The MLR was calculated as the ratio between the absolute monocyte and lymphocyte counts, and the NLR was calculated as the ratio between the absolute neutrophil and lymphocyte counts.

Exclusion criteria included the presence of other conditions such as malignancy, hematologic disease, use of immunosuppressants, active infection within 1 month before admission, severe hepatic or renal dysfunction, major trauma, or surgery; and missing blood parameter data at admission.

All patients were categorized into two groups: mild-moderate injury and severe injury. In addition, they were further divided into two groups: hospitalized patients and discharged patients. These groups were then compared in terms of demographic and laboratory parameters.

Statistical analysis

Statistical analyses were performed using Statistical Package for Social Sciences for Windows, Version 25.0, Armonk, NY, USA: IBM Corp. Qualitative data were expressed as frequency and percentage (n [%]), while quantitative data were expressed as mean and standard deviation (SD). The Chi-square test was used for qualitative variables, one-way analysis of variance (F test) for quantitative variables, and *post hoc* Tukey honestly significant difference test for pairwise comparisons. Receiver operating characteristic (ROC) curves were generated, and the area under the curve (AUC) and 95% confidence intervals were calculated for the studied hemogram parameters. Sensitivity and specificity values were determined based on cut-off points identified as the value closest to 1 in the ROC curves. A $p < 0.05$ was considered significant.

Results

EGD was performed on 168 patients due to CSI. Among them, 123 patients exhibited corrosive damage during endoscopy. Thirty-three patients were excluded due to comorbid diseases and missing data.

Of the 90 patients included in the study, 43 were male, and 47 were female. The youngest patient was 21 years old, and the oldest was 85 years old. The mean age was 52.5 ± 14 years. The distribution of patients according to Zargar's classification and the severity of internalization are presented in table 1.

A summary of the comparison of demographic and laboratory parameters is provided in table 2. Mean age, NLR, MLR, RDW, and PLT were increased in hospitalized patients with severe injuries. Mean age, NLR, MLR, and RDW were also increased in patients with severe esophageal injuries (Table 2).

The ROC curve analysis for NLR and MLR demonstrated statistically significant predictive value for both the severity of injury and hospitalization (Tables 3 and 4, Figs. 1 and 2).

In the ROC analyses conducted for hospitalized patients, the AUC was found to be 0.821 (95%

Table 1. Endoscopic findings and the distribution of the patients according to the Zargars' classification

Grade	Outpatient	Inpatient	Total	p
Zargar Grade				0.000
Grade 1	25	0	25	
Grade 2a	19	4	23	
Grade 2b	1	18	19	
Grade 3a	0	12	12	
Grade 3b	0	11	11	
Total	45	45	90	
Severity				0.000
Mild-moderate	45	22	67	
Severe	1	22	23	
Total	46	44	90	

confidence interval: 0.72-0.91) for the selected cut-off value of 2.1 for NLR, with a sensitivity of 77.7% and specificity of 77.1%. For MLR, the AUC was found to be 0.784 (95% confidence interval: 0.68-0.88), with a sensitivity of 77.3% and specificity of 77.1% for the selected cut-off value of 0.26 (Fig. 1).

In the ROC analyses conducted for patients with severe esophageal injuries, the AUC was found to be 0.841 (95% confidence interval: 0.75-0.93) for the selected cut-off value of 2.3, with a sensitivity of 76.2% and specificity of 80.0% for NLR. For MLR, the AUC was found to be 0.796 (95% confidence interval: 0.69-0.88), with a sensitivity of 76.2% and specificity of 78.3% for the selected cut-off value of 0.27 (Fig. 2).

Discussion

Esophageal corrosive injuries are common clinical problems, often requiring immediate attention. It is essential to identify the severity of injury accurately to determine appropriate treatment and predict the prognosis of patients. However, endoscopy, the gold standard for evaluating corrosive injuries, is an invasive procedure that may not always be feasible or safe in certain clinical scenarios, such as unstable patients or those with contraindications. Therefore, the search for non-invasive markers to predict the severity of corrosive injuries is of great importance. In this study, we investigated the utility of two easily accessible and cost-effective markers, NLR and MLR, in predicting the severity of esophageal injuries in patients with CSIs. Our results indicate that both NLR and MLR are significantly associated with the severity of corrosive injuries and hospitalization.

Table 2. Comparison of the demographic and laboratory parameters of all patients

Variables	Outpatient (n = 45)	Inpatient (n = 45)	p	Mild-moderate (n = 46)	Severe (n = 46)	p
Sex (m)	19	24	0.55	32	11	0.23
Age (y)	45.2 ± 12.7	55.5 ± 10.6	0.013	46.1 ± 13.1	55.05 ± 10.7	0.015
NLR	1.81 ± 0.6	5.06 ± 3.5	0.000	2.0 ± 1.8	4.9 ± 3.3	0.000
MLR	0.24 ± 0.1	0.53 ± 0.40	0.000	0.27 ± 0.2	0.51 ± 0.34	0.007
RDW	12.8 ± 1.1	13.7 ± 2.3	0.000	12.8 ± 1	13.8 ± 2.38	0.000
PLT	231.4 ± 55.8	268.89 ± 96.2	0.000	241.8 ± 71.3	258.7 ± 89	0.12
MPV	10.3 ± 1.9	10.6 ± 1.1	0.051	9.9 ± 1.9	10.8 ± 162	0.058

NLR: neutrophil-to-lymphocyte ratio; MLR: monocyte-to-lymphocyte ratio; RDW: red cell distribution width; PLT: platelet; MPV: mean platelet volume.

Table 3. ROC curve was used to evaluate the predictive value of neutrophil-to-lymphocyte ratio and monocyte-to-lymphocyte ratio for hospitalization

Test variable (s)	AUC	Asymptotic significant	CI	Cut off	sens	1-spes
NLR	0.821	0.000	0.72-0.91	2.1	77.7	77.1
MLR	0.784	0.001	0.68-0.88	0.26	77.3	77.1

AUC: area under curve; Sig: significance; CI: confidence interval; Sens: sensitivity; Spes: specificity; NLR: neutrophil-to-lymphocyte ratio; MLR: monocyte-to-lymphocyte ratio.

Table 4. ROC curve was used to evaluate the predictive value of neutrophil-to-lymphocyte ratio and monocyte-to-lymphocyte ratio for the severity of esophageal injury

Test variable (s)	AUC	Asymptotic significant	CI	Cut-off	Sens	Spes
NLR	0.841	0.000	0.75-0.93	2.3	76.2	85.4
MLR	0.794	0.000	0.70-0.89	0.26	81.2	69.4

AUC: area under curve; Sig: significance; CI: confidence interval; Sens: sensitivity; Spes: specificity; NLR: neutrophil-to-lymphocyte ratio; MLR: monocyte-to-lymphocyte ratio.

Looking at the studies on CSI, most of them have been conducted in children, while the incidence of caustic substance ingestion in adults is not low. Each year, a total of 5,000-15,000 CSI cases are reported.⁹ CSI poses a significant health challenge, particularly in developing countries. The severity of injury and subsequent complications can vary greatly among individuals, making it crucial to identify patients at risk. Clinical signs alone are often insufficient for accurately predicting the extent of corrosive injury, and endoscopy, while considered the gold standard, has its limitations, including invasiveness and

false-negative results.¹⁰ Upper gastrointestinal endoscopy is necessary to assess the severity of the injury. However, EGD is not always easily accessible, and it is invasive, expensive, and yields false-negative results in 70% of cases.¹¹ In addition, some researchers are cautious about early EGD in cases of suspected esophageal injury due to the risk of perforation. The supine position during esophagoscopy under general anesthesia can cause reflux of the corrosive agent from the stomach to the esophagus, increasing the risk of damage. Furthermore, EGD does not always provide an accurate assessment of the degree of esophageal injury.^{12,13} Therefore, there is a need for a more easily accessible, inexpensive, and safe test to determine the severity of esophageal damage. Identifying such a test that can at least predict which patients should undergo endoscopy would be valuable.

In this study, the gender distribution between the groups was equal. Patients with severe injury and/or hospitalized patients were older than the other groups. The NLR, MLR, RDW, and PLT count showed significant differences when evaluating endoscopic severity and CBC parameters in hospitalized patients. The mean PLT count was higher in patients with mild-moderate injury and discharged patients, although it was only statistically significant for hospitalization.

RDW, NLR, MLR, and PLT count are quantitative measurements of variability that is routinely reported in CBC, widely available, inexpensive, and highly reproducible tests. RDW, calculated by dividing the MPV by the SD of erythrocytes and then multiplying by 100 to express the data as a percentage, has long been used for the differential diagnosis of anemias and has emerged as a new risk marker for many different diseases associated with acute and chronic




Figure 1. Receiver operating characteristic curve analysis for hospitalization in patient with caustic substance intake.




Figure 2. Receiver operating characteristic curve analysis for the severity of esophageal injury in patient with caustic substance intake.

inflammation.^{14,15} In our study, we found some important results. RDW was higher in hospitalized patients and patients with severe esophageal injury compared to others. We speculate that the inflammation caused by the caustic substance may shorten the half-life of erythrocytes, alter membrane characteristics, and increase RDW values.

NLR is an inexpensive inflammatory indicator that can be calculated by dividing the absolute neutrophil count by the absolute lymphocyte count from a CBC. Numerous publications in the literature have shown that NLR is associated with various inflammatory diseases, and it has become widely accepted as a useful tool for evaluating inflammatory activity.¹⁶ Azab et al. determined the normal range of NLR to be

between 1.7 and 2.28 in their study with 9427 individuals, while Aydın et al. from Turkey determined the normal range to be between 1.0 and 2.3.^{17,18} In our study, we found that NLR was significantly higher in patients with severe esophageal injury and hospitalized patients compared to the others. Furthermore, we demonstrated that an NLR over 2.1 could be used to predict hospitalization with 77.7% sensitivity and 77.7% specificity, and an NLR of 2.3 could be used to predict severe esophageal injury with 76.2% sensitivity and 85.4% specificity.

MLR has gained interest as a biomarker. As a new, inexpensive, and readily available inflammatory and infectious biomarker, MLR has been shown to be applicable in predicting and prognosticating various inflammatory diseases. MLR reflects the balance of change between innate and adaptive immunity and provides a simple indicator of immune status and inflammation level.¹⁹⁻²¹ To the best of our knowledge, no studies have reported on the predictive value of MLR for the severity of esophageal injury in patients with CSI. In our study, we found that MLR was significantly higher in patients with severe esophageal injury and hospitalized patients compared to the others. In addition, we demonstrated that MLR values above 0.26, as determined by the ROC curve, can predict hospitalization with 77.3% sensitivity and 77.1% specificity, and predict severe esophageal injury with 81.2% sensitivity and 69.4% specificity.

PLT, which can be easily and promptly evaluated upon admission in most clinical settings, was significantly decreased in hospitalized patients, although there was no thrombocytopenia in our study. There is now a large body of evidence showing that platelets play a central role in inflammatory reactions. Moreover, in some studies, thrombocytopenia has been associated with increased mortality, highlighting the beneficial roles of platelets in inflammation.²² The measurement of platelets upon admission to the emergency room may be one of the most accurate and convenient parameters for precisely assessing the prognosis of patients with caustic injury.

Regarding the relationship between age and the severity of esophageal injury, the literature presents conflicting results. In our study, we found that patients with severe esophageal injury and hospitalized patients were older than the others.^{10,23} Based on this finding, we can predict that older patients may develop more severe esophageal injury.

It is important to acknowledge certain limitations of our study, including its retrospective design and

relatively limited sample size. We could not analyze some contributing factors such as the specific caustic substances ingested. However, this study is the first one in the literature to demonstrate the correlation between NLR and MLR values and the severity of esophageal injury.

Conclusions

Although EGD is considered the gold standard for determining the severity of esophageal injury, it has several disadvantages as previously mentioned. Therefore, the necessity of endoscopy in patients with caustic substance ingestion is still controversial, and there is no definitive recommendation. Our study highlights the potential of NLR and MLR as valuable and easily accessible diagnostic markers for predicting the severity of esophageal injury in patients with CSI. By incorporating these markers into clinical practice, healthcare providers may be able to better assess the extent of injury, guide treatment decisions, and improve patient outcomes. Further research is needed to fully elucidate the mechanisms underlying the associations observed in this study and to validate our findings in larger, prospective cohorts.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of human subjects and animals. Los autores declaran que para esta investigación no se han realizado experimentos en seres humanos ni en animales.

Confidentiality, informed consent, and ethical approval. The authors have obtained approval from the Ethics Committee for the analysis of routinely collected and anonymized clinical data; therefore, individual informed consent was not required. Relevant ethical recommendations have been followed. The Research Ethics Board of the Research Institute of the Tokat Gaziosmanpasa University Faculty of Medicine approved the study (study code 23-KAEK-276), which was conducted in accordance with the Declaration of Helsinki.

Declaration on the use of artificial intelligence.

The authors declare that no generative artificial intelligence was used in the writing or creation of the content of this manuscript.

References

1. Boskovic A, Stankovic I. Predictability of gastroesophageal caustic injury from clinical findings: is endoscopy mandatory in children? *Eur J Gastroenterol Hepatol.* 2014;26:499-503.
2. Aydin E, Özcan R, Emre S, Ellicevik M, Tutuncu C, Kurugoglu S, et al. Corrosive ingestion in pediatric age group: analysis of 681 patients. *Turk Assoc Pediatr Surg.* 2012;26:26-31.
3. Gaudreault P, Parent M, McGuigan MA, Chicoine L, Lovejoy FH. Predictability of esophageal injury from signs and symptoms: a study of caustic ingestion in 378 children. *Pediatrics.* 1983;71:767-70.
4. Kumar R, Geuna E, Michalarea V, Guardascione M, Naumann U, Lorente D, et al. The neutrophil-lymphocyte ratio and its utilisation for the management of cancer patients in early clinical trials. *Br J Cancer.* 2015;112:1157-65.
5. Fan Z, Li Y, Ji H, Jian X. Prognostic utility of the combination of monocyte-to-lymphocyte ratio and neutrophil-to-lymphocyte ratio in patients with NSTEMI after primary percutaneous coronary intervention: a retrospective cohort study. *BMJ Open.* 2018;8:e023459.
6. Hemond CC, Glanz BI, Bakshi R, Chitnis T, Healy BC. The neutrophil-to-lymphocyte and monocyte-to-lymphocyte ratios are independently associated with neurological disability and brain atrophy in multiple sclerosis. *BMC Neurol.* 2019;19:23.
7. Mercan R, Bitik B, Tufan A, Bozbulut UB, Atas N, Ozturk MA, et al. The association between neutrophil/lymphocyte ratio and disease activity in rheumatoid arthritis and ankylosing spondylitis. *J Clin Lab Anal.* 2016;30:597-601.
8. Zargar SA, Kochhar R, Mehta S, Mehta SK. The role of fiberoptic endoscopy in the management of corrosive ingestion and modified endoscopic classification of burns. *Gastrointest Endosc.* 1991;37:165-9.
9. Yagmur Y, Ozturk H, Kara IH, Kiraz M. The effect of nitric oxide on esophageal alkali burns. *Turk J Gastroenterol.* 1999;10:185-9.
10. Aydin E, Beser OF, Sazak S, Duras E. Role of RDW in prediction of burn after caustic substance ingestion. *Children (Basel).* 2017;5:5.
11. Lupa M, Magne J, Guarisco JL, Amedee R. Update on the diagnosis and treatment of caustic ingestion. *Ochsner J.* 2009;9:54-9.
12. Celik B, Nadir A, Sahin E, Kaptanoglu M. Is esophagoscopy necessary for corrosive ingestion in adults? *Dis Esophagus.* 2009;22:638-41.
13. Kay M, Wyllie R. Caustic ingestions and the role of endoscopy. *J Pediatr Gastroenterol Nutr.* 2001;32:8-10.
14. Demirkol S, Balta S, Cakar M, Unlu M, Arslan Z, Kucuk U. Red cell distribution width: a novel inflammatory marker in clinical practice. *Cardiol J.* 2013;20:209.
15. Lippi G, Targher G, Montagnana M, Salvagno GL, Zoppini G, Guidi GC. Relation between red blood cell distribution width and inflammatory biomarkers in a large cohort of unselected outpatients. *Arch Pathol Lab Med.* 2009;133:628-32.
16. Zahorec R. Neutrophil-to-lymphocyte ratio, past, present and future perspectives. *Bratisl Lek Listy.* 2021;122:474-88.
17. Azab B, Camacho-Rivera M, Taioli E. Average values and racial differences of neutrophil lymphocyte ratio among a nationally representative sample of United States subjects. *PLoS One.* 2014;9:e112361.
18. Aydin I, Agilli M, Aydin FN, Kurt Y, Cayci T, Tas A, et al. The reference ranges of neutrophil-lymphocyte ratio in different age groups. *Gulhane Med J.* 2015;57:414-8.
19. Kwon BS, Jeong DH, Byun JM, Lee TH, Choi KU, Song YJ, et al. Prognostic value of preoperative lymphocyte-monocyte ratio in patients with ovarian clear cell carcinoma. *J Cancer.* 2018;9:1127-34.
20. Lin Y, Peng Y, Chen Y, Li S, Huang X, Zhang H, et al. Association of lymphocyte to monocyte ratio and risk of in-hospital mortality in patients with acute type a aortic dissection. *Biomark Med.* 2019;13:1263-72.
21. Huang Y, Liu A, Liang L, Jiang J, Luo H, Deng W, et al. Diagnostic value of blood parameters for community-acquired pneumonia. *Int Immunopharmacol.* 2018;64:10-5.
22. Koupenova M, Vitseva O, Mackay CR, Beaulieu LM, Benjamin EJ, Mick E, et al. Platelet-TLR7 mediates host survival and platelet count during viral infection in the absence of platelet-dependent thrombosis. *Blood.* 2014;124:791-802.
23. Mohammadi AB, Nahandi MZ, Ostadi A, Ghorbani A, Hallaj S. Endoscopic, laboratory, and clinical findings and outcomes of caustic ingestion in adults: a retrospective study. *Gastroenterol Hepatol Bed Bench.* 2022;15:59-65.

Referral and treatment delay for breast cancer treated at a third level oncology hospital of IMSS, 2019-2020

Retraso en la referencia y el tratamiento por cáncer de mama en un hospital oncológico de tercer nivel del IMSS, 2019-2020

Saraí García-Salas^{1*}, Patricia Pérez-Martínez², and Yazmín L. Martínez-Sánchez³

¹Occupational Medicine, Unidad Médica Familiar No. 17; ²Health Education Division, Unidad Médica de Alta Especialidad, Hospital de Oncología, Centro Médico Nacional Siglo XXI; ³Epidemiology Division, Unidad Médica de Alta Especialidad, Hospital de Oncología, Centro Médico Nacional Siglo XXI. Instituto Mexicano del Seguro Social, Mexico City, Mexico

Abstract

Objective: To compare the referral delay (Dref) and the treatment delay (Dtx) of workers with breast cancer (BrCa) treated at a third level oncology hospital (HOnco) in 2019 and 2020. **Methods:** A retrospective cohort study was carried out in workers who attended the HOnco in 2019 and 2020 diagnosed with BrCa. Both, the Dref and the Dtx were defined as the elapsed days from suspicion consultation until the referral to HOnco and from this to the beginning of treatment. The data was gathered from the clinical file. In order to compare the variables, χ^2 and U Mann y Whitney were used with $p \leq 0.05$. **Results:** With a total of 542 cases, the Dref was 21 days and the Dtx was 35 days. When comparing 2019 and 2020, significant differences were found in the Dref (17 vs. 24 days) advanced stage (32.2% vs. 51%) and primary private care (14.5% vs. 38.8%). **Conclusions:** Dref for BrCa was significantly higher for 2020.

Keywords: Breast cancer. Referral delay. Treatment delay. Temporary disability.

Resumen

Objetivo: Comparar el retraso en la referencia (Rref) y el retraso en el tratamiento (Rtx) de las trabajadoras con cáncer de mama (CaMa) tratadas en un hospital oncológico de tercer nivel (HOnco) en 2019 y 2020. **Métodos:** Se realizó un estudio de cohorte retrospectivo en trabajadoras con diagnóstico de CaMa que acudieron al HOnco en 2019 y 2020. El Rref y el Rtx se definieron como los días transcurridos desde la consulta de sospecha hasta el envío al HOnco y de este al inicio del tratamiento. Para comparar las variables se usaron χ^2 y U de Mann-Whitney, con $p \leq 0.05$. **Resultados:** En un total de 542 casos, el Rref fue de 21 días y el Rtx de 35 días. Al comparar 2019 y 2020 se encontraron diferencias significativas en Rref (17 vs. 24 días), estadio avanzado (32.2% vs. 51%) y atención privada previa (14.5% vs. 38.8%). **Conclusiones:** El Rref por CaMa fue significativamente mayor en el año 2020.

Palabras clave: Cáncer de mama. Retraso de referencia. Retraso de tratamiento. Incapacidad temporal para el trabajo.

*Correspondence:

Saraí García-Salas
E-mail: saraigarcias@imss.gob.mx
2444-0507/© 2023 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 13-12-2022
Date of acceptance: 26-10-2023
DOI: 10.24875/CIRUE.M23000950

Cir Cir (Eng). 2026;94(3):310-315
Contents available at PubMed
www.cirugiaycirujanos.com

Introduction

Cancer is one of the greatest public health challenges. Breast cancer (BC) has a worldwide incidence of 47.8 per 100,000 inhabitants and a mortality rate of 13.6 per 100,000 inhabitants.¹ Survival in highly developed populations reaches 75-90%, whereas in less developed countries it is 47-76%.² In Latin America, it is more frequent in women aged 40 to 75 years.³ The *Instituto Mexicano del Seguro Social* (IMSS) reports that patients with BC are in stages III and IV in 57.5% of cases.⁴

Medical care at the IMSS

The IMSS divides medical care into 3 levels: the first provides comprehensive and continuous medical care to the patient; the second provides diagnostic, therapeutic, or rehabilitation care according to the patient's condition; and the third consists of highly specialised medical units, with advanced technology and maximum diagnostic and therapeutic resolution capacity.⁵ The referral and counter-referral system was created to facilitate the channeling of patients between levels of care, reducing waiting periods and administrative procedures.⁶

Nearly 40% of women diagnosed with BC are of productive working age; of these, nearly 80% are salaried employees,⁷ and the prescription of temporary work disability (TWD) is part of the treatment of insured women. In 2016, Campos Raymundo⁸ reported up to 504 days of TWD due to BC at the HONco.

Types of delay in the care of oncological diseases

Several published articles describe the problem of delay in cancer care, differentiating according to the place where the delay originates:

- Patient delay: the period during which the patient has symptoms but does not seek medical assessment.
- Medical delay: the period during which health personnel do not have an accurate diagnosis of the patient's disease.
- Referral delay (Rref): the period during which the first-contact physician refers the patient to the specialist physician.
- Treatment delay (Rtx): the period during which the patient waits from the consultation with the

specialist physician until the start of definitive oncological treatment.

- Health system delay: the period during which the processes and procedures of the system cause a delay before the patient receives oncological treatment.⁹

The objective of the study was to compare the Rref and Rtx of female workers with BC treated at an HONco in 2019 and 2020.

Methods

We conducted a retrospective analytical cohort study from January 2019 to December 2020 in women diagnosed with BC at the HONco. Records from that study period were reviewed for patients with this diagnosis who also had the 1F suffix in their social security number, which generally identifies women in the worker category; students or those who, upon verification, were not workers were excluded. The referral forms from the units to the HONco, known as "referral sheets", were then searched, and those corresponding to patients treated at another third-level unit who attended the HONco only for complementary treatment were eliminated. Rref days were calculated from the date of referral to the date of the first consultation at the HONco, and Rtx days were calculated from that latter date to the date of the first day on which treatment was received. During these waiting periods, information on TWD was sought in a single subsidy and benefits system. Other data were searched for in the clinical record, such as age, clinical stage, histopathological diagnosis, medical care in the private setting, and types of treatment received. For quantitative variables, the Kolmogorov-Smirnov test was applied as a normality test, showing a non-normal distribution; these data were reported as median and interquartile range 25 to 75 (IQR₂₅₋₇₅). Qualitative variables were expressed as frequency and percentage. Hypothesis tests for 2 quantitative variables were performed using the Mann-Whitney U test, and hypothesis tests for 2 qualitative variables were performed using the chi-square test. A result was considered significant when the p value was < 0.05.

Results

A total of 542 female workers with BC were included, with a median age of 47 years (IQR₂₅₋₇₅: 41-54). The most frequent histological diagnosis was ductal carcinoma in 465 patients (86.2%). Clinical stage IIA was

the most frequent, reported in 154 patients (28.4%), followed by IIB in 103 patients (19%). It was identified that 72.8% of patients had received care for BC at another IMSS unit before arriving at the HOnco, and 25.5% had received some form of care in a private setting. Regarding TWD data, 264 insured women (48.7%) had a disability period before their first consultation at the HOnco, and the range of days granted was up to 18 days. When the patient started treatment, the number of TWD cases increased to 508 (93.7%), with a median of 9 days (IQR₂₅₋₇₅: 2-28). The treatment received upon arrival at the HOnco was radiotherapy in 228 patients (42%), optimal medical therapy in 170 (31.4%), and surgical treatment in 144 (26.6%).

The number of days from referral to the first consultation at the HOnco was 21 days (IQR₂₅₋₇₅: 12-34). The median number of days elapsed from the first consultation at the HOnco to the start of any of the treatments was 35 days (IQR₂₅₋₇₅: 18.7-65.7) (Table 1).

When comparing 2019 and 2020, the latter during the COVID-19 pandemic, it was identified that in 2019 the median Rref was 17 days (IQR₂₅₋₇₅: 9-31), whereas in 2020 it was 24 days (IQR₂₅₋₇₅: 16-37; $p = 0.000$). Female workers with TWD in 2019 numbered 147 (43.5%), and in 2020 they numbered 117 (47.8%; $p = 0.687$). The days of TWD granted during the Rref period were similar in both years (IQR₂₅₋₇₅: 0-18 days for 2019 and 0-17 days for 2020; $p = 0.626$) (Table 2).

The median number of days elapsed from the first consultation at the HOnco to the start of treatment was 37 days (IQR₂₅₋₇₅: 19-67) in 2019 and 35 days (IQR₂₅₋₇₅: 19-64; $p = 0.570$) in 2020. The TWD days granted to female workers during the Rtx period were 7 days (IQR₂₅₋₇₅: 2-28) in 2019 and 11 days (IQR₂₅₋₇₅: 3-28; $p = 0.011$) in 2020.

Regarding Rtx days by type of treatment, the most marked difference was observed in patients who received chemotherapy, since in 2020 the waiting time almost doubled (20 vs. 32 days). This was not the case for radiotherapy, for which the time was longer in 2019 (60 vs. 49 days; $p < 0.000$) (Table 3).

Cases were grouped into early stages (I, IIA, and IIB) and advanced stages (IIIA, IIIB, IIIC, and IV). In 2019, advanced stages accounted for 96 cases (32.3%), whereas in 2020 they accounted for 125 cases (51%; $p = 0.000$). For TWD days in early stages, an IQR₂₅₋₇₅ of 0-12 days was found, and in late-stage disease an IQR₂₅₋₇₅ of 0-24 days was found ($p = 0.000$). Patients were categorised according to the medical care received before referral into two groups. In the private service, there were 43 cases

Table 1. Sociodemographic, clinical, and temporary work disability characteristics in women with breast cancer

Variable	n (%) or IQR
Total	542
Age, years	47 (41-54)
Year of breast cancer diagnosis	
2019	297 (54.8)
2020	245 (45.2)
Histological diagnosis	
Ductal carcinoma	467 (86.2)
Lobular carcinoma <i>in situ</i>	29 (5.4)
Lobular carcinoma	28 (5.2)
Mucinous carcinoma	6 (1.1)
Papillary carcinoma	6 (1.1)
Medullary carcinoma	3 (0.6)
Metaplastic carcinoma	2 (0.4)
Intraductal carcinoma <i>in situ</i>	1 (0.2)
Tubular carcinoma	1 (0.2)
Clinical stage	
IA	64 (11.8)
IIA	154 (28.4)
IIB	103 (19.0)
IIIA	94 (17.3)
IIIB	61 (11.3)
IIIC	24 (4.4)
IV	42 (7.7)
Treatment received at the HOnco	
Surgical treatment	144 (26.6)
Medical treatment	170 (31.4)
Radiotherapy	228 (42.0)
Care before referral	
IMSS	395 (72.8)
Private service	138 (25.5)
Ministry of Health	6 (1.1)
Other	3 (0.6)
Referral delay, days	21 (12-34)
Treatment delay, days	35 (18.7-65.7)
TWD during the referral delay	
Yes	264 (48.7)
No	278 (51.3)
TWD during the treatment delay	
Yes	508 (93.7)
No	34 (6.3)

HOnco: Hospital de Oncología; IMSS: Instituto Mexicano del Seguro Social; TWD: temporary work disability; IQR: interquartile range.

(14.5%) in 2019 and 95 cases (38.8%) in 2020 ($p = 0.000$) (Table 4).

Discussion

The general characteristics of cancer are consistent with the global literature. The main histological diagnosis, according to Barraga-Ruiz et al.,³ is ductal

Table 2. Referral delay and history of temporary work disability in 2019 and 2020

Variables	2019	2020	p
Referral delay, days	17 (IQR ₂₅₋₇₅ : 9-31)	24 (IQR ₂₅₋₇₅ : 16-37)	0.000
Temporary work disability during referral delay			0.687
Yes	147 (43.5%)	117 (47.8%)	
No	150 (56.5%)	128 (52.2%)	
Duration of temporary work disability during referral delay, days	0 (IQR ₂₅₋₇₅ : 0-18.5)	0 (IQR ₂₅₋₇₅ : 0-17)	0.626

TWD: temporary work disability; IQR: interquartile range.

Table 3. Treatment delay and history of temporary work disability in 2019 and 2020

Variables	2019	2020	p
Treatment delay, days	37 (IQR ₂₅₋₇₅ : 19-67)	35 (IQR ₂₅₋₇₅ : 19-64)	0.570
Temporary work disability during treatment delay			0.045
Yes	284 (95.6%)	224 (91.4%)	
No	13 (4.4%)	21 (8.6%)	
Duration of temporary work disability during treatment delay, days	7 (IQR ₂₅₋₇₅ : 2-28)	11 (IQR ₂₅₋₇₅ : 3-28)	0.011
Treatment time, days			0.00
Surgical	49 (IQR ₂₅₋₇₅ : 30-70)	60 (IQR ₂₅₋₇₅ : 14-46)	
Medical	20 (IQR ₂₅₋₇₅ : 23-88)	32 (IQR ₂₅₋₇₅ : 20-87)	
Radiotherapy	49 (IQR ₂₅₋₇₅ : 10-37)	33 (IQR ₂₅₋₇₅ : 20-51)	

TWD: temporary work disability; IQR: interquartile range.

Table 4. Comparison of breast cancer care in 2019 and 2020

Variables	2019	2020	p
Clinical stages, n (%)			0.000
Early	201 (67.7)	120 (49)	
Advanced	96 (32.3)	125 (51)	
Previous care, n (%)			0.00
Private medical service	43 (14.5)	95 (38.8)	
IMSS and public institutions	254 (85.5)	150 (61.2)	

IMSS: Instituto Mexicano del Seguro Social.

carcinoma, of which we found a frequency of 86.2%. The most frequent clinical stage was IIA in both years, at 28.4%, considering that the hospital is a third-level centre.

The age reported in the national literature is 40 to 75 years. The age range of the female workers in this analysis was 41 to 54 years, with a minimum of 18 and a maximum of 77 years.³

The Rref time was 21 days, which exceeds the opportunity days for referral for treatment mentioned in NOM-041-SSA1-2011,¹⁰ which indicates that it should be 10 days. We found that it was longer in 2020, which could be due to the difference in medical care during that year because of the COVID-19 health contingency, since medical care received by patients at an IMSS unit was up to 72.8%. However, in 2019, medical care in the private medical service was 14.5%, and in 2020 it was up to 38.8%. When performing the Spearman test, we found a positive and significant correlation between Rref and clinical stage, but not between type of care and clinical stage or Rref. The factors that influenced the identification of more advanced clinical stages between the 2 years were not an objective proposed in the study, but they could be addressed in other investigations. When comparing Rref with that reported by Allgar and Neal,⁹ who conducted a questionnaire in England among patients with various types of cancer, including BC, and found 13.2 days, even this value is above that established in NOM-041-SSA1-2011. Therefore, it would be important to reconsider these cut-off points to establish the timeliness of referral for this disease, generating real indicators that guide institutions' efforts to reduce them to a minimum for the benefit of the health of patients with BC.

Regarding the clinical stages of BC received at the HOnco, in 2019 a higher percentage of patients was observed in early stages, with 83.5%, and 16.5% in advanced stages; in 2020, these figures were 68.2% and 31.8%, respectively. As mentioned by Vanni et al.¹¹ in a cross-sectional study in which they administered a questionnaire to patients with BC in Rome, patients in isolation presented sentinel lymph node involvement more frequently than those from the previous year, and this will have long-term repercussions. Maringe et al.,¹² who collected information from medical records and compared pre-pandemic data with data during the pandemic, also reported that there will be an increase in deaths per year due to cancer diagnosis. The clinical stage of the disease also played an important role in disability, since a greater number of TWD days was reported in patients with advanced stages.

Medical care received by patients was provided at an IMSS unit in up to 72.8% of cases. However, in

2019, medical care in the medical service was 14.5%, and in 2020 it reached 38.8% of insured women, playing an important role in diagnosis.

Rtx had a median of 35 days, and NOM-041-SSA1-2011 states that treatment in an oncology centre should begin in < 15 days.¹⁰ The TWD time while waiting to receive some form of treatment was 8 days in 2019 and 16 days in 2020, a statistically significant difference. The number of days exceeding the standard has repercussions on the prescription of disability and, therefore, on cost. A study conducted by *Instituto Nacional de Cancerología* in 2012 through a questionnaire administered to patients reported an Rtx of 141 days. This was not the case in the present study, since the data were obtained from primary sources, avoiding the risk of recall and measurement bias.¹³

The Rtx time varied according to the type of treatment. For medical treatment (chemotherapy), the median was 28 days; in the different medical notes, a delay was found in the performance of the interconsultation by the breast tumour service. The radiotherapy service had a median of 33 days, and in the review of the clinical record it was found that those patients referred from the second level still had a pending chemotherapy cycle, delaying the start of radiotherapy, and that the large number of patients attending this unit only for radiotherapy made the wait to start treatment longer. Surgical Rtx had a median of 51 days; procedures were affected by the performance of studies such as slide review, genotype expression, and laboratory and imaging studies. The comparison between 2019 and 2020 showed an increase in days for medical treatment and surgical treatment. Papautsky and Hamlish¹⁴ administered a questionnaire to patients and found that 30% perceived a delay in therapeutic care.

TWD time had a median of 9 days for optimal medical therapy and 14 days for radiotherapy, due to treatment effects rather than clinical stage.

Among the limitations of the study, the type of population included should be noted, since, as patients considered active workers, there are likely to be important differences compared with the rest of the patients with BC, mainly in the variable analysed regarding prior care in a private setting, since this was a group that was able to meet this need by its own means, especially in 2020, when we can consider that fear of COVID-19 infection led them to attend private and less crowded places. Likewise, because this was a third-level hospital, external validity is limited.

Conclusions

BC is a disease in which timely diagnosis is the best way to reduce mortality; therefore, the performance of a greater number of screening studies should be prioritised at the first level of care. Rref and Rtx are greater than those reported in other countries and exceed the days referred to in the regulations. For 2020, the greatest backlog was found, leaving a large workload for the following years in the different oncology services, both for this cancer and probably for the rest of cancers. The results indicate that strategies to improve the care of patients with BC should be reinforced by shortening the waiting time for consultation with the oncology specialist and the time required to perform the complementary diagnostic studies needed for therapeutic decision-making. This will also have an effect on reducing TWD days, giving female workers a greater opportunity to return to work.

Funding

The authors received no funding for the research or for the publication of this article.

Conflicts of interest

The authors declared no conflicts of interest regarding the research or publication of this article.

Ethical considerations

Protection of human subjects and animals. The authors declare that the procedures followed conformed to the ethical standards of the responsible human experimentation committee, in accordance with the World Medical Association and the Declaration of Helsinki. The procedures were authorised by the institution's Ethics Committee.

Confidentiality, informed consent, and ethical approval. The authors obtained approval from the Ethics Committee for the analysis of routinely obtained and anonymised clinical data. Due to the nature of the study, individual informed consent was not required. The relevant ethical recommendations were followed.

Declaration on the use of artificial intelligence. The authors declare that no type of generative artificial intelligence was used for the writing or creation of content for this manuscript.

References

1. Cancer Today. Francia: IARC. Disponible en: <http://gco.iarc.fr/today/home>.
2. Sankaranarayanan R, Swaminathan R, Brenner H, Chen K, Chia KS, Chen JG, et al. Cancer survival in Africa, Asia, and Central America: a population-based study. *Lancet Oncol*. 2010;11:165-73.
3. Barraga-Ruiz JA, Becerra-Alcántara GI, González-López NJ. Guía de práctica clínica: diagnóstico y tratamiento del cáncer de mama en segundo y tercer nivel de atención, evidencias y recomendaciones. México: Secretaría de Salud; 2009.
4. Sociedad Mexicana de Oncología AC. Prevención y diagnóstico oportuno en cáncer. México; 2016. Disponible en: https://www.smeo.org.mx/descargables/copredoc_guia.pdf.
5. Diario Oficial de la Federación. Norma Oficial Mexicana NOM-041-SSA2-2011, para la prevención, diagnóstico, tratamiento, control y vigilancia epidemiológica del cáncer de mama. México: DOF; 2011. Disponible en: http://dof.gob.mx/nota_detalle.php?codigo=5194157&fecha=09/06/2011.
6. Guerra y Guittens JL. Referencia y contrarreferencia de un primer nivel a un segundo y tercer nivel de atención de pacientes de la clínica Moctezuma, [tesis de especialidad]. Ciudad de México: Universidad Nacional Autónoma de México; 1995.
7. Torp S, Syse J, Paraponaris A, Gudbergsson S. Return to work among self-employed cancer survivors. *J Cancer Surviv*. 2017;11:189-200.
8. Campos-Raymundo A. Factores asociados a incapacidad temporal prolongada en trabajadoras atendidas en UMAE Hospital de Oncología CMN Siglo XXI, con diagnóstico de cáncer de mama [tesis de especialidad]. Ciudad de México: Universidad Nacional Autónoma de México; 2016.
9. Allgar VI, Neal RD. Delays in the diagnosis of six cancers: analysis of data from the national survey of NHS patients: cancer. *Br J Cancer*. 2005;92:1959-70.
10. Diario Oficial de la Federación. Reglamento de Prestaciones Médicas del Instituto Mexicano del Seguro Social. México: DOF; 2006. Disponible en: <http://www.imss.gob.mx/sites/all/statics/pdf/reglamentos/4045.pdf>.
11. Vanni G, Tazzioli G, Pellicciaro M, Materazzo M, Paolo O, Cattadori F, et al. Delay in breast cancer treatments during the first COVID-19 lockdown. A multicentric analysis of 432 patients. *Anticancer Res*. 2020;40:7119-25.
12. Maringe C, Spicer J, Morris M, Purushotham A, Nolte E, Sullivan R, et al. The impact of the COVID-19 pandemic on cancer deaths due to delays in diagnosis in England, UK: a national, population-based, modelling study. *Lancet Oncol*. 2020;21:1023-34.
13. Unger-Saldaña K, Peláez-Ballesteros I, Infante-Castañeda C. Development, and validation of a questionnaire to assess delay in treatment for breast cancer. *BMC Cancer*. 2012;12:626.
14. Papautsky EI, Hamlish T. Patient-reported treatment delays in breast cancer care during the COVID-19 pandemic. *Breast Cancer Res Treat*. 2020;184:249-54.

Liver transplant program outcomes at a Colombian center: 14 years of experience

Resultados del programa de trasplante de hígado en un centro colombiano: 14 años de experiencia

Maricel Licht-Ardila^{1*}, Giseth A. Mosquera-Velasquez², Angie V. Abril-Lopera²,
Alexandra Hurtado-Ortiz¹, Luis A. Olarte-Licht², Santiago Flórez-Suárez¹,
Lizeth C. Padilla-Padilla³, Elizabeth C. David-Gonzalez³, Joel Duarte-Yepes³,
Alejandra Mendoza-Monsalve¹, Skarlet M. Vásquez², Gabriel J. Arenas-Peña³,
Mauricio Vanegas-Ballesteros³, and Cristian E. Tarazona-Leon³

¹Department of Epidemiology, Fundación Cardiovascular de Colombia; ²Department of Health Sciences, Universidad Autónoma de Bucaramanga (UNAB); ³Department of Hepatic Transplantation, Fundación Cardiovascular de Colombia. Santander, Colombia

Abstract

Objectives: The objective of the study was to analyze outcomes of a liver transplant program over 14 years and evaluate associations between patient characteristics, surgical factors, complications, and mortality. **Methods:** A retrospective cohort study was conducted at a transplant center, including patients from 2011 to 2025. Descriptive statistics, χ^2 , Mann–Whitney U tests, and multivariate logistic regression were used to identify mortality predictors. Kaplan–Meier survival analysis was also performed. **Results:** The study included 109 liver transplant recipients. Rejection rates were higher among deceased patients (30.3% vs. 13.2%, $p = 0.034$). One-year survival improved from 63.3% (2011–2014) to 89.1% (2023–2025), with 100% rejection-free survival in the most recent cohort. Complications like multi-organ failure, bronchopneumonia, acute renal failure, ventilatory failure with coma, and leukoencephalopathy significantly increased mortality risk (odds ratio 4.27, $p = 0.002$). **Conclusions:** One-year survival has improved, with reduced rejection rates reflecting better post-transplant care. However, complications remain a major mortality risk. Shorter ischemia times indicate optimized surgical and logistical procedures.

Keywords: Liver transplantation. Mortality. Survival. Complications. Graft rejection. Tissue donors.

Resumen

Objetivos: Analizar los resultados de un programa de trasplante hepático durante 14 años y evaluar la relación entre las características del paciente, los factores quirúrgicos, las complicaciones y la mortalidad. **Métodos:** Estudio de cohorte retrospectivo en un centro de trasplantes, incluyendo pacientes entre 2011 y 2025. Se usaron estadísticas descriptivas con pruebas χ^2 y U de Mann–Whitney para identificar predictores de mortalidad. Se aplicó análisis de supervivencia de Kaplan–Meier. **Resultados:** Se incluyeron 109 receptores. El rechazo fue mayor en los fallecidos (30.3% vs. 13.2%; $p = 0.034$). La supervivencia al año mejoró del 63.3% (2011–2014) al 89.1% (2023–2025), con un 100% de supervivencia libre de rechazo en la cohorte más reciente. Las complicaciones como falla multiorgánica, bronconeumonía, insuficiencia renal aguda y falla

*Correspondence:

Maricel Licht-Ardila
E-mail: mlichtar@gmail.com

Date of reception: 03-04-2025

Date of acceptance: 01-08-2025

DOI: 10.24875/CIRUE.M25000951

Cir Cir (Eng). 2026;94(3):316-325

Contents available at PubMed

www.cirugiaycirujanos.com

2444-0507/© 2025 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

ventilatoria con coma aumentaron el riesgo de mortalidad (OR: 4.27; $p = 0.002$). **Conclusiones:** La supervivencia al año ha mejorado y la reducción del rechazo refleja un mejor cuidado postrasplante. Sin embargo, las complicaciones siguen siendo clave en la mortalidad. Los tiempos de isquemia más cortos indican optimización quirúrgica y logística.

Palabras clave: Trasplante hepático. Mortalidad. Supervivencia. Complicaciones. Rechazo del injerto. Donantes.

Introduction

Liver transplantation has become a crucial therapeutic intervention for patients with end-stage liver disease, significantly improving survival rates and quality of life.¹ Worldwide, it is estimated that more than 41,000 people received a liver transplant in 2023, with the American region being the location with the highest number of procedures (around 35%).² Since the first transplant in 1963, the procedure has evolved considerably in both surgical techniques and post-operative management.³ However, it remains a complex intervention with a high risk of complications. At present, patient survival rates at 1, 3, and 5 years are approximately 91.8%, 83.8%, and 76.1%, respectively.^{4,5} In Colombia, despite the implementation of this intervention more than four decades ago,⁶ challenges persist, including donor availability, access to immunosuppressive therapies, and the management of post-operative complications.⁷

In Colombia, organ donation and transplantation are regulated by a legal framework aimed at ensuring organ availability and improving access to transplants. The legislation follows the principle of presumed consent, meaning individuals are considered donors unless they explicitly choose otherwise. To optimize the donation process, strategies such as awareness campaigns, strengthening the transplant network, and enhancing organ procurement logistics have been implemented.⁸ However, liver donation remains a significant challenge due to multiple barriers faced by medical personnel, including insufficient training in brain death certification, difficulties in communicating with families, and limitations in hospital infrastructure. These factors negatively impact donation rates, making it essential to address them from the healthcare team's perspective to enhance access to liver transplantation and improve care for patients on the waiting list.⁹

The long-term success of this intervention largely depends on the appropriate use of immunosuppressive therapies, which prevent graft rejection and improve patient survival. Various immunosuppression protocols combine drugs such as calcineurin

inhibitors, antimetabolites, and corticosteroids, each with specific mechanisms of action aimed at modulating the immune response. The selection of an immunosuppressive regimen should be based on multiple factors, including recipient characteristics, rejection risk, and the presence of comorbidities, highlighting the importance of an individualized approach to post-transplant management.¹⁰

Common complications after liver transplantation include infections, acute graft rejection, renal dysfunction, biliary stenosis, and pulmonary or neurological disorders.¹¹ Bacterial infections, particularly in the immediate post-operative period, are a major cause of morbidity. Acute rejection usually occurs within the 1st few weeks and requires prompt treatment to preserve graft function. Renal dysfunction is often linked to nephrotoxic immunosuppressants and intraoperative hemodynamic fluctuations. Biliary stenosis, affecting around 13% of recipients, is one of the most significant complications.^{12,13} Neurological complications, though less common, range from mild manifestations to potentially life-threatening conditions, reflecting their multifactorial etiology. Given these challenges, this study aims to analyze the key outcomes of a liver transplant program over 14 years in Colombia, identifying the association between patient characteristics, surgical factors, post-operative complications, and mortality after transplantation.

Methods

A retrospective cohort study was conducted at a high-complexity institution, including patients who underwent liver transplantation between May 2011 and March 2025. The study was carried out at a liver transplant center of excellence, supported by a multidisciplinary team of hepatologists, internists, hepatobiliary surgeons, nurses, nutritionists, psychologists, social workers, psychiatrists, anesthesiologists, and nephrologists. Post-transplant follow-up was structured to ensure close monitoring and timely interventions. During the 1st month, patients were evaluated weekly, followed by biweekly assessments if recovery progressed favorably. Subsequently, follow-up visits

were scheduled monthly, then every 3 months, and eventually every 5 months as the post-transplant period lengthened. In cases of liver function abnormalities or immunosuppression imbalances, weekly assessments continued until stabilization was achieved.

Participants

The liver transplantation program includes patients who meet the Milan criteria, which specify that candidates must have either a single liver tumor measuring < 5 cm in diameter or 2-3 tumors, each measuring < 3 cm at the time of diagnosis.^{14,15} For this study, all patients enrolled in the liver transplantation program were included in the study. In addition to liver cancer cases that meet these criteria, the program also accommodates patients with other liver diseases requiring transplantation, such as cirrhosis of various etiologies (including viral hepatitis B and C, alcoholic liver disease, metabolic disorders, primary and secondary biliary cirrhosis, and autoimmune hepatitis), polycystic liver disease with associated symptoms and deterioration of quality of life, acute or fulminant hepatitis, and primary biliary cirrhosis in non-cirrhotic patients who experience significant quality of life impairment. Patients < 13 years and those with liver cancer who did not meet Milan criteria were excluded. Participant selection was based on a registry from the liver transplantation center of excellence, following a consecutive selection process to verify eligibility.

Variables

Sociodemographic variables included age (years); sex (male or female); type of social security (contributory, subsidized, particular and other); educational level (primary, secondary, technical, professional, postgraduate and no information); marital status (single, married, separated, widowed, common law union, not disclosed); area of residence (rural or urban); and occupation (homemaker, unemployed, Business person, employee, Independent, others, Disability pensioner).

Clinical variables as personal history, including alcohol consumption (daily consumption: more than seven drinks per week; occasional consumption: less than once per week; weekly consumption: 1-7 drinks per week; does not consume), smoking history (former smoker: quit more than 6 months ago; never smoked; current smoker), diabetes (including whether insulin-dependent), dyslipidemia, obesity, history of hepatitis

(A, B, C, or unspecified), heart failure, acute myocardial infarction, and cirrhosis. Family history, including liver cancer, pancreatic cancer, and other types of cancer were also measured.

Variables related to the procedure's characteristics were collected, such as ischemia times (cold and warm), transplant classification (adult or pediatric), donor type (living or deceased), and transplant outcomes such as the need for re-transplantation. Regarding complications, rejection occurrence was assessed and classified as acute, chronic, along with its potential causes, including treatment adherence, administrative factors related to insurance, subtherapeutic drug levels, or other causes. Biliary complications were also analyzed, specifically strictures, bile leaks, and other related issues.

In addition, complications were considered, including the necessity for surgical reintervention. These were categorized based on temporal occurrence, distinguishing between reintervention due to hemorrhage within the initial 48 h and reintervention within the 1st month. Furthermore, other observed complications encompassed multi-organ failure, unspecified bronchopneumonia, acute renal failure, ventilatory failure with coma, leukoencephalopathy, and acute myocardial infarction. Post-operative complications were additionally classified using the Clavien-Dindo classification, which standardizes the assessment of surgical adverse event severity.¹⁶ The primary outcome measure was all-cause mortality following liver transplantation, assessed both during hospitalization and throughout the follow-up period. Mortality was ascertained through institutional medical records and follow-up telephone communications, conducted as part of the liver transplant program's ongoing surveillance. To determine mortality, hospital records and updated medical histories were reviewed at each follow-up visit.

Data sources

The information was obtained from medical records, including surgical notes, progress notes, and outpatient consultations documented by healthcare personnel. These data are stored in the transplant center of excellence's database, maintained on the research electronic data capture platform under project PID 690.^{17,18} To minimize information bias and ensure data quality, strategies such as double data entry, cross-validation of records, and training of personnel responsible for data extraction were implemented.

Statistical analysis

The descriptive analysis was performed first, assessing normality using the Kolmogorov-Smirnov test. The median was calculated, along with the interquartile ranges. Categorical variables were presented as frequencies and percentages. A bivariate analysis was conducted using mortality as the outcome variable, applying the χ^2 test for categorical variables, and the Mann-Whitney U test for those continuous without a normal distribution. A multivariate logistic regression model was developed to identify independent predictors of mortality. Variable selection followed a backward elimination approach, initially including factors with a $p < 0.25$. Only variables with a $p < 0.05$ were retained in the final analysis. Model fit was assessed using the Hosmer-Lemeshow test and the pseudo R^2 statistic.

Furthermore, a survival analysis was performed using the Kaplan-Meier estimator to evaluate the probability of developing complications over time (mortality, graft rejection, biliary and vascular complications). Survival data of liver transplant patients were analyzed across four different periods: 2011-2014, 2015-2018, 2019-2022, and 2023-2025. Comparisons between these periods were performed using the log-rank test to assess differences in survival distributions. All analyses were conducted using RStudio software and Stata version 16 (StataCorp, College Station, TX, USA).

Results

A total of 109 patients who underwent liver transplantation were analyzed, of whom 76 (67%) survived, and 33 (33%) died, mostly of men (63.3%) with a global median of age of 53 years (range: 42-63). Most patients are affiliated with the contributory health insurance system (60.6%), followed by the subsidized system (33.9%). Regarding education, the majority completed primary or secondary school (31.2% each), while 21.1% have professional, technical (9.17%), or postgraduate education (0.91%), 41.3% were married, and 89.9% live in urban areas (Table 1).

In the bivariate analysis, the mean age was significantly higher in patients who died during the follow-up after liver transplantation compared to those who survived (59 vs. 51.5; $p = 0.031$). A higher proportion of surviving patients were affiliated with the subsidized healthcare regime (36.8%) compared to those who died (27.7%; $p = 0.031$). In addition, a greater

Table 1. Sociodemographic characteristics of patients with liver transplant

Variable	Alive n = 76 (%)	Deceased n = 33 (%)	Total n = 109 (%)	p
Sex				
Female	29 (38.1)	11 (33.3)	40 (36.7)	0.631
Male	47 (61.8)	22 (66.6)	69 (63.3)	
Age*	51.5 (40.5-60.0)	59 (47-63)	53 (42-63)	0.031
Type of security and social				
Contributory	45 (59.2)	21 (63.6)	66 (60.6)	0.031
Subsidized insurance	28 (36.8)	9 (27.7)	37 (33.9)	
Particular	3 (3.95)	0 (0.00)	3 (2.75)	
Other	0 (0.00)	3 (9.09)	3 (2.75)	
Level of education				
Primary	22 (28.9)	12 (31.1)	34 (31.2)	< 0.001
Secondary	30 (39.4)	4 (12.1)	34 (31.2)	
Technical	9 (11.8)	1 (3.03)	10 (9.17)	
Professional	15 (19.7)	8 (24.2)	23 (21.1)	
Postgraduate	0 (0.00)	1 (3.03)	1 (0.91)	
No information	0 (0.00)	7 (6.42)	7 (6.42)	
Marital status				
Single	22 (28.5)	5 (15.1)	27 (24.8)	0.386
Married	27 (35.3)	18 (54.5)	45 (41.3)	
Separated	4 (5.26)	1 (3.03)	5 (4.59)	
Widowed	2 (2.63)	2 (6.06)	4 (3.67)	
Common law union	19 (25.0)	6 (18.1)	25 (22.9)	
Not disclosed	2 (2.63)	1 (3.03)	3 (2.75)	
Area of residence				
Rural	9 (11.8)	2 (6.06)	11 (10.1)	0.357
Urban	67 (88.1)	31 (93.4)	98 (89.9)	
Occupation				
Home	21 (27.6)	10 (30.3)	31 (28.4)	0.535
Unemployed	6 (7.89)	1 (3.03)	7 (6.42)	
Business person	8 (10.53)	2 (6.06)	10 (9.17)	
Employee	12 (15.7)	3 (9.09)	15 (13.8)	
Independent	14 (18.4)	8 (24.4)	22 (20.2)	
Others	15 (19.7)	8 (24.4)	23 (21.1)	
Disability pensioner	0 (0.00)	1 (3.03)	1 (0.91)	

*Median. IQR: interquartile range.

proportion of surviving patients had secondary or technical education compared to those who died (51.2% vs. 15.1%, respectively; $p < 0.001$) (Table 1).

When evaluating the personal and family history of patients with liver transplant, alcohol consumption was reported by 49.54%, while 78.89% had never smoked. Diabetes (22.94%), obesity (11.01%), and

cirrhosis (49.54%) were common, with no significant differences between groups. Notably, Hepatitis B was more frequent in survivors (15.79% vs. 3.03%; $p = 0.050$). Other comorbidities, including heart failure and dyslipidemia, were only present in patients who did not survive (Table 2).

In the analysis of clinical outcomes and post-transplant complications, the median total ischemia time was 393 min (321-515), with cold ischemia at 343 min (280-440) and warm ischemia at 60 min (50-67), showing no significant differences between groups. All transplants were from deceased donors, and 8.26% of patients required re-transplantation. A significantly higher occurrence of rejection was observed in deceased patients (30.30%) compared to survivors (13.15%) ($p = 0.034$). Other post-transplant complications were significantly more common in deceased patients (45.45%) compared to survivors (17.10%) ($p < 0.001$). Reintervention percentages due to bleeding (11.92%), post-operative bleeding at 48 h (11.01%), and re-intervention within the 1st month (27.78%) were higher in deceased patients (33.33%), but without significant differences between groups (Table 3). Similarly, according to the Clavien-Dindo, the majority were classified as grade IIIb (52.14%), followed by IIIa (25.64%) and IVa (11.11%). Grade II complications accounted for 5.98%, while the most severe events – grades IVb and V – each represented 2.56% of the total (Table 4).

During the 2011-2014 period, the 1-year survival percent was 63.3% (95% confidence interval [CI]: 43.6-77.7%), with a progressive decline in the first 6 months post-transplant. For the 2015-2018 period, the 1-year survival percentage improved to 84.2% (95% CI: 58.6-94.6%). In the 2019-2022 period, the 1-year survival percentage was 74.1% (95% CI: 48.5-88.3%), showing a slight decrease compared to the 2015-2018 period. Finally, during the 2023-2025 period, the 1-year survival percentage reached 89.1% (95% CI: 73.4-95.7%) ($p = 0.093$) (Fig. 1).

Kaplan-Meier survival analysis assessed the cumulative incidence of graft rejection across four periods: 2011-2014, 2015-2018, 2019-2022, and 2023-2025, with a 365-day follow-up. The probability of remaining rejection-free at 1 year was 70.4% (95% CI: 49.4-84.0) in the 2011-2014 cohort ($n = 29$), with a decline within the first 100 days (100-74.3%). In 2015-2018 ($n = 19$), rejection-free survival improved to 1 year at 78.6% (95% CI: 52.5-91.4), stabilizing after the first 100 days. The 2019-2022 cohort ($n = 20$) exhibited a further increase to 88.2% (95% CI: 59.8-96.9), with fewer early rejection events. Notably, in 2023-2025 ($n = 40$),

Table 2. Personal and family history of patients with liver transplant

Variable	Alive (n = 76) (%)	Deceased (n = 33) (%)	Total (n = 109) (%)	p
Personal history				
Alcohol consumption				
Daily consumption	18 (23.68)	10 (30.30)	28 (25.69)	0.807
Occasional consumption	7 (9.21)	4 (12.12)	11 (10.09)	
Weekly consumption	11 (14.47)	4 (12.12)	15 (13.76)	0.228
Does not consume	40 (52.63)	15 (45.45)	55 (50.46)	
Smoking history				
Former smoker	13 (17.11)	8 (24.24)	21 (19.27)	0.484
Never smoked	61 (80.26)	25 (75.76)	86 (78.89)	
Currently smokes	2 (2.63)	0	2 (1.83)	0.231
Diabetes	15 (19.74)	10 (30.30)	25 (22.94)	
Dyslipidemia	4 (5.26)	0	4 (3.67)	0.275
Obesity	7 (9.21)	5 (15.15)	12 (11.01)	
Hepatitis A	1 (1.32)	0 (0.00)	1 (0.92)	0.697
Hepatitis B	12 (15.79)	1 (3.03)	13 (11.93)	
Hepatitis C	4 (5.26)	1 (3.03)	5 (4.59)	0.521
Unspecified hepatitis	9 (11.84)	4 (12.12)	13 (11.93)	
Heart failure	2 (2.63)	0	2 (1.83)	0.484
Myocardial infarction	1 (1.32)	1 (3.03)	2 (1.83)	
Cirrhosis	39 (51.32)	15 (45.45)	54 (49.54)	0.574
Family history				
Liver cancer	1 (1.32)	0	1 (0.92)	0.697
Pancreatic cancer	1 (1.32)	0	1 (0.92)	0.697
Other cancer	1 (1.32)	0	1 (0.92)	0.697

Daily consumption: more than 7 drinks per week; Occasional consumption (less than once a week); Weekly consumption (1-7 drinks per week); Former smoker (quit more than 6 months ago).

rejection-free survival reached 100%, with no recorded rejection events, suggesting significant advancements in graft survival outcomes, $p = 0.009$ (Fig. 2).

Biliary and vascular complication-free survival varied across periods. For biliary complications, survival declined to 70.0% at 365 days (95% CI: 48.7-83.8%) in 2011-2014, 67.1% (95% CI: 40.9-83.7%) in 2015-2018, 70.3% (95% CI: 41.9-86.7%) in 2019-2022, and improved to 91.4% (95% CI: 75.7-97.2%) in 2023-2025 ($p = 0.076$) (Fig. 3A). Vascular complication-free survival followed a similar pattern: 89.4% at 31 days (95% CI: 70.6-96.5%) in 2011-2014, 89.16% (95% CI: 63.2-97.2%) in 2015-2018, 90.0% (95% CI: 65.6-97.4%) in 2019-2022, and 92.4% (95% CI: 78.1-97.5%) in 2023-2025, $p = 0.975$ (Fig. 3B).

The cold ischemia time (in blue) shows a gradual reduction over the periods, with medians of 461.50 min in 2011-2014, 398.00 min in 2015-2018, 395.50 min in 2019-2022, and 319.00 min in 2023-2025, indicating a decreasing trend in this parameter ($p < 0.001$). On the other hand, the warm ischemia time (in brown) remains relatively stable across the different periods, with

Table 3. Clinical outcomes and post-transplant complications in liver transplant patients

Variable	Alive (n = 76) (%)	Deceased (n = 33) (%)	Total (n = 109) (%)	p
Ischemia times (min)	385 (320-504.5)	440 (330-515)	393 (321-515)	0.856
Cold*	330.5 (271-435)	405 (307.5-456)	343 (280-440)	0.166
Warm*	60 (50-67)	59 (52.5-65)	60 (50-67)	0.856
Transplant outcomes				
Re-transplantation	6 (7.89)	3 (9.09)	9 (8.26)	0.550
Transplant classification				
Adult	72 (94.74)	31 (93.94)	103 (94.50)	0.591
Pediatric	4 (5.26)	2 (6.06)	6 (5.50)	
Donor type				
Living	0	0	0	
Deceased	76 (100)	33 (100)	109 (100)	0.034
Rejection occurrence	10 (13.15)	10 (30.30)	20 (18.35)	
Rejection type				
Acute	5 (6.57)	7 (21.21)	12 (11.01)	0.325
Chronic	5 (6.57)	3 (9.09)	8 (7.34)	
Rejection cause				
Treatment adherence	2 (2.63)	0	2 (1.83)	0.306
Administrative (insurance- related)	1 (1.31)	0	1 (0.92)	
Subtherapeutic drug levels	1 (1.31)	0	1 (0.92)	
Other	1 (1.31)	2 (6.06)	3 (2.75)	
Biliary complications				
Stricture	9 (11.84)	9 (27.27)	18 (16.51)	0.645
Bile leaks	0 (0.00)	2 (6.06)	2 (1.83)	
Other	3 (3.94)	2 (6.06)	5 (4.59)	
Other complications	13 (17.10)	15 (45.45)	28 (25.69)	< 0.001
Post-transplant re interventions				
Reintervention due to bleeding	8 (10.52)	5 (15.15)	13 (11.92)	0.528
Reintervention due to bleeding at 48 h	8 (10.52)	4 (12.12)	12 (11.01)	0.521
Reintervention within the 1 st month	19 (25.00)	11 (33.33)	30 (27.78)	0.321

*Median. IQR: interquartile range.

Table 4. Complications according to the Clavien-Dindo classification

Classification	Number of complications	%
II	7	5.98
IIIa	30	25.64
IIIb	61	52.14
IVa	13	11.11
IVb	3	2.56
V	3	2.56
Total	117	100

values close to 35 min, suggesting that there have been no significant changes in this variable ($p = 0.028$). Regarding the total ischemia time (in green), a progressive reduction is observed, decreasing from 605.00 min

in 2011-2014 to 579.00 min in 2023-2025 ($p = 0.038$) (Fig. 4).

In the logistic regression analysis, the presence of other complications was identified as an independent predictor of mortality in patients undergoing liver transplantation (Odds ratio = 4.28, $p = 0.002$) (95% CI: 1.71-10.68). Besides, patients who experienced post-transplant complications were 4.28 times more likely to die compared to those without complications. The overall model was statistically significant (Likelihood-ratio $X^2 [1] = 9.83$, $p = 0.0017$), although it explained a relatively small proportion of the variance in mortality (Pseudo $R^2 = 0.0749$) $p < 0.001$.

Discussion

The evolution of liver transplant outcomes reflects a significant improvement in the transplant team's




Figure 1. Kaplan-Meier survival curves for mortality within 1 year after liver transplant. Cumulative survival probability within 1-year post-liver transplantation, comparing different time periods. The number of patients at risk is shown at time intervals.




Figure 2. Kaplan-Meier Survival curves for graft rejection within 1 year after liver transplant. Cumulative probability of freedom from graft rejection within 1-year post-liver transplantation, comparing different time periods. The number of patients at risk is presented at time intervals.

experience, leading to better clinical results and increased survival probabilities. In our study, the median age was 7.5 years higher in patients who died compared to those who survived, with a statistically significant difference, similar to findings reported by Su et al.¹⁹ One-year survival consistently improved, rising from 63.3% in 2011-2014 to 89.1% in 2023-2025. Comparable results were observed in a meta-analysis by Barbetta et al., showing the same trend over time.²⁰ This increase indicates advancements in patient selection, optimization of surgical procedures, and greater efficacy in post-operative management. Likewise, Colombian transplant centers have reported improved perioperative protocols and early detection of complications, contributing to better outcomes.²¹

The findings from other Latin American transplant centers further support the regional advancement of transplantation practices. For instance, a recent publication by Varón-Vega et al. presents the experience of a high-altitude lung transplant center in Bogotá, Colombia, offering valuable insights into the development and consolidation of complex transplant programs in resource-constrained settings across Latin America.²² These institutional reports highlight the importance of local expertise, multidisciplinary care, and adaptability in improving outcomes and building sustainable transplant programs in the region.

A notable finding in this study is the progressive reduction in cold ischemia time, with the median duration decreasing from 461.5 min in the 2011-2014 period to 319.0 min in 2023-2025. This trend reflects advancements in organ preservation protocols and logistical efficiencies in transplantation. In contrast, warm ischemia time remained relatively stable, and its variation did not show a significant association with patient mortality. This result is consistent with previous reports, suggesting that while minimizing cold ischemia time is essential for graft viability and long-term outcomes, variations in warm ischemia time within clinically acceptable limits may not have a substantial impact on post-transplant survival.^{23,24}

An improvement in rejection prevention was evident, with a lower incidence observed among surviving patients compared to those who did not survive during the studied periods. The absence of rejection episodes within the 1st year post-transplant, observed in the most recent evaluation period, reflects advancements in immunosuppressive management and post-transplant care. Optimized immunosuppressive protocols, tailored treatment strategies, and strict adherence to follow-up regimens have likely contributed to these findings. Research in Latin America has demonstrated that personalized immunosuppressive approaches play a crucial role in reducing rejection rates and prolonging graft survival. These results emphasize the need for ongoing refinement of immunosuppressive therapies and patient monitoring to enhance long-term transplant success.^{25,26}

Besides, patients who experienced post-transplant complications were 4.28 times more likely to die compared to those without complications, highlighting the need to further strengthen strategies for the prevention and management of these complications.²⁷ This finding underscores the importance of close monitoring during the 1st post-operative month, particularly given that vascular and biliary




Figure 3. A and B: Kaplan-Meier survival curves for biliary and vascular complication. Kaplan-Meier survival analysis for the absence of complications at 1 year (biliary) and 1 month (vascular) following liver transplantation.




Figure 4. Cold, warm, and total ischemia times in liver transplantation. The boxplots show the evolution of cold, warm, and total ischemia times over four distinct periods; they show a general trend of improvement in ischemia times, with notable consistency in warm ischemia time.

complications primarily occurred during this period.⁷ Period-based analysis also reveals a decrease in the incidence of rejection and an improvement in survival, which may be attributed to the accumulated clinical experience of the treating team, the adoption of new technologies, and enhancements in perioperative management protocols.²⁸ In Colombia, hospitals implementing multidisciplinary post-transplant follow-up programs have reported a significant reduction in early post-operative complications and better long-term patient outcomes.²⁹

It is also important to consider the potential influence of the COVID-19 pandemic on post-transplant outcomes, particularly in the earlier years of the 2020s.

Several studies have reported that patients with underlying liver disease, including cirrhosis or metabolic-associated fatty liver disease, faced increased risks of severe COVID-19 and mortality, which may have indirectly affected transplant survival rates during that period. In addition, drug-induced liver injury and systemic inflammatory responses related to SARS-CoV-2 have been implicated in liver dysfunction, complicating the clinical course of transplant recipients. The heightened vulnerability of these patients to infections, immune dysregulation, and liver injury during the pandemic years could have contributed to increased post-transplant morbidity and mortality, warranting further investigation in future studies.³⁰

These findings reaffirm that the accumulated experience over the years has been a key factor in improving liver transplant outcomes.³¹ It is recommended to continue optimizing ischemia times, strengthening measures to prevent acute rejection, and refining strategies to reduce post-operative complications. Future research should focus on identifying prognostic factors that enhance the accuracy of rejection and post-operative complication risk prediction, as well as evaluating the impact of novel immunosuppressive therapies and surgical techniques on long-term outcomes. In addition, comparative studies between different transplant centers in Latin America could provide valuable insights into best practices and opportunities for standardizing care protocols across the region.³²

This study has some limitations inherent to its retrospective design, as well as the fact that the data come from a single institution, which may restrict the generalizability of the findings to other settings. Nevertheless, it has several strengths. It provides a detailed analysis of long-term outcomes in liver transplant patients, supported by a rigorous follow-up protocol and a multidisciplinary approach.

Conclusions

The results of this study demonstrate a progressive improvement in 1-year survival, along with a reduction in rejection incidence, suggesting advances in post-transplant management. However, post-transplant complications, excluding biliary and vascular complications, remain a significant risk factor for mortality. Finally, the progressive reduction in cold and total ischemia times reflects an optimization of surgical and logistical procedures in liver transplantation.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of human subjects and animals. The authors declare that the procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation and with

the World Medical Association and the Declaration of Helsinki. The procedures were authorized by the Institutional Ethics Committee.

Confidentiality, informed consent, and ethical approval. The study was approved by the Research Ethics Committee (REC) of the Fundación Cardiovascular de Colombia under the approval code CEI-2024-016-8. All patient data were anonymized to ensure confidentiality, and the study was conducted in accordance with the principles outlined in the Declaration of Helsinki. This article complies with the provisions set forth in Resolution 008430 of 1993 by the Ministry of Health of Colombia, which establishes guidelines for the protection of the rights of research subjects in scientific studies. Informed consent was obtained from all participants before their inclusion in the study, and no identifiable information was disclosed in this publication. The datasets generated and analyzed during the study are available on reasonable request from the corresponding author.

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing or creation of the content of this manuscript.

References

1. Cristin DJ, Forman LM, Jackson WE. Beyond survival: targeting health-related quality of life outcomes after liver transplantation. *Clin Liver Dis (Hoboken)*. 2021;17:359-64.
2. Observatorio Global de Donación y Trasplante (GODT), Organización Nacional de Trasplantes (ONT). 2023 Global Report on Donation and Transplantation. Madrid, España: GODT-ONT; 2024.
3. Clemente Ricote G, Díaz Sánchez A. Complicaciones precoces tras el trasplante hepático. ¿Podemos llegar a predecirlas? *Rev Esp Enferm Dig*. 2008;100:121-8.
4. Fasullo M, Patel M, Khanna L, Shah T. Post-transplant biliary complications: advances in pathophysiology, diagnosis, and treatment. *BMJ Open Gastroenterol*. 2022;9:e000778.
5. Zuluaga Arbeláez N, Morales Ortiz A, Calle Tavera L, Londoño Castillo J, Escobar Pérez R, Licona Vera E. Complicaciones biliares postrasplante hepático: fugas y estenosis/gastric cancer - update on diagnosis and treatment: literature review. *Gastroenterol Latinoam*. 2024;35:160-5.
6. Restrepo J. Historia de los trasplantes de órganos en Colombia. *Rev Colomb Cir*. 1990;5 Suppl 3:60-1.
7. Tarazona-León C, Vanegas-Ballesteros M, Vergara FA, Camelo-Pardo G, Manrique-Hernández EF, Oliveros-Araujo KV. Complicaciones vasculares postrasplante hepático, una situación con alta morbimortalidad. *Rev Colomb Cir*. 2024;39:738-44.
8. Instituto Nacional de Salud (Colombia). Marco Legal de la Red de Donación y Trasplante en Colombia. Bogotá: Instituto Nacional de Salud; 2022.
9. Sierra Mendoza R, López Noguera JS, Sosa Bermúdez NE, Ruvalcaba Ledezma JC, Hernández Ceruelos MC. Donación de órganos desde una perspectiva del personal médico. *J Negat No Posit Results*. 2021;6:307-20.
10. González-Domínguez E, Córdova D, Abad P, González E, González C, Cordero JJ, et al. Protocolo de inmunosupresión en trasplante hepático: revisión de la literatura. *Rev Hepatol*. 2022;3:40-56.
11. McElroy L, Daud A, Davis A, Lapin B, Baker T, Abecassis M, et al. A meta-analysis of complications following deceased donor liver transplant. *Am J Surg*. 2014;208:605-18.
12. Tarazona-León C, Vanegas Ballesteros M, Vergara FA, Camelo Pardo G, Manrique Hernández EF. Complicaciones biliares postrasplante hepático en una institución de Colombia. *Rev Cir*. 2024;76:563-9.

13. Dacha S, Barad A, Martin J, Levitsky J. Association of hepatic artery stenosis and biliary strictures in liver transplant recipients. *Liver Transpl.* 2011;17:849-54.
14. Mazzaferro V, Regalia E, Doci R, Andreola S, Pulvirenti A, Bozzetti F, et al. Liver transplantation for the treatment of small hepatocellular carcinomas in patients with cirrhosis. *N Engl J Med.* 1996;334:693-9.
15. Toniutto P, Fumolo E, Fornasiere E, Bitetto D. Liver transplantation in patients with hepatocellular carcinoma beyond the Milan criteria: a comprehensive review. *J Clin Med.* 2021;10:3932.
16. Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg.* 2004;240:205-13.
17. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform.* 2009;42:377-81.
18. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. The REDCap consortium: building an international community of software platform partners. *J Biomed Inform.* 2019;95:103208.
19. Su F, Yu L, Berry K, Liou IW, Landis CS, Rayhill SC, et al. Aging of liver transplant registrants and recipients: trends and impact on waitlist outcomes, post-transplantation outcomes, and transplant-related survival benefit. *Gastroenterology.* 2016;150:441-53.e6; quiz e16.
20. Barbeta A, Aljehani M, Kim M, Tien C, Ahearn A, Schilperoort H, et al. Meta-analysis and meta-regression of outcomes for adult living donor liver transplantation versus deceased donor liver transplantation. *Am J Transplant.* 2022;22:2790-803.
21. Fernández Aguilar JL, Suárez-Muñoz MA, Santoyo Santoyo J, Sánchez Pérez B, Pérez Daga A, Ramírez Plaza CP, et al. Aggressive management of the arterial complications of liver transplantation. Impact upon survival and biliary complications. *Cir Esp.* 2010;87:155-8.
22. Varón-Vega FA, Rincón A, Tuta-Quintero E, González-Avilés C, Prada LP, Téllez LJ. Survival analysis in a high-altitude lung transplant program: insights from a real-life observational study. *Cir Cir.* 2025;93:267-72.
23. Alfageme Zubillaga M, Royuela Vicente A, Cortés León C, Minaya Bernedo J, Hernández Guilbert PM. Trasplante hepático: influencia del tiempo de isquemia y la edad del donante en la rigidez del injerto medida mediante elastografía tipo ARFI. *Seram.* 2018;2(1).
24. Raimondi G, Wood KJ, Perelson AS, Arciero JC. Editorial: transplant rejection and tolerance-advancing the field through integration of computational and experimental investigation. *Front Immunol.* 2017;8:616.
25. Mishra S, Taneja S. Algorithmic approach to deranged liver functions after transplantation. *J Clin Exp Hepatol.* 2024;14:101317.
26. Briceño Delgado J. Implementation of multimodal enhanced recovery protocols in liver transplantation: is it possible in Spain? *Cir Esp.* 2024;102:67-8.
27. Guerra JF, Quezada JL, Cancino A, Arrese M, Wolff R, Benítez C, et al. Trasplante hepático: evolución, curva de aprendizaje y resultados después de los primeros 300 casos. *Rev Med Chil.* 2019;147:955-64.
28. Salazar-Flórez JE, Arango-Flórez EM, Morales-Toro M, Rojas-Copete AM, Leal-Castillo DA, Toro-Rendón LG. Síndrome metabólico después de trasplante hepático en pacientes del Centro Especializado San Vicente Fundación, Rionegro, Antioquia, Colombia, 2013-2017. *Rev Peru Med Exp Salud Publica.* 2022;39:474-9.
29. Burbano Pinchao JF. Complicaciones Presentadas en el Postoperatorio Temprano del Trasplante Hepático en el Hospital Universitario Fundación Santa Fe de Bogotá [Tesis de Grado]. Bogotá: Universidad del Rosario; 2017.
30. Abdo-Francis JM, Moreno-Alcantar R, Pérez-Hernández JL, Remes-Troche JM, Velarde-Ruiz Velasco A, Cerda-Reyes E, et al. Impact of COVID-19 on pre-existing liver disease. *Cir Cir.* 2024;92:131-6.
31. Muñoz Maya OG, Insuasty Enríquez MR, Marín JI, Hoyos Duque SI, Guzmán Luna CE, Mena Hurtado Á, et al. Biliary duct complications after orthotopic liver transplants: experience from the pablo tobón uribe hospital - Medellín. *Rev Col Gastroenterol.* 2011;26:8-17.
32. Santos-Sánchez OM, Muñoz Maya OG, Marín-Zuluaga JI, Restrepo Gutiérrez JC. Trasplante hepático en un centro de Colombia: comparación de dos periodos. *Rev Col Gastroenterol.* 2019;34:117-24.

Identifying the best cutoff value of the fecal occult blood immunochemical test in the detection of advanced and neoplastic colorectal lesions

Identificación del mejor punto de corte de la prueba inmunoquímica de sangre oculta en heces en la detección de lesiones colorrectales avanzadas y neoplásicas

Patricia Gaggero^{1*}, Eduardo Fenocchi¹, Cecilia Silva¹, Juan C. López-Alvarenga², Natalia Lambert¹, Fabián Batista¹, Óscar Teramoto-Matsubara³, Sergio Sobrino-Cossío⁴, Jonathan R. White⁵, and Adolfo Parra-Blanco⁵

¹Digestive Cancer Centre, National Cancer Institute, Montevideo, Uruguay; ²Population Health & Biostatistics, School of Medicine, University of Texas, Rio Grande Valley, Edinburg, Texas, USA; ³Endoscopy Service, Hospital ABC, Mexico City, Mexico; ⁴Endoscopy Service, Hospital Angeles del Pedregal, Mexico City, Mexico; ⁵NIHR Nottingham Biomedical Research Centre, Department of Gastroenterology, Nottingham University Hospitals NHS Trust and University of Nottingham, Nottingham, United Kingdom

Abstract

Objective: Screening is an effective tool for detecting colorectal lesions in asymptomatic subjects. There is a positive correlation between fecal immunochemical test (FIT) values and the size of tumors. Despite the efficacy of screening, the detection of colorectal cancer (CRC) remains low. The primary objective was to evaluate the best FIT cutoff value for detecting advanced adenomas and CRC among individuals at average risk in a country with a high incidence and morbidity from CRC. **Methods:** This observational and prospective study analyzed consecutive cases in 1461 asymptomatic subjects with a positive FIT (≥ 100 ng hemoglobin [Hb]/mL) referred for colonoscopy (2012-2015) at a tertiary center in Uruguay. **Results:** Colorectal lesions were detected in 35.3% (516/1461) of cases, with a mean age of 62.8 ± 8.3 years. About 53.2% were men and 65.1% of the tumors were located in the left side of the colon. The size of the lesion and FIT values ($p = 0.001$) were positively correlated. Laterally spreading tumors predominated in the right colon (586 ng Hb/mL; 95% Confidence interval [CI] 443.4-760). One hundred and thirty-five (26%) lesions were advanced adenomas (15 ± 6.7 mm); 694.6 ng/mL; 95% CI 632.4-756.9). The highest diagnostic yield (0.5112) for advanced adenomas was at a FIT level of 400 ng Hb/mL (accuracy 88.6%). There were significant differences in FIT values early and advanced CRC (927 ng/mL; [95% CI: 637-1082] vs. 1453 [95% CI: 1352-1594; $p = 0.001$]). **Conclusions:** A FIT value of 400 ng Hb/mL was the best diagnostic yield to detect advanced adenomas and CRC. This value is useful during the COVID-19 pandemic as it allows prioritization of colonoscopy to those most at risk of significant disease, thus reducing risks to both patients and healthcare workers.

Keywords: FIT. Fecal occult blood immunochemical test. Advanced adenomas. Colorectal cancer.

Resumen

Objetivo: El cribado es una herramienta eficaz para detectar lesiones colorrectales en sujetos asintomáticos. Existe una correlación positiva entre los valores de la prueba inmunoquímica fecal (FIT) y el tamaño de los tumores. A pesar de la eficacia del cribado, la detección del cáncer colorrectal (CCR) sigue siendo baja. El objetivo principal fue evaluar el mejor valor de corte FIT para detectar adenomas avanzados y CCR entre personas con riesgo promedio en un país con alta incidencia y morbilidad por CCR. **Métodos:** Este estudio observacional y prospectivo analizó 1461 sujetos asintomáticos con FIT positivo (≥ 100 ng de hemoglobina [Hb]/mL) remitidos para colonoscopia (2012-2015) en un centro terciario en Uruguay. **Resultados:** Se detec-

*Correspondence:

Patricia Gaggero
E-mail: patriciagaggero07@gmail.com

Date of reception: 29-12-2023

Date of acceptance: 12-01-2024

DOI: 10.24875/CIRUE.M24000952

Cir Cir (Eng). 2026;94(3):326-334

Contents available at PubMed

www.cirugiaycirujanos.com

2444-0507/© 2024 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

taron lesiones colorrectales en el 35.3% (516/1461) de los casos, con una edad media de 62.8 ± 8.3 años. Alrededor del 53.2% eran hombres y el 65.1% de los tumores estaban ubicados en el lado izquierdo del colon. El tamaño de la lesión y los valores FIT ($p = 0.001$) se correlacionaron positivamente. Los tumores con extensión lateral predominaron en el colon derecho (586 ng Hb/mL; intervalo de confianza [IC] del 95%: 443.4-760). Ciento treinta y cinco (26%) lesiones fueron adenomas avanzados (15 ± 6.7 mm); 694.6 ng/ml; IC 95% 632.4-756.9). El rendimiento diagnóstico más alto (0.5112) para los adenomas avanzados se obtuvo con un nivel FIT de 400 ng Hb/ml (precisión del 88.6%). Hubo diferencias significativas en los valores de FIT de CCR temprano y avanzado (927 ng/mL; [IC 95%: 637-1082] vs. 1453 [IC 95%: 1352-1594; $p = 0.001$]). **Conclusiones:** Un valor FIT de 400 ng Hb/mL fue el mejor rendimiento diagnóstico para detectar adenomas avanzados y CCR. Este valor es útil durante la pandemia de COVID-19, ya que permite priorizar la colonoscopia a aquellos con mayor riesgo de padecer una enfermedad importante, reduciendo así los riesgos tanto para los pacientes como para los trabajadores de la salud.

Palabras clave: Prueba inmunoquímica de sangre oculta en heces. Adenomas avanzados. Cáncer colorrectal.

Introduction

The incidence and mortality of cancer have increased in lower-income countries. This is likely due to epidemiological and demographic transition (aging and population growth) as well as exposure to risk factors such (alcohol and smoking).¹ In 2018, the number of new cases detected was 18.1 million with, 9.6 million cancer-related deaths. The incidence in the United States of America (USA) was 21% with a mortality of 14.4%.²⁻⁴ The organs most commonly involved in Central and South America were as follows: prostate, lung, colon, and rectum colorectal cancer (CRC) and stomach in men, and breast, cervix/uterus, CRC, and lung in women. In 2018, the overall incidence and mortality reported were 10.2% and 9.2%, respectively, for both men and women.²

Cancer incidence has increased in Central/South America and Eastern Europe but is decreasing in developed countries such as the USA, Canada, Australia, New Zealand, and the rest of Europe).² The incidence rates among men and women in Latin America were 10.3 and 16.8, respectively. Incidence in the countries with the highest rates was as follows: Uruguay (34.2 men and 24.7 women), Brazil (27.7 and 21.5), and Argentina (25.2 in men).

Mortality is extremely high in Central (8-9 fold) and South America (3-5 fold) when compared to Europe and USA⁵. Although it has generally been reported as being < 10 , mortality is high in Uruguay (17.7 among men and 12.0 among women), Cuba (11.3 women), and Argentina (14.6 men).^{6,7}

Uruguay and Argentina have implemented national programs for the early detection of CRC.⁷⁻¹⁰ Detailed knowledge of the natural history of cancer development (the adenoma-carcinoma sequence) allows for

early detection at the subclinical stage and appropriate therapeutic intervention.¹¹

In sporadic CRC, the tiered non-linear accumulation of genetic and chromosomal alterations occurs in approximately 7-10 years;^{11,12} but this time period is shorter for hereditary CRC.

CRC is a common public health problem (5% incidence). Screening is effective when the appropriate population is targeted. Endoscopic removal of adenomas reduces the incidence of CRC (76%) and its mortality (53%).¹³⁻¹⁵ Although colonoscopy is the most commonly used method for detecting adenomas and CRC, it is not the most cost-effective tool.^{16,17} Screening programs have been implemented in the WHO population programs.^{18,19}

The fecal occult blood (FOB) or fecal immunochemical test (FIT) is currently the best screening method²⁰. FIT accurately detects globin component of hemoglobin (Hb) to the nearest nanogram (ng) per milliliter (mL) of feces using immunoassay. Colonoscopy is indicated when the patients' FIT values are ≥ 100 ng Hb/mL, due to the high sensitivity of detecting disease.^{16,21,22}

For adequate population screening, FIT positivity rates should range around 3-6%.²¹⁻²⁵ In the case of CRC, sensitivity (Sn), specificity (Sp), positive predictive values, and negative predictive values (NPV) of FIT were reported as 100%, 90%, 16%, and 100%, respectively; and 74%, 93%, 45%, and 98%, respectively, for all neoplasms. FIT detects virtually all CRCs and 74% of significant colorectal lesions.²⁶

Screening should be started at the age of 50 and be suspended either at 75 years of age or when the patient is expected to live < 10 years. Between 75 and 85 years, the risk to benefit ratio should be individualized. There is no benefit beyond the age of 85.¹¹ In the US, screening is opportunistic, unlike other

countries where it is more systematic and organized to cover more of the population.

The primary objective was to evaluate the best FIT cutoff value for detecting advanced adenomas and malignancies among individuals at average risk in a country with a high incidence and morbidity from CRC.

Methods

Consecutive patients with an average risk of CRC and a positive FIT (> 100 ng Hb/mL) scheduled for a colonoscopy at the Centre for Digestive Cancer of the National Cancer Institute of Montevideo, Uruguay, were recruited. All individuals provided written consent and this study followed the standards for reporting diagnostic accuracy. This cross-sectional, prospective, observational, and single-center study was carried out between 2012 and 2015. The study is part of the systematic program for the prevention and early detection of CRC under the Ministry of Health of Uruguay.

Selection criteria

- Inclusion: asymptomatic individuals, over 50 years of age, positive FIT (> 100 ngHb/mL), and informed written consent.
- Exclusion: personal or family history of CRC, digestive bleeding, colorectal resective surgery, inflammatory bowel disease and/or red flag signs (weight loss, anemia, and abdominal masses), and incomplete examination due to poor bowel cleansing.

Mass media campaign (television, radio, and press) were used to invite the population to participate in the national CRC screening program. The process for collecting samples, storing at -10°C , and returning it within 5 days was explained to each patient. There were no restrictions on diet or medication.

Patients with positive FIT results were scheduled for a colonoscopy within 8 weeks, which was performed after informed consent and after bowel preparation. The quality of FIT measurements was ensured through daily calibration and external validation using samples supplied by the manufacturer every 6 months. The colonoscopies were performed by three experienced endoscopists with a minimal total experience of 3000 colonoscopies. The cold technique was typically used to remove lesions < 5 mm and endoscopic mucosal resection to remove larger ones.

Clinical and demographic data were recorded, including age, sex, FIT values (ng Hb/mL), presence

of colorectal lesions (type – Paris classification –, size and location; in addition to laterally spreading tumors (LST) > 10 mm), and histological types (subtypes and degree of dysplasia/differentiation).

The adenomas were classified as advanced or not advanced. Early CRC did not extend beyond the submucosa (Tis-T1); for invasive, CRC malignant cells were observed in the *muscularis propria* or beyond. The size of the polyps was defined *ex vivo* after resection and before fixation with 10% formaldehyde.

Definition of variables

- Subjects at average risk of CRC: Asymptomatic adult individuals over 50 years of age, with no personal/family history of CRC and/or adenomas or inflammatory bowel disease⁷.
- Advanced adenomas: Adenomas ≥ 10 mm and/or with villous component $\geq 25\%$ and/or high-grade dysplasia. Non-advanced adenoma: 1-2 adenomas each < 10 mm in size.⁷
- (FIT-OC-Eiken [Eiken Chemical Co., Eiken, Japan]) was used during this study. The reported sensitivity and specificity was 96-98% and 91-95%, respectively.^{27,28} The detection range was between 0 and 2000 ng Hb/mL of buffer, which enabled the cutoff point to be set for a positive result. The cutoff point was 100 ng Hb/mL as recommended by the manufacturers.¹⁴

Statistical analysis


The data were captured in a Lotus Approach v. 9.7 Windows database and included age, sex, topography, size, histopathological diagnosis, and FIT quantification. To evaluate the differences between independent samples, Chi-squared tests and the exact Fisher's test were used for categorical variables and Student's t-test was applied for quantitative variables, with a significance level of 0.05. One-way analyses of variance and covariance were used to adjust the contrast between lesions in terms of age, sex, and positive findings of colorectal lesions. A hyperbolic function was calculated for two parameters.

The Sn, Sp, PPV, NPV, and accuracy of different FIT cutoff points were calculated, and the ROC curve plotted ($y = 0.5 \ln(x) + 1$ Area Under the Curve $\times 0.5112$) for the best FIT cutoff point in the detection of advanced adenoma. SPSS v. 17.0 and Epi Info v. 3.3.2.0 as statistical software were used.

Table 1. Type of colorectal lesions detected through colonoscopy in asymptomatic individuals

Paris classification	Type of colorectal polyps detected in 516 colonoscopies				p
	A Pedunculated Polyp (Ip)	B Semi-pedunculated polyp (Isp)	C Sessile Polyp (Is)	D Lateral spreading tumor	
Number of lesions	142	114	248	12	-
Age (years)					
Average	62.1	62.8	62.4	65.5	0.5
SD	88	8.2	8.1	8.3	
95% CI	60.6-63.6	61.1-64.5	61.3-63.5	60.8-70.3	
Size (mm)					0.001
Average	12.9	8.6	5.6	14.8	
SD	6.2	5.4	4.3	9.5	
IC95%	11.9-13.1	7.4-9.7	5.0-6.3	12.1-17.5	
FIT (ng Hb/mL)					0.001
Average	631.3 ^a	452.1 ^b	401.4 ^c	779.2 ^d	
SD	484	407	368	525	
IC95%	537.6-705	366.2-558	348.7-454.5	544-1614	

The letters (a, b, c or d) show the statistically significant differences between the groups, where the letter a differs from the other groups (b, c or d). Example: a = a, but a ≠ b or c or d. FIT: fecal occult blood immunochemical test; SD: standard deviation; 95% CI: confidence interval 95%.

**Figure 1.** Differences between FIT levels and the type of colorectal lesion.

Results

The study included 1461 colonoscopies of subjects with average cancer risk and with FIT values ≥ 100 ng. Complete FIT and colonoscopy outcomes were available for 1461 patients who met the inclusion criteria for the final analysis.

No colorectal lesions (advanced adenomas/adenomas/early CRC or advanced CRC) were detected in 945 cases (64.6%). The average age of those subjects

was 61 ± 9 years (55% men) and their average FIT was 317 ± 180 (95% CI 295.9-338.5). No visible lesions were found in 485 (237 ± 140 ng; 95% CI 21.3-270.7). Median FITs were 237, 370, 382, and 188 for diverticulitis ($n = 396$), angiodysplasias ($n = 36$), non-specific rectocolitis ($n = 10$), and colonic ulcers ($n = 7$), respectively. Other findings included three cases of pinworm (*Enterobius vermicularis*), two submucosal lesions, and two cases of chronic idiopathic ulcerative colitis.

Colorectal lesions were detected in 516 cases (35.3%). About 25.7% of the patients presented with > 2 lesions. For the purpose of FIT value analysis, the largest lesion was used. The average age was 62.8 ± 8.3 years and 53.2% were male. Majority of lesions were located in the left colon (336, 65.1%) and with 180 (34.9%) in the right colon. The morphological Paris classification of lesions is described in table 1. There was a statistical difference between lesion types and FIT levels ($p = 0.0001$) and also size ($p = 0.001$). Figure 1 demonstrates differences between FIT levels and lesion type.

The most common lesions confirmed by histology were tubular adenoma ($n = 167$), tubulovillous adenoma ($n = 104$), hyperplastic ($n = 63$), advanced CRC ($n = 54$) villous adenoma ($n = 38$), early CRC ($n = 17$), and serrated adenomas.¹² Lesion size and FIT levels were positively correlated ($p = 0.001$), see table 2. Patient's age did not correlate with FIT levels. There was no significant difference between FIT value and LST or early CRC size.

Table 2. Colorectal lesions and FIT levels

Lesions	FIT Levels (ng Hb/mL)								p
	100-200	201-300	301-400	401-500	501-600	601-700	701-800	> 801	
Number	169	62	45	24	18	21	28	149	
Average size (mm) (SD)	6.37 (3.3)	7.1 (4.1)	7.5 (5.3)	10.8 (8.5)	12.6 (7.8)	12.6 (7.9)	14.2 (8)	22 (13.1)	0.001
95% CI	4.8-7.8	4.7-9.6	4.7-10.4	6.9-14.7	8-17.1	8.4-16.8	10.6-17.9	20.5-23.6	

FIT: fecal occult blood immunochemical test; SD: standard deviation; 95% CI: confidence interval 95%.

Table 3. Diagnostic accuracy to correctly diagnose colorectal lesions based on the 400 ng/mL FIT cutoff value

Tumor	ROC Curve (AUC)	IC 95%
Adenoma	0.691	0.656, 0.724
Advanced adenoma	0.831	0.798, 0.861
Laterally Spreading Tumors	0.907	0.878, 0.933
Early CRC	0.996	0.985, 0.999
Advanced CRC	1.0	0.993, 1.0
Laterally Spreading Tumors + early CRC	1.0	0.992, 1.0
All the lesions	0.777	0.749, 0.802

AUC: area under the curve; 95% CI: 95% confidence interval; CRC: colorectal cancer.

LST lesions predominated in the right colon ($n = 12$; 66.6%) and early CRC in the left ($n = 17$; 88%). There were no significant differences between size (17.5 vs. 15, $p = 0.1$) and FIT values (average 626 vs. 927 ng/mL, $p = 0.1$).

There were 135 cases of advanced adenomas, with an average size of 15 ± 6.7 mm (95% CI 13.6-16.3) and FIT levels of 694.6 ng (95% CI 632.4-756.9). There was no correlation between advanced adenoma size and FIT levels ($R = 0.22$). The best cutoff value for detecting significant pathology was 400 ng/mL (Sn 0.59, Sp 0.84) with false-negative rates of 0.16. The positive likelihood ratio (LR+) and negative likelihood ratio (LR-) were 3.81 and 0.48, respectively. Table 3 and figure 2 illustrate the diagnostic accuracy for a 400 ng/mL cutoff and ROC curves. Diagnostic accuracy for different FIT levels is demonstrated in table 4.

There were 71 cases of CRC (17 early and 54 advanced) with an average age of 65.8 ± 8 years (1:1 male-female ratio) predominating in the left colon (88% early and 80% advanced). There were

significant differences between FIT values for early and advanced CRC (average 927 ng; [95% CI: 637-1082] vs. 1453 [95% CI: 1352-1594; $p = 0.001$]). The degree of tumor differentiation was well differentiated ($n = 31$), moderately differentiated ($n = 34$), and poorly differentiated ($n = 6$).

11/12 LST were treated endoscopically and one was surgically resected. 14/17 early CRC were treated by endoscopic resection and the remainder required surgery. All advanced CRCs were referred to the oncology team.

Discussion

A FIT cutoff value of 100 ng Hb/mL is highly sensitive for detecting CRC and advanced adenomas. The Sn, Sp, PPV, and NPV to detect CRC were 100%, 90%, 16%, and 100%, respectively, and for advanced neoplasms 74%, 93%, 45%, and 98%, respectively. A FIT detects 100% of CRC and 74% of colorectal lesions.²⁹

Although the frequency of advanced adenomas and malignant lesions was higher in our studied population, there was not a correlation between the size (average 15 ± 6.7 mm) and the FIT level of 694.6 ng ($R = 0.22$). However, the most significant accuracy was seen with a value of 400 ng Hb/mL (Sn 0.59, Sp 0.84) with a 0.16 false-negative rate, a positive likelihood ratio (LR+) and negative likelihood ratio (LR-) of 3.81 and 0.48, respectively.

Nagorni et al.³⁰ reported a higher detection rate of advanced adenoma among individuals with a positive FIT than in symptomatic subjects undergoing colonoscopy (11.6% vs. 5.1%).³⁰ Similarly, our group reported a higher detection rate in colonoscopy in asymptomatic subjects (positive FIT) compared with symptomatic patients without FIT testing (8.9 vs. 4.35%) in 3234 consecutive cases at the Cancer Detection Center in Montevideo, Uruguay.³¹ qFIT (quantitative)




Figure 2. ROC curve for the detection of colorectal lesions using FIT. **A:** simple adenomas, **B:** advanced adenomas, **C:** laterally spreading tumors, **D:** early CRC, **E:** advanced CCR, **F:** laterally spreading tumors plus early CRC, and **G:** all.

showed a directly proportional relationship between the size of the lesion and the amount of bleeding. For each mm increase, the value of Hb in FIT increased by 35 ± 9 ng ($p < 0.001$) with the best diagnostic yield of 15 mm, where the values stabilized and the curve almost flattened between 1100 and 1900 ng/Hb.

The frequency of advanced adenomas/CRC was high (13.7%) in our study when compared to previous studies. Van Rossum et al.²⁹ reported a CRC detection rate of 0.45, 0.44, 0.39, 0.39, 0.39, 0.39, 0.39, and 0.37% using values of 50, 75, 100, 125, 150, 175, 200, and 225 ng Hb/mL, respectively. However, the number of colonoscopies required to detect one case was 15.3 (50 ng Hb/mL), 12.4 (75 ng Hb/mL), 11.7 (100 ng Hb/mL), 10.3 (125 ng Hb/mL), 9.8 (150 ng Hb/mL), 9.0 (175 ng Hb/mL), 8.3 (200 ng Hb/mL), and 8.1 (225 ng Hb/mL) with very wide 95% confidence intervals (95% CI). In this study, the Sp to detect advanced adenoma using a value of 400 ng Hb/mL was over 96% (96–98.7%).

Hernandez et al.³² reported that the FIT cutoff points with the highest yield for advanced neoplasm and

CRC were 272.57 ± 597.28 (ROC Curve 0.97) and $998.00 \pm 1,075.44$ (ROC Curve 0.72), respectively. In our study, the cutoff points with the highest yield (the highest true-positive rate together with the lowest false-positive rate) for predicting advanced adenoma were 395 ng Hb/mL, 927 Hb/mL ng for early CRC, and 1453 ng Hb/mL for advanced CRC.

The assessment of FIT cutoff values and advanced adenomas is often limited in studies due to the number of colonoscopies performed, type of FIT immunoassay used, and the inclusion of both the high-risk screening population and symptomatic patients. These cofounders limit the applicability of findings to screening populations.

Yuan et al. reported that different cutoff values were associated with the performance of qFIT (quantitative). The sensitivity and specificity of qFIT for advanced neoplasia (AN) was 51.3% and 86.4%, respectively, with the best cutoff level of 400 ng/mL. The sensitivity and specificity for CRC were 61.0% and 89.1%, with the best cutoff level of 500 ng/mL.³³

Table 4. Diagnostic accuracy of FIT for lesions at different cutoff values

Diagnosis	200	300	400	500	600	700	800	900	1000
Adenoma									
Sensitivity	58.0	43.0	29.0	21.0	17.0	13.4	9.4	6.4	5.0
Specificity	70.0	91.2	97.0	100	100	100	100	100	100
Accuracy	65.1	71.8	69.7	68.2	66.6	65.2	63.6	62.5	62.0
Advanced adenoma									
Sensitivity	78.6	67.0	59.5	51.5	48.5	46.3	38.2	33.1	29.4
Specificity	70.0	91.2	100	100	100	100	100	100	100
Accuracy	72.0	85.5	88.3	88.6	88.0	87.4	85.5	84.3	83.5
Laterally spreading tumors									
Sensitivity	91.7	88.3	-	66.7	41.7	-	-	-	33.3
Specificity	70.0	91.2	-	97.0	100	-	-	-	100
Accuracy	74.0	91.0	-	96.2	98.5	-	-	-	98.2
Early CRC									
Sensitivity	100	100	95.4	-	81.8	60.0	50	-	31.8
Specificity	70.0	91.2	97.0	-	100	100	100	-	100
Accuracy	71.3	91.6	97	-	99.1	98.0	97.6	-	96.5
Advanced CRC									
Sensitivity	100	100	100	100	94.3	92.4	88.7	79.2	70.770.3
Specificity	70.3	91.2	97.0	100	100	100	100	100	100
Accuracy	73.3	92.1	97.4	100	95.4	99.2	98.8	97.8	97.0
CRC + Lateral Spreading									
Sensitivity	97.0	94.1	94.1	85.2	-	64.7	47.0	35.3	32.3
Specificity	70.7	91.2	92.0	97.0	-	100	100	100	100
Accuracy	72.0	91.4	91.8	96.2	-	97.5	96.2	95.4	95.2
All lesions									
Sensitivity	70.0	58.3	47.8	39.7	-	36.2	33.4	23.0	20.5
Specificity	70.2	91.2	97.0	100	-	100	100	100	100
Accuracy	70.0	73.5	70.0	67.5	65.3	63.5	61.0	48.5	57.1

Shahidi and Cheung³⁴ evaluated FIT performance in an average risk Canadian population; using cutoff values of > 100 ng Hb/mL, this increased the PPV for advanced adenoma by 6.5% and CRC by 1.5% but the main trade-off was the potential for missed lesions. A 50 ng Hb/mL cutoff value has been shown to be most cost-effective value.

These findings could be highly important in the decision-making process during the SARS-COV-2 (COVID-19) pandemic, helping clinicians to risk stratify which invasive aerosol generating procedures are need to be undertaken in timely manner and which can be safely postponed to a later date. During the most difficult times in the COVID-19, pandemic screening programs, elective colonoscopies in patients with positive FITs were postponed and only emergency procedures undertaken.³⁵

Unfortunately, it is still too early to know the impact on the subsequent health, quality of life, and endoscopic lesion detection of the postponed individuals with positive FIT who were scheduled for colonoscopy.

In our study, the detection rate of colorectal lesions was 35% (post-test likelihood) using a non-invasive test (FIT) in subjects at average risk. We observed significant differences in location (left colon), FIT values ($p = 0.0001$), and lesion size ($p = 0.001$), but not for age or gender.

Nationwide policies have implemented CRC screening programs in many countries¹¹. CRC along with breast cancer screening is the only malignancies where screening is effective in the prevention or progression of disease. In 90% of cases, colorectal lesions are in the pre-cancerous (adenomatous) phase when patients are usually asymptomatic; therefore, screening can be highly effective.²¹

Screening in asymptomatic subjects reduces the incidence and mortality of CRC. Colonoscopy is the most commonly used method following a positive FIT for lesion detection and removal.¹³ However, it is not a cost-effective tool for screening as over 75% of the detected lesions are not CRC.²⁹

Uruguay and Argentina have implemented screening programs for CRC. Uruguay initiated a study to promote screening and healthy behavior in the general population and those most at risk (14,000 people)⁹ Uruguay has had a National Policy for the Detection of (CRC Program) since 1998. It includes the annual screening of asymptomatic individuals with ages ranging from 50 to 70 years using an immunochemical test (iFOBT). This recently changed to FIT as this more specific to the lower GI tract. Colonoscopy is then recommended for positive cases.^{9,10} The widespread use of FIT is vitally important in countries where cancer poses a major health problem.¹³

The compliance rates for FIT and colonoscopy in a feasibility study involving 11,734 asymptomatic individuals over the age of 50 were 90.1% and 75.1%, respectively. FIT was positive in 1170 (11.1%) responders and colonoscopy was performed in 879 (75.1%), revealing 330 (38%) lesions, including 54 advanced cancers, 47 early cancers, 131 advanced adenomas, and 98 low-risk non-advanced adenomas. Predictive values for cancer were 0.95 and 8.6% and for advanced adenomas 1.24 and 11.2%.¹⁴

In one Dutch study, 8806 (3.3%) of 265 881 participants had a positive FIT. Three thousand two hundred and fifty-four (1.2%) AN and 557 (0.2%) CRC were reported in colonoscopy. Prognostic models for detecting AN and CRC in round 3 were performed. The predicted risk ranged from 0.4% to 36.7% for AN and from 0.0% to 5.5% for CRC. In external validation, the model retained similar discrimination accuracy for AN (C-statistic 0.77, 95% CI 0.66 to 0.87) and CRC (C-statistic 0.78, 95% CI 0.66 to 0.91). The models accurately predict the risk of subsequent AN and CRC, discriminate those outcomes with a high degree of concordance, and allow for clinically meaningful risk stratification.³⁶

FIT and colonoscopy have been shown in randomized trials to reduce the incidence of CRC and its associated mortality. The incidence of advanced adenoma and CRC in preventive screening (Positive Fecal Tests) is depending on the number of people who undergo endoscopy. Even though, they have been showed to decrease CRC mortality but overestimate the rate of adenomas and cancer.³⁷

Our study has also shown that the population most at risk of malignancies is an economically active population (> 60 years) who are also more likely to have comorbidity. Thus, we must consider this in addition to the patient's individual risk and the odds based on FIT values.

Future risk stratification will require evaluating the current cutoff point (FIT > 100 ng) to enhance the

performance of the test, while not overwhelming the health-care system. The likelihood of detecting colorectal neoplasms was 25% before the test and 37.7% after the test (LR+ 5.7) in 63,393 subjects with average risk in Uruguay¹⁰ suggesting that deferring the colonoscopy would not be the most appropriate strategy. It is important to consider that the age group in which colonoscopy is indicated matches the age group with a higher potential risk of complications from COVID-19.

There are, however, some limitations of this study, with the main one being that the short and long-term patient outcomes are unknown for those who did not proceed to colonoscopy with a positive FIT. Furthermore, the results from this tertiary center may not be generalizable to smaller hospitals with different patient demographics as other risks influence mortality such as associated diseases such as cardiovascular disease or infection, population pyramid, and the prevalence of overweight or obese individuals.³⁷⁻³⁹

The main findings of the study are well-timed to guide service delivery during the COVID-19 pandemic, with a diagnostic yield of 400 ng Hb/mL demonstrating likely advanced adenomas or CRC. The indirect benefit of implementing such a cutoff is that it would reduce the burden of care and the risk of transmission of the virus during the epidemic in subjects at higher risk of comorbidity. Although lower cutoff levels would increase the detection rate of lesions, this would be at the expense of a higher number of colonoscopies. Prioritization of patients based on this cut-off point (400 ngHb/mL) in the challenging new setting for endoscopy in the fight against CRC, appears to be a feasible strategy that can lower the risks to both patients and healthcare workers.

Funding

The authors declare that no funding was received.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of human subjects and animals. The authors declare that no experiments on humans or animals were performed for this research.

Confidentiality, informed consent, and ethical approval. The authors have obtained approval from the Ethics Committee for the analysis of routinely

collected and anonymized clinical data; therefore, individual informed consent was not required. Relevant ethical recommendations have been followed.

Declaration on the use of artificial intelligence.

The authors declare that no generative artificial intelligence was used in the writing or creation of the content of this manuscript.

References

- Forman D, Sierra MS. Cancer in central and South America: introduction. *Cancer Epidemiol*. 2016;44 Suppl 1:S3-10.
- Bray F, Ferlay J, Soerjomataram I, Siegel R, Torre L, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2018;68:394-424.
- Maule M, Merletti F. Cancer transition and priorities for cancer control. *Lancet Oncol*. 2012;13:745-6.
- Ferlay J, Colombet M, Soerjomataram I, Mathers C, Parkin DM, Piñeros M, et al. Estimating the global cancer incidence and mortality in 2018: GLOBOCAN sources and methods. *Int J Cancer*. 2019;144(8):1941-53.
- Goss PE, Lee BL, Badovinac-Crnjevic T, Strasser-Weippl K, Chavarri-Guerra Y, Louis JS, et al. Planning cancer control in Latin America and the Caribbean. *Lancet Oncol*. 2013;14:391-436.
- Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380:2224-60.
- Centers for Disease Control and Prevention. Section 9: natural history and spectrum of disease. In: *Principles of Epidemiology in Public Health Practice, An Introduction to Applied Epidemiology and Biostatistics*. 3rd ed., Atlanta: Centers for Disease Control and Prevention; 2012.
- Programa Nacional de Prevención y Detección Temprana del Cáncer Colorectal. Argentina, Ministerio de Salud Resolución Ministerial 2173/2013; 2015. Available from: <https://www.msal.gov.ar/inc/index.php/programas/programanacionaldeprevencionydetecciontempranadelcancercolorectal>
- Programa Nacional de Control del Cáncer en el Uruguay, Período 2005-2009. Uruguay Ministerio de Salud Pública Dirección General de la Salud; 2008. Available from: https://www.bvsoncologia.org.uy/pdfs/detacados/pronaccan_2005-2010.pdf
- Fenocchi E, Martínez EL, Tolve J, Montano D, Rondan M, Parra-Blanco A, et al. Screening for colorectal cancer in Uruguay with an immunochemical faecal occult blood test. *Eur J Cancer Prev*. 2016;15:384-90.
- Rex DK, Boland CR, Dominitz JA, Giardiello FM, Johnson DA, Kaltenbach T, et al. Colorectal cancer screening: recommendations for physicians and patients from the U.S. Multi-society task force on colorectal cancer. *Gastrointest Endosc*. 2017;86:18-33.
- Levin B, Lieberman DA, MacFarland B, Smith RA, Brooks D, Andrews KS, et al. Screening and surveillance for the early detection of colorectal cancer and adenomatous polyps, 2008: a joint guideline from the American cancer society, the us multi-society task force on colorectal cancer, and the American college of radiology. *CA Cancer J Clin*. 2008;58:130-60.
- Brenner H, Chang-Claude J, Seiler CM, Rickert A, Hoffmeister M. Protection from colorectal cancer after colonoscopy: a population-based, case-control study. *Ann Intern Med*. 2011;154:22-30.
- Singh H, Nugent Z, Demers AD, Kliever EV, Mahmud SM, Bernstein CN. The reduction in colorectal cancer mortality after colonoscopy varies by site of the cancer. *Gastroenterology*. 2010;139:1128-37.
- Zauber AG, Winawer SJ, O'Brien MJ, Lansdorp-Vogelaar I, Van Ballegooijen M, Hankey BF, et al. Colonoscopic polypectomy and long-term prevention of colorectal-cancer deaths. *N Engl J Med*. 2012;366:687-96.
- Eishi Y. Fecal Occult Blood Tests and Their Pitfalls. Japan: JICA; 2000.
- Von Karsa L, Patnick J, Segnan N, Atkin W, Halloran S, Lansdorp-Vogelaar I. European guidelines for quality assurance in colorectal cancer screening and diagnosis: overview and introduction to the full supplement publication. *Endoscopy*. 2013;45:51-9.
- Miller AB. Implementation of colon cancer screening: techniques, costs, and barriers. *Gastroenterol Clin N Am*. 2008;37:83-95.
- Karsa LV, Lignini TA, Patnick J, Lambert R, Sauvaget C. The dimensions of the CRC problem. *Best Pract Res Clin Gastroenterol*. 2010;24:381-96.
- Ransohoff D, Lang C. Screening for colorectal cancer with the fecal occult blood test: a background paper. American college of physicians. *Ann Intern Med*. 1997;126:811-22.
- Nakama H, Zhang B, Fattah AS. A cost-effective analysis of the optimum number of stool specimens collected for immunochemical occult blood screening for colorectal cancer. *Eur J Cancer*. 2000;36:647-50.
- Allison JE, Feldman R, Tekawa IS. Hemoccult screening in detecting colorectal neoplasm: sensitivity, specificity, and predictive value. Long-term follow-up in a large group practice setting. *Ann Intern Med*. 1990;112:328-3.
- Morikawa T, Kato J, Yamaji Y, Wada Y, Mitsushima T, Shiratori Y. A comparison of the immunochemical fecal occult blood test and total colonoscopy in the asymptomatic population. *Gastroenterology*. 2005;129:422-8.
- Young GP, St John JB, Winawer SJ, Rozen P. Choice of fecal occult blood tests for colorectal cancer screening: recommendations based on performance characteristics in population studies: a WHO (World Health Organization) and OMED (World organization for digestive endoscopy) report. *Am J Gastroenterol*. 2002;97:2499-507.
- Smith A, Young GP, Cole SR, Bampton P. Comparison of a brush-sampling fecal immunochemical test for hemoglobin with a sensitive guaiac-based fecal occult blood test in detection of colorectal neoplasia. *Cancer*. 2006;107:2152-9.
- Levy Z, Rozen P, Hazazi R, Vilkin A, Waked A, Maoz E, et al. Can quantification of faecal occult blood predetermine the need for colonoscopy in patients at risk for non-syndromic familial colorectal cancer? *Aliment Pharmacol Ther*. 2006;24:1475-81.
- Lambert R, Provenzale D, Ectors N, Vainio H, Dixon MF, Atkin W, et al. Early diagnosis and prevention of sporadic colorectal cancer. *Endoscopy*. 2001;33:1042-64.
- Martínez ME, Baron JA, Lieberman DA, Schatzkin A, Lanza E, Winawer SJ, et al. A pooled analysis of advanced colorectal neoplasia diagnoses after colonoscopic polypectomy. *Gastroenterology*. 2009;136:832-41.
- Van Rossum LG, Van Rijn AF, Laheij RJ, Van Oijen MG, Fockens P, Jansen JB, et al. Cutoff value determines the performance of a semi-quantitative immunochemical faecal occult blood test in a colorectal cancer screening programme. *Br J Cancer*. 2009;101:1274-81.
- Nagorni A, Katia V, Zirkovic J. Advanced colorectal adenoma. *Arch Oncol*. 2004;12:59-60.
- Fenocchi F, Burgos B, Gaggero P, Sobrino-Cossio S, Lopez-Alvarenga JC, Lambert N. W1489: utility of faecal immunological test for diagnostic of colorectal neoplastic lesions. *Gastrointest Endosc*. 2010;71:AB341.
- Hernández V, Cubiella J, González-Mao MC, Iglesias F, Rivera C, Iglesias MB, et al. Fecal immunochemical test accuracy in average-risk colorectal cancer screening. *World J Gastroenterol*. 2014;28:20:1038-47.
- Yuan SY, Wu W, Fu J, Lang YX, Li JC, Guo Y, et al. Quantitative immunochemical fecal occult blood test for neoplasia in colon cancer screening. *J Dig Dis*. 2019;20:78-82.
- Shahidi N, Cheung WY. Colorectal cancer screening: opportunities to improve uptake, outcomes, and disparities. *World J Gastrointest Endosc*. 2016;8:733-40.
- Gralnek IM, Hassan C, Beilenhoff U, Antonelli G, Ebigbo A, Pellisé M, et al. ESGE and ESGENA position statement on gastrointestinal endoscopy and the COVID-19 pandemic. *Endoscopy*. 2020;52:483-90.
- Meester RG, Van de Schootbrugge-Vandermeer HJ, Breekveldt EC, De Jonge L, Toes-Zoutendijk E, Kooyker A, et al. Faecal occult blood loss accurately predicts future detection of colorectal cancer. A prognostic model. *Gut*. 2023;72:101-108.
- Helsing LM, Kalager M. Colorectal cancer screening-approach, evidence, and future directions. *NEJM Evid*. 2022;1:EVIDra2100035.
- Kaufmann R, Pontet-Ubal N. The challenge of tackling the obesity economic burden: the case of Uruguay [version 1; peer review: awaiting peer review]. *Emerald Open Res*. 2019;1:11.
- Lloyd-Sherlock P, Ebrahim S, Martínez R, McKee M, Ordunez P. Reducing the cardiovascular disease burden for people of all ages in the Americas region: analysis of mortality data, 2000-15. *Lancet Glob Health*. 2019;7:e604-12.

Risk factors associated with mortality from explosions in civilian and military populations

Factores de riesgo asociados a mortalidad por explosiones en poblaciones civil y militar

Patrizio Petrone^{1,2*}, Jordi Marin-García¹, Javier Dagnesses-Fonseca¹, John McNelis³, and Corrado P. Marini³

¹Department of Surgery, NYU Grossman Long Island School of Medicine, Mineola; ²Research Committee, Panamerican Trauma Society; Department of Surgery, ³Department of Surgery, Albert Einstein College of Medicine, Jacobi Medical Center. Bronx, New York, USA

Abstract

Trauma from explosions is among the most serious injuries faced by healthcare professionals. There are few studies that describe the risk factors for mortality, or the patterns of blast injuries in the civilian compared to the military population. Review of articles extracted from Medline, PubMed, and Scopus published (01/2014-03/2024). Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines were followed. Data on mortality and injury patterns were extracted from the articles that met the inclusion criteria to describe the type of injuries and to identify risk factors for mortality. Sixteen articles were included. Average age of civilians was 40.0 years $\pm$ 3.7, with men representing 79.4% compared to a mean age of 24.5 $\pm$ 1.4 and 79.0% for the military patients. While there was no difference in the distribution of injuries by body regions, the civilian population was more likely to sustain severe cranial and thoracic injuries (31.9% and 20.4% vs. 18.1% and 9.9%, respectively). Civilian population had a non-statistically significant lower risk of death (odds ratio = 0.96; 95% confidence interval = 0.82, 1.13). The findings of this review indicate that clinical experience derived from explosions occurring in one setting may not be fully generalizable to explosions occurring in other contexts, as demographic, environmental, and contextual factors contribute to distinct injury patterns.

Keywords: Explosions. Injury patterns. Injury severity. Blast injuries. Mass-casualty incidents. Mortality. Risk factors.

Resumen

Los traumatismos por explosiones se encuentran entre las lesiones más graves a las que se enfrentan los profesionales sanitarios. Existen pocos estudios que comparen los factores de riesgo de mortalidad o los patrones de lesiones por explosión en la población civil en comparación con la población militar. Se realizó una revisión sistemática de artículos publicados en Medline, PubMed y Scopus, de enero de 2014 a marzo de 2024, siguiendo las directrices PRISMA. Se extrajeron datos sobre mortalidad y tipos de lesiones de los estudios que cumplieron los criterios de inclusión. Se incluyeron 16 artículos. La edad media de los civiles fue de 40.0 $\pm$ 3.7 años, con un 79.4% de varones, frente a 24.5 $\pm$ 1.4 años y un 79.0% de varones en los militares. No hubo diferencias en la distribución de las lesiones por regiones corporales, pero la población civil presentó una mayor probabilidad de sufrir lesiones craneales y torácicas graves (31.9% y 20.4% frente a 18.1% y 9.9%, respectivamente). La población civil tuvo un riesgo de muerte menor, aunque no estadísticamente significativo (OR: 0.96; IC 95%: 0.82-1.13). Esta revisión sugiere que la experiencia clínica derivada de las explosiones en un entorno específico podría no ser completamente extrapolable a otros contextos, debido a diferencias demográficas, ambientales y operativas que influyen en los patrones de lesión.

Palabras clave: Explosiones. Patrones de lesiones. Gravedad de las lesiones. Lesiones por explosión. Incidentes con múltiples víctimas. Mortalidad. Factores de riesgo.

*Correspondence:

Patrizio Petrone

E-mail: patrizio.petrone@gmail.com

Date of reception: 21-11-2025

Date of acceptance: 14-02-2026

DOI: 10.24875/CIRUE.M26000953

Cir Cir (Eng). 2026;94(3):335-343

Contents available at PubMed

www.cirugiaycirujanos.com

2444-0507/© 2026 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Traumatic injuries remain one of the leading causes of morbidity and mortality in the world population.¹ Although trauma and acute care surgeons have experience with more common injury mechanisms, such as motor vehicle accidents and interpersonal violence, they are not as familiar with less frequent, but more lethal, blast-related injuries.^{2,3} Blast or explosion-related injuries encompass a wide spectrum of trauma resulting from explosive events and are typically classified by their setting in military (war-related) defined as those occurring in the context of organized armed conflict involving military forces, and civilian (non-war-related) contexts, as those occurring outside active combat zones, including terrorism, industrial accidents, and unintentional explosions; by injury mechanism (primary to quinary); and by the nature of the explosion as accidental blast – such as gas line ruptures, boiler failures, or chemical plant accidents – usually occur in industrial or residential settings and often involve thermal burns, blunt force trauma, and inhalation injuries from structural collapse or fire. In contrast, intentional explosions – such as those from improvised explosive devices (IEDs), car bombs, or suicide attacks – are engineered for maximum lethality, combining multiple mechanisms including blast overpressure, penetrating trauma from shrapnel, and complex polytrauma. Distinguishing between these events is essential not only due to their intent but also due to differing injury patterns: accidental explosions tend to injure bystanders, whereas intentional blasts are often premeditated to affect densely populated or enclosed areas, amplifying casualty severity. Recognizing these distinctions is critical for anticipating injury profiles and guiding effective emergency response and trauma care.

Explosions result in multisystem injuries through a range of mechanisms, including both direct and indirect blast effects. These effects generate shearing and tearing forces that can lead to tissue penetration, organ perforation or rupture, and thermal burns.⁴ In addition, blast waves may produce internal injuries that are often occult, difficult to detect due to their non-visible nature and the unique injury mechanisms involved.^{5,6} Explosive involve also has the potential to cause mass casualty incidents (MCI) given their non-discriminatory and spreading impact across wide areas.⁴ Victims located near the epicenter typically sustain immediate fatal or severe injuries, while those further from the blast radius are more likely to suffer minor to moderate trauma

(Fig. 1).⁷ As a general rule, for every fatality, at least two individuals sustain injuries.⁸

The global incidence of deaths due to armed conflict and terrorism has been rising, with an estimated 129,700 fatalities reported in 2017 alone.⁹ As blast injuries increasingly occur in civilian settings,^{10,11} there is a growing need for emergency personnel to be proficient in the assessment, management, and triage of blast-related trauma and potential MCIs.¹²⁻¹⁷ Nonetheless, accidental explosions remain a more frequent source of blast injuries among civilians. A notable example is the 2020 Beirut harbor explosion, in which the detonation of stored ammonium nitrate – located adjacent to a fireworks warehouse – resulted in over 200 deaths and tens of thousands of injuries.¹⁸

The distribution of injuries to the different body regions varies and is often directly related to the mechanism of the explosion and the location of the explosive material.¹⁹ The placement of IEDs at ground level as opposed to a higher level leads to a different distribution of body regions injuries. For example, in the 2013 Boston Marathon bombing, two IEDs were placed at ground level causing predominantly lower extremity injuries.¹⁶ Lung injuries and tympanic membrane perforations were the most common following the 2004 Madrid train bombings from over-pressurization in confined space.²⁰ In contrast, the Beirut port shock wave occurred in an open-air space, but quickly spread to the densely populated capital, where it was amplified by reflecting off surfaces. The pressure of the blast wave caused extensive damage to buildings and glass windows, resulting in blunt head injuries from falling debris and masonry, and penetrating wounds to the face and extremities.¹⁹

To date, relatively few studies have systematically investigated the risk factors associated with adult mortality following blast trauma, particularly with respect to differences in injury patterns between civilian and military populations. These distinctions may be critical in anticipating injury severity and guiding appropriate medical response. Within this context, the present study aims to examine and compare the injury profiles and associated mortality risk factors from both accidental and intentional explosions across civilian and military settings.

Methods

Data sources and search strategy

A literature search was performed using Medline, PubMed, Scopus, and through searching citations of




Figure 1. Type of injuries according to distance from the explosion.

other original articles. MeSH terms utilized were explosions, blast, blast waves, blast injuries, injury patterns, and mass-casualty incidents. Articles published in English and Spanish were selected from 1/1/2014-3/31/2024 (Fig. 2). The authors attempted to register this systematic literature review in PROSPERO, as it does not currently accept registrations for scoping reviews, literature reviews, or mapping reviews.

Study selection criteria

Only original and human subject research articles were included, encompassing both prospective and retrospective descriptive studies. Eligible studies were required to report injury profiles, defined as the distribution of injuries sustained from explosions stratified by injury severity score (ISS) and abbreviated injury score (AIS) body region. Exclusion criteria included studies involving non-human subjects, clinical case reports, literature reviews, pediatric populations, and publications in languages other than English or Spanish.

Data extraction

The first author (P. Petrone) supervised the entire process, from article selection to data extraction methodology. Data extraction was conducted in two phases: One international research fellow (Jordi Marin-García) independently extracted the first set of data, which was then verified by PP. In turn, PP extracted the second set of data, which was subsequently verified by JMG. For the articles published in Spanish, three of the first three authors are native Spanish speaker, ensuring accurate interpretation. Data were extracted using a standardized table that included the following

variables: Author, year of publication, study type, number of patients, age, gender, injury types, and injury severity (as measured by ISS and AIS). To ensure consistency and accuracy, any disagreement between the authors was resolved through a structured consensus process. Initial differences were identified during independent data review and addressed through open discussion. When informal consensus could not be achieved, a third senior investigator was consulted to provide an impartial opinion. This stepwise approach ensured methodological rigor, minimized bias, and maintained alignment with the study's predefined criteria and objectives.

This review was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-analyses and Assessing the methodological quality of systematic reviews guidelines to ensure methodological rigor and transparency. A comprehensive literature search was performed to answer the following PICO-formulated research question (Population, Intervention, Comparison, and Outcome):

- Population: Victims of explosion-related injuries (civilian and military).
- Intervention/exposure: Blast exposure in different operational contexts (war-related vs. non-war-related).
- Comparison: Injury pattern differences between groups.
- Outcome: Distribution and severity of injury by anatomical region.

An assessment of risk of bias was performed per individual study, using the Newcastle - Ottawa Quality Assessment Scale for both case control studies and cohort studies as valid evaluation tool.

Given the heterogeneity in study designs, populations, and outcome measures, data from the included studies were synthesized using a narrative synthesis approach. The involved systematic categorization of key themes, patterns, and findings across studies, with specific attention to contextual variables, such as setting, intent of explosion, and injury mechanisms, enabled a comprehensive synthesis of the available evidence and facilitated the identification of recurring trends, knowledge gaps, and clinically relevant associations.

To assess the certainty of the overall body of evidence, each outcome was evaluated using established domains: risk of bias, inconsistency, indirectness, imprecision, and publication bias. Certainty ratings were assigned as high, moderate, low, or very low, and any decision to downgrade the certainty was explicitly documented and justified to ensure transparency.




Figure 2. Preferred reporting items for systematic reviews and meta-analyses flowchart.

The AIS is an anatomically based injury severity scoring system that rates injury by body region on a six-point scale that provides data to calculate the ISS.^{21,22} There are six regions of the body according to the ISS classification to which injuries can be assigned, although the AIS dictionary 2005 – Update 2008 is divided into nine anatomical chapters.^{22,23}: head or neck injuries, facial injuries, thoracic injuries, abdominal injuries, pelvic, extremity, or pelvic girdle injuries, external injuries, and other traumatic injuries.

Statistical analysis

The data set was assessed for normality using the Shapiro-Wilk test, and continuous data were analyzed using Welch's t-test.^{24,25} The Welch's t-test was

selected instead of the Student's t-test because it provides better control for type I errors when homoscedasticity (the assumption that the variance of the data is homogeneous) is violated, while performing the Student's t-test when homoscedasticity is maintained.^{26,27} Discrete variables were compiled into contingency tables to determine the odds ratio (OR). Fisher's exact test was used instead of the chi-square for greater reliability.²⁸ Continuous variables are reported as mean with standard deviation (SD). Statistical significance was conferred if $p < 0.05$.

Results

Of the 10,028 articles initially identified through searches of Medline, PubMed, and Scopus, ten met the predefined

inclusion criteria. An additional six eligible articles were identified through citation tracking of original studies, resulting in a total of 16 articles included in the final review (Fig. 2). Duplicate publications were identified and removed using automated software (Covidence; Melbourne VIC 3000, Australia). All reviewers independently evaluated these 16 studies and agreed on their inclusion following a detailed assessment. Notably, only one of the included articles directly compared outcomes between civilian and military populations (Table 1).^{7,19,29-42}

Study demographics

Of the 16 studies included, 9 (56%) investigated the effects of explosions in the military populations, while the remaining seven focused on civilian populations. Across all studies, a total of 14,942 patients were affected by explosions – 9,240 civilians and 5,702 military personnel. Collectively, 27,270 injuries were reported: 17,620 in the civilian population resulting from both terrorist and accidental explosions, and 9,650 in the military population. Demographic characteristics of the study populations are summarized in table 2. The average age of victims in the civilian population was significantly higher than that of victims of military casualties (40 ± 3.7 vs. 24.5 ± 1.4 ; $p < 0.05$). However, there was no statistically significant difference in sex distribution, with a predominance of male victims in both groups (79.4% vs. 79.0%; $p > 0.05$). Mortality rates were also comparable between the two populations (4.6% vs. 4.7%; $p > 0.05$). No independent risk factors for mortality were identified that distinguished civilian from military outcomes (OR = 0.96, 95% confidence interval [CI] = 0.82, 1.13; $p = 0.659$), nor were there significant predictors for the occurrence of severe injuries, defined as ISS ≥ 15 (OR = 1, 95% CI = 0.82, 1.23; $p = 0.952$).

Injury profile and severity of injuries

The overall distribution of injuries among civilians did not differ significantly from that observed in the military population. Injuries involving the head–face–neck, thoracic, and abdominal regions occurred in 29.8% vs. 34%, 8.8% vs. 7.7%, and 7.3% vs. 6.5% of cases, respectively. However, extremity injuries were significantly more frequent among military casualties than among civilians (39.8% vs. 30.8%; $p < 0.05$). Conversely, head and thoracic injuries were more prevalent in the civilian population (31.9% vs. 18.1%

Table 1. Studies included by author and by type of population

Author (year of publication)	Study period	Type of population	n = 14,942
Rozenfeld et al. ²⁹	2004-2011	Military	474
Rozenfeld et al. ²	2012-2013	Military	19
Shakargy et al. ⁴⁶	2003-2014	Military	844
Heidenberg et al. ⁷	2000-2005	Civilian	1261
Yu et al. ³²	2015	Civilian	75
Sharrock et al. ³³	2003-2014	Military	80
Ashkenazi et al. ³⁴	1994-2005	Civilian	66
Rozenfeld et al. ³⁵	1997-2016	Civilian	1025
Schweizer et al. ³⁶	2001-2018	Military	739
Maddry et al. ³⁷	2011-2014	Military	915
Nunziato et al. ³⁸	2015	Civilian	2682
Al-Hajj et al. ³⁹	2020	Civilian	791
Tapia et al. ⁴⁰	2007-2020	Military	1526
Gebzan et al. ¹⁹	2020	Civilian	315
Rozenfeld et al. ⁴¹	1997-2018	Civilian- Military	3908
Shakargy et al. ⁴⁶	1982-2021	Military	222

Table 2. Demographics of the study population

Variable	Civilian events	Military events
Age	40 years (SD $\pm$ 3.65)	24.5 years (SD $\pm$ 1.41)
Sex, n (%)		
Male	7366 (79.4)	4505 (79.0)
Female	1742 (20.6)	1197 (21.0)
Mortality, n (%)	421 (4.6)	267 (4.7)
ISS	8 (IQR = 2)	9 (IQR = 10.5)

SD: standard deviation; ISS: injury severity score; IQR: interquartile range.

and 20.4% vs. 9.9%, respectively), while severe extremity injuries occurred more commonly in military personnel (51.8% vs. 30.9%). None of the included studies specified whether the explosive devices involved were high-order or low-order explosives.

Discussion

The objective of this review was to compare the risk factors associated with mortality among patients injured

by different types of explosions across civilian and military settings. Interestingly, the finding data suggests that, despite clear differences in the context and underlying causes of explosions between these populations, there are several similarities in injury patterns and outcomes. These commonalities may indicate the degree of transferability in treatment strategies and clinical management approaches across both settings.

As one would expect, the victims of war-related explosions were young adults with a mean age of 24.5 years, while civilian victims had a mean age of 40.0 years. The patterns of injuries in the civilian population showed a greater proportion of severe injuries to the head and thoracic body regions and a lower incidence of severe injuries to the pelvic and extremities compared to the military population. These differences can be ascribed to the types of explosive devices responsible for the explosions in the civilian and military environment and the universal use of protective equipment by the military personnel. IEDs and landmines, also known as antipersonnel mines, are among the most common causes of localized military explosions aimed at increasing the logistic burden for caring for the victims by causing either traumatic amputations of the lower extremities or massive lower extremities and pelvic injuries, as opposed to the civilian explosions that predispose the victims to head and thoracic injuries.^{41,43} Some degree of similarity between intentional and terrorism explosions is observed when high explosives are placed in maximum proximity to potential victims; in this setting, a massive amount of the explosive energy is transferred to victims.^{41,43}

The placement of an explosive device in enclosed environments, such as within a vehicle during assassination attempts, or in confined public spaces like buses, bars, or coffee shops during terrorist attacks, amplifies the impact of the blast by redirecting and concentrating the pressure wave toward victims inside the space.^{44,45} In contrast, unintentional outdoor explosions, like the one mentioned in Beirut, tend to produce a more heterogeneous injury pattern due to the dissipation of the blast wave in open air.²⁹

Explosions typically occur in one of four primary contexts: Domestic settings (e.g., gas leaks or residential fires), industrial environments involving occupational hazards (e.g., mining, demolition, or the production of chemical products and fuels), armed conflict zones, and acts of terrorism. An additional critical factor influencing the resulting injury patterns is the physical environment in which the explosion takes place, as factors such as confinement, proximity, and structural

surroundings significantly affect the blast's impact. Underwater blast waves cause greater damage due to the water's incompressibility.^{30,31} These waves travel faster and farther than in the air, leading to more severe injuries over longer distances.³² Personnel floating in water are more prone to abdominal than thoracic injuries, while fully submerged victims face equal risk of both, even at 3 times the blast distance.⁴⁶

Explosion-related injuries are classified into four categories based on the mechanism of injury (Table 3 and Fig. 3) and by distance from the blast epicenter (Fig. 1):

- Primary blast injuries result from the direct interaction of the blast shock wave with the body. Gas-containing organs, such as the middle ear, lungs, and gastrointestinal tract, are at particular risk. In contrast, solid organs, including the skin, are more resistant to the blast wave. Notably, patients with isolated primary blast injury may present with minimal signs of trauma despite significant internal damage.
- Secondary blast injuries are caused by energized fragments propelled by the blast. These include both *primary fragments* (components of the explosive device itself) and *secondary fragments* (environmental debris such as glass, metal, or building materials). These injuries often result in penetrating trauma, lacerations, traumatic amputations, and traumatic brain injury, and can affect any part of the body.
- Tertiary blast injuries occur when the victim is physically displaced by the blast wind and thrown against a hard surface or when heavy objects are propelled into the body. These injuries are commonly associated with blunt trauma, crush injuries, and structural collapse. The combination of primary and tertiary forces may lead to complex injuries such as limb avulsion, crush syndrome, and compartment syndrome.
- Quaternary blast injuries result from the thermal effects of the blast, combustion gases, and chemical or environmental contaminants. Victims may present with thermal burns, asphyxiation, and toxic syndromes secondary to fuel, metal exposure, or environmental contamination.

Limitations of the evidence and review process

Several limitations were identified in the available body of evidence. First, there was substantial heterogeneity

Table 3. Classification of trauma explosion

Mechanism	Cause	Organs affected	Typical injuries
Primary	Increased pressure	Predominates in organs with air: lungs, middle ear, gastrointestinal tract	Tympanic rupture Lung injuries Intestinal perforation TBI Eye injury
Secondary	Splinters/shrapnel	Whole body	Penetrating injuries Lacerations Compartment syndrome
Tertiary	The victim is thrown against a stationary object, or objects are thrown towards people	Whole body	Blunt trauma (contusions, fractures, crush injuries) Polytrauma Amputations Compartment syndrome Crush syndrome TBI
Quaternary	Other explosion-related injuries (e.g., heat)	Whole body Burns: unprotected skin (face, neck, hands)	Burns Inhalation Smoke poisoning Infections
Quinary	Bacteria, chemicals, radiologic materials	Hyperinflammatory or hypermetabolic responses	Shock, fever, coagulopathy without clear trauma mechanisms

TBI: traumatic brain injury.

across studies in terms of study design, populations examined, injury classification systems, and outcome measures, which limited direct comparability and precluded pooled analysis. Many of the included studies were retrospective or descriptive, often with small sample sizes and limited external validity. In addition, methodological details were frequently underreported, reducing the ability to assess study quality and risk of bias comprehensively.

Contextual factors – such as the type of explosive device used, the environmental setting of the incident (e.g., open vs. confined spaces), and the use of personal protective equipment – were inconsistently reported, further complicating interpretation of findings. Notably, within the civilian population, most studies did not differentiate between injuries resulting from accidental explosions and those related to acts of

terrorism, limiting the specificity of conclusions drawn for this subgroup.

Regarding the review process itself, several limitations must also be acknowledged. Although a comprehensive and systematic search strategy was employed, it is possible that relevant studies published in non-English languages or non-Spanish languages, as well as those in the grey literature, were inadvertently excluded. Screening and data extraction were conducted independently by the authors, with discrepancies resolved through consensus or third-party adjudication. Nonetheless, the potential for subjectivity in data interpretation remains and may have influenced the synthesis of findings. The authors emphasize that this review provides a descriptive synthesis rather than a meta-analytic comparison.




Figure 3. Type of explosion injuries.

Conclusions

The findings of this review indicate that clinical experience derived from explosions occurring in one setting – such as military or combat zones – may not be fully generalizable to explosions occurring in other contexts, such as civilian or accidental events, as demographic, environmental, and contextual factors contribute to distinct injury patterns.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare no conflicts of interest.

Ethical considerations

Protection of human subjects and animals. The authors declare that no experiments on humans or animals were performed for this research.

Confidentiality, informed consent, and ethical approval. This study does not involve personal patient data, medical records, or biological samples, and does not require ethical approval. SAGER guidelines do not apply.

Declaration on the use of artificial intelligence. The authors declare that no generative artificial intelligence was used in the writing or creation of the content of this manuscript.

References

1. Injuries and Violence. Available from: <https://www.who.int/news-room/fact-sheets/detail/injuries-and-violence> [Last accessed on 2024 Mar 13].

2. Rozenfeld M, Givon A, Peleg K. Violence-related versus terror-related stabbings: significant differences in injury characteristics. *Ann Surg.* 2018;267:965-70.
3. Almog G, Kedar A, Bala M. When a vehicle becomes a weapon: intentional vehicular assaults in Israel. *Scand J Trauma Resusc Emerg Med.* 2016;24:149.
4. DePalma RG, Burris DG, Champion HR, Hodgson MJ. Blast injuries. *N Engl J Med.* 2005;352:1335-42.
5. Champion HR, Holcomb JB, Young LA. Injuries from explosions: physics, biophysics, pathology, and required research focus. *J Trauma.* 2009;66:1468-77.
6. Almog G, Luria T, Richter E, Pizov R, Bdolah-Abram T, Mintz Y, et al. Can external signs of trauma guide management? Lessons learned from suicide bombing attacks in Israel. *Arch Surg.* 2005;140:390-3.
7. Heldenberg E, Givon A, Simon D, Bass A, Almog G, Peleg K, et al. Civilian casualties of terror-related explosions: the impact of vascular trauma on treatment and prognosis. *J Trauma Acute Care Surg.* 2016;81:435-40.
8. Coupland RM, Meddings DR. Mortality associated with use of weapons in armed conflicts, wartime atrocities, and civilian mass shootings: literature review. *BMJ.* 1999;319:407-10.
9. GBD 2017 Causes of Death Collaborators. Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet.* 2018;392:1736-88.
10. Hojman H, Rattan R, Osgood R, Yao M, Bugaev N. Securing the emergency department during terrorism incidents: lessons learned from the Boston marathon bombings. *Disaster Med Public Health Prep.* 2019;13:791-8.
11. De Ceballos JP, Turégano-Fuentes F, Perez-Diaz D, Sanz-Sanchez M, Martín-Llorente C, Guerrero-Sanz JE. 11 March 2004: the terrorist bomb explosions in Madrid, Spain - an analysis of the logistics, injuries sustained and clinical management of casualties treated at the closest hospital. *Crit Care.* 2005;9:104-11.
12. SALT mass casualty triage: concept endorsed by the American College of Emergency Physicians, American College of Surgeons Committee on Trauma, American Trauma Society, National Association of EMS Physicians, National Disaster Life Support Education Consortium, and State and Territorial Injury Prevention Directors Association. *Disaster Med Public Health Prep.* 2008;2:245-6.
13. Kashuk JL, Halperin P, Caspi G, Colwell C, Moore EE. Bomb explosions in acts of terrorism: evil creativity challenges our trauma systems. *J Am Coll Surg.* 2009;209:134-40.
14. Kilner TM, Brace SJ, Cooke MW, Stallard N, Bleetman A, Perkins GD. In 'big bang' major incidents do triage tools accurately predict clinical priority?: a systematic review of the literature. *Injury.* 2011;42:460-8.
15. Kapur GB, Hutson HR, Davis MA, Rice PL. The United States twenty-year experience with bombing incidents: implications for terrorism preparedness and medical response. *J Trauma.* 2005;59:1436-44.
16. Gates JD, Arabian S, Biddinger P, Blansfield J, Burke P, Chung S, et al. The initial response to the Boston marathon bombing: lessons learned to prepare for the next disaster. *Ann Surg.* 2014;260:960-6.
17. Jacobs LM, Wade DS, McSwain NE, Butler FK, Fabbri WP, Eastman AL, et al. The Hartford consensus: THREAT, a medical disaster preparedness concept. *J Am Coll Surg.* 2013;217:947-53.
18. Mansour HA, Bitar E, Fares Y, Makdessi AA, Maalouf A, Ghoul ME, et al. The Beirut port explosion: injury trends from a mass survey of emergency admissions. *Lancet Lond Engl.* 2021;398:21-2.
19. Gebran A, Abou Khalil E, El Moheb M, Albaini O, El Warea M, Ibrahim R, et al. The Beirut port explosion injuries and lessons learned: results of the Beirut blast assessment for surgical services (BASS) multicenter study. *Ann Surg.* 2022;275:398-405.
20. Gutierrez De Ceballos JP, Turégano Fuentes F, Perez Diaz D, Sanz Sanchez M, Martín Llorente C, Guerrero Sanz JE. Casualties treated at the closest hospital in the Madrid, March 11, terrorist bombings. *Crit Care Med.* 2005;33 Suppl 1:S107-12.
21. Gennarelli TA, Wodzin E. AIS 2005: a contemporary injury scale. *Injury.* 2006;37:1083-91.
22. Gennarelli TA, Wodzin E. Abbreviated Injury Scale 2005: Update 2008. Barrington, Ill.: Association for the Advancement of Automotive Medicine; 2008.
23. Baker SP, O'neil B, Haddon WJ Jr., Long WB. The injury severity score: a method for describing patients with multiple injuries and evaluating emergency care. *J Trauma.* 1974;14:187-96.
24. Shapiro SS, Wilk MB. An analysis of variance test for normality (complete samples). *Biometrika.* 1965;52:591-611.
25. Welch BL. The generalisation of student's problems when several different population variances are involved. *Biometrika.* 1947;34:28-35.
26. Overall JE, Atlas RS, Gibson JM. Tests that are robust against variance heterogeneity in k x 2 designs with unequal cell frequencies. *Psychol Rep.* 1995;76:1011-7.
27. Delacre M, Lakens D, Leys C. Why psychologists should by default use Welch's t-test instead of Student's t-test. *Int Rev Soc Psychol.* 2017;30:92-101.
28. Ramsey PH, Ramsey PP. Evaluating the normal approximation to the binomial test. *J Educ Stat.* 1988;13:173-82.
29. Rozenfeld M, Peleg K, Radomislensky I, Ragoler M, Israel Trauma Group (I.T.G.), Bodas M. Injuries from explosions: more differences than similarities between various types. *Disaster Med Public Health Prep.* 2023;17:e154.
30. Elsayed NM. Toxicology of blast overpressure. *Toxicology.* 1997;121:1-15.
31. Blast Injuries: Preparing for the Inevitable. Available from: <https://www.ebmedicine.net/topics/trauma/blast-injury> [Last accessed on 2024 Feb 04].
32. Yu M, Lv Q, Ding H, Zeng X, Cao J, Liu J, et al. Evaluation of blast injury patients from the 2015 Tianjin explosions in China. *Burns.* 2016;42:1133-40.
33. Sharrock AE, Remick KN, Midwinter MJ, Rickard RF. Combat vascular injury: influence of mechanism of injury on outcome. *Injury.* 2019;50:125-30.
34. Ashkenazi I, Sevi R, Turégano-Fuentes F, Walsh MS, Olsha O, Schecter WP, et al. Hemodynamic consequences of extremity injuries following a terrorist bombing attack: retrospective cohort study. *Eur J Trauma Emerg Surg.* 2019;45:865-70.
35. Rozenfeld M, Givon A, Rivkind A, Bala M, Peleg K, Alfici R, et al. New trends in terrorism-related injury mechanisms: is there a difference in injury severity? *Ann Emerg Med.* 2019;74:697-705.
36. Schweizer MA, Janak JC, Graham B, Mazuchowski EL, Gurney JM, Shackelford SA, et al. Nonfatal motor vehicle related injuries among deployed US Service members: characteristics, trends, and risks for limb amputations. *J Trauma Acute Care Surg.* 2019;87:907-14.
37. Maddy JK, Arana AA, Mora AG, Perez CA, Cutright JE, Kester BM, et al. Advancing prehospital combat casualty evacuation: patients amenable to aeromedical evacuation via unmanned aerial vehicles. *Mil Med.* 2021;186:e366-72.
38. Nunziato CA, Riley CJ, Johnson AE. How common are civilian blast injuries in the national trauma databank, and what are the most common mechanisms and characteristics of associated injuries? *Clin Orthop Relat Res.* 2021;479:683-91.
39. Al-Hajj S, Farran SH, Zgheib H, Tfaily MA, Halaoui A, Wehbe S, et al. The Beirut ammonium nitrate blast: a multicenter study to assess injury characteristics and outcomes. *J Trauma Acute Care Surg.* 2023;94:328-35.
40. Tapia G, Naylor JF, April MD, Schauer SG. Battle injury patterns sustained noncombatant military occupational specialty service members. *Med J (Ft Sam Houston Tex).* 2022;2:78-82.
41. Rozenfeld M, Givon A, Shenhar G, Renert L, Peleg K. A new paradigm of injuries from terrorist explosions as a function of explosion setting type. *Ann Surg.* 2016;263:1228-34.
42. Roy MC. Blast Injury: what to expect in civilian vs. Military contexts. *ICU Manag Pract.* 2015;15 (Accessed on Thu, 31 Dec 2015).
43. Peleg K, Jaffe DH, Israel Trauma Group. Are injuries from terror and war similar? A comparison study of civilians and soldiers. *Ann Surg.* 2010;252:363-9.
44. Noji EK, Lee CY, Davis T, Peleg K. Investigation of federal bureau of investigation bomb-related death and injury data in the United States between 1988 and 1997. *Mil Med.* 2005;170:595-8.
45. Kosashvili Y, Loebenberg MI, Lin G, Peleg K, Zvi F, Kluger Y, et al. Medical consequences of suicide bombing mass casualty incidents: the impact of explosion setting on injury patterns. *Injury.* 2009;40:698-702.
46. Shakargy JD, Gendler S, Talmi T, Shushan G, Radomislensky I, Tsur AM, et al. Blast injury patterns among Israel defense forces fatalities. *Mil Med.* 2023;188:e1788-94.

Boosting the findings of cellular indices and acute phase reactants as predictors of abscess after appendectomy

Impulsando los hallazgos de índices celulares y reactantes de fase aguda como predictores de absceso tras apendicectomía

Nelson A. Milanés-González^{1*}  and José E. Velázquez-Amador² 

¹Research Unit, Universidad de Colima, Colima, Colima; ²Research Unit, Universidad Autónoma de Guadalajara, Zapopan, Jalisco, Mexico

To the Editor,

I would like to congratulate you on the publication of the article by Moreno-Alfonso et al.¹ entitled “Cellular indices and acute-phase reactants as predictors of abscess after appendectomy,” which is undoubtedly relevant both to the management of the complications of appendicitis mentioned in the present case and to other surgical and nonsurgical conditions. This article reaffirms the importance of the complete blood count as the main initial paraclinical test, with very low cost and high benefit in the management of patients with appendicitis. As is well known, appendicitis has a high incidence in today’s world and is the leading cause of surgical emergency in both adults and children.² In the latter, it is even more relevant, since the approach is slowed down because the clinical history is mainly obtained indirectly and because clinical SIGNS may be masked. This leads to the presentation of complications such as perforation or abscess, before and after surgical intervention, as mentioned in the publication.

The literature had previously addressed the importance of acute-phase reactants as predictors of complications in different diseases; however, one of the highlights of the article is the comparison with three parameters that are emerging today, namely the neutrophil-to-lymphocyte ratio, the platelet-to-lymphocyte

ratio, and the direct bilirubin-to-lymphocyte ratio. The neutrophil-to-lymphocyte ratio had already demonstrated its usefulness in differentiating complicated from uncomplicated appendicitis.³ This contribution by the authors will undoubtedly promote the development of new studies to enrich the evidence currently available on these markers. To support this, we would like to make a couple of comments regarding the methodology and the results obtained in the study.

First, the selection of patients should be mentioned, including children younger than 15 years, which is to be expected, since this specific population is more susceptible to complications and, therefore, to greater morbidity and mortality; however, it would be interesting to apply their method in a different age group, such as older adults, who are also considered at risk.

Second, in the method section, the treatment given to patients in a homogeneous manner, based on cefotaxime and metronidazole, is mentioned. This is also a standard measure in many patients with this disease, but it would be useful to mention whether there is any literature on similar results using another antibiotic regimen.

Third, as part of the appendectomy approach, a minimally invasive technique is mentioned: video-assisted transumbilical appendectomy, which has demonstrated, per se, a lower incidence of trans- and

*Correspondence:

Nelson A. Milanés-González

E-mail: nelsonantonio971@gmail.com

2444-0507/© 2026 Academia Mexicana de Cirugía. Published by Permanyer. This is an open access article under the terms of the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Date of reception: 04-02-2026

Date of acceptance: 28-02-2026

DOI: 10.24875/CIRUE.M26000954

Cir Cir (Eng). 2026;94(3):344-345

Contents available at PubMed

www.cirugiaycirujanos.com




Figure 1. Representative pyramid of the findings presented by the authors regarding the main objective of the article. OR, odds ratio.

postoperative complications compared with the usual laparoscopic and laparotomy techniques. Although the authors mention that there were no significant differences regarding the occurrence of abscess after appendectomy, other publications did report a difference, which could indicate selection bias.⁴

Finally, we would like to highlight the results reported by the authors with a figure of our own creation (Fig. 1).

We can only congratulate the authors on their successful and interesting publication, which will be considered for future studies in the field of surgery. We look forward to their response.

Funding

The authors declare that they have not received funding.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Ethical considerations

Protection of human subjects and animals. The authors declare that no experiments were performed on humans or animals for this research.

Confidentiality, informed consent, and ethical approval. The study does not involve personal data, medical records, or human biological samples; therefore, ethical approval is not required. The SAGER guidelines do not apply.

Declaration on the use of artificial intelligence. The authors declare that no type of generative artificial intelligence was used for the writing or creation of content in this manuscript.

References

1. Moreno-Alfonso JC, Lozano-Eslava LC, Barbosa-Velázquez S, Molina-Caballero A, Yáñez-Irazábal MC, Pérez-Martínez A. Cellular indices and acute phase reactants as predictors of abscess after appendectomy. *Cir Cir.* 2025;93:652-8.
2. Borruel Nacenta S, Ibáñez Sanz L, Sanz Lucas R, Depetris MA, Martínez Chamorro E. Update on acute appendicitis: typical and atypical findings. *Radiologia (Engl Ed).* 2023;65(Suppl 1):S81-91.
3. Chandni, Wasay A, Chaudhary MA, Qureshi MU. Preoperative neutrophil-to-lymphocyte ratio as a predictive marker of postoperative infectious complications in pediatric patients with acute appendicitis. *Cureus.* 2024;16:e71353.
4. Koontz CS, Smith LA, Burkholder HC, Higdon K, Aderhold R, Carr M. Video-assisted transumbilical appendectomy in children. *J Pediatr Surg.* 2006;41:710-2.